

**EMBEDDINGS OF FRECHET SPACES
IN UNIFORM FRECHET ALGEBRAS**

Jorge Mujica

RELATÓRIO TÉCNICO Nº 32/90

Abstract. It is shown that every Fréchet space is topologically isomorphic to a complemented subspace of a uniform Fréchet algebra.

Universidade Estadual de Campinas
Instituto de Matemática, Estatística e Ciência da Computação
IMECC - UNICAMP
Caixa Postal 6065
13.081 - Campinas - SP
BRASIL

O conteúdo do presente Relatório Técnico é de única responsabilidade do autor.

Setembro - 1990

EMBEDDINGS OF FRECHET SPACES IN UNIFORM FRECHET ALGEBRAS

by Jorge Mujica

Abstract

It is shown that every Fréchet space is topologically isomorphic to a complemented subspace of a uniform Fréchet algebra.

Introduction

A well known result of Milne [10] asserts that every Banach space is isometrically isomorphic to a complemented subspace of a uniform Banach algebra. Since Enflo [6] has constructed a Banach space without the approximation property, Milne's theorem implies the existence of a uniform Banach algebra without the approximation property.

In this note we first present Milne's proof from the point of view of infinite dimensional holomorphy, and next follow the same approach to extend Milne's theorem to the realm of Fréchet spaces.

I am grateful to Aleksander Pelczynski for observing that Milne's theorem is closely connected with a result in [13]. That observation ultimately led to the proof of Milne's theorem presented here. I am also grateful to João B. Prolla for some helpful comments.

1. Notation and Terminology

Throughout this note $C(X)$ denotes the locally multiplicatively con-

vex algebra of all complex-valued continuous functions on a topological space X , with the compact-open topology. A uniform Banach algebra (resp. uniform Fréchet algebra) is a closed subalgebra of $C(X)$ which contains the constants and separates the points of X , for a compact Hausdorff space X (resp. hemicompact k -space X).

If U is an open subset of a complex locally convex space E , then $H(U)$ denotes the locally multiplicatively convex algebra of all complex-valued holomorphic functions on U , with the compact-open topology, whereas $H^\infty(U)$ denotes the Banach algebra of all bounded members of $H(U)$, with the sup norm.

If E is a real topological vector space, then the Cartesian product $E \times E$, with the operations

$$(x, y) + (s, t) = (x + s, y + t),$$

$$(\alpha + \beta i)(x, y) = (\alpha x - \beta y, \alpha y + \beta x),$$

is a complex topological vector space, called the complexification of E , and denoted by $E_{\mathbb{C}}$. If E is a real Banach space, then $E_{\mathbb{C}}$ is a complex Banach space for the norm

$$\|(x, y)\| = \sup_{\theta} \max\{\|x \cos \theta - y \sin \theta\|, \|y \cos \theta + x \sin \theta\|\}.$$

Furthermore, the mapping $x \rightarrow (x, 0)$ embeds E isometrically into $E_{\mathbb{C}}$, and the projection $(x, y) \rightarrow (x, 0)$ has norm one.

We refer to [9] of [14] for the terminology from topological vector spaces, to [7] for the terminology from topological algebras, and to [5] or [12] for the terminology from infinite dimensional complex analysis.

2. Embeddings of Banach Spaces in Uniform Banach Algebras

2.1 Milne's Theorem [10]. Every Banach space is isometrically isomorphic to a 1-complemented subspace of a uniform Banach algebra.

Proof. Consider first the case of a complex Banach space E . If $U_{E'}$ denotes the open unit ball of the dual E' , then the bidual E'' is isometrically isomorphic to a 1-complemented subspace of $H^\infty(U_{E'})$. Indeed, if $S : E'' \hookrightarrow H^\infty(U_{E'})$ is the inclusion mapping, and $T : H^\infty(U_{E'}) \rightarrow E''$ is defined by $Tf = Df(0)$ for every $f \in H^\infty(U_{E'})$, then $T \circ S(x'') = x''$ for every $x'' \in E''$, S is an isometry and $\|T\| = 1$. If A is the subalgebra of $H^\infty(U_{E'})$ generated by $S(E)$, then $Tf \in E$ for every $f \in A$. Indeed, each $f \in A$ can be written as a sum $f = \sum_{m=0}^N f_m$, where

$$f_m(x') = \sum_{j=1}^{j_m} c_{mj}(x'(x_{mj}))^m,$$

with $c_{mj} \in \mathbb{C}$ and $x_{mj} \in E$. Since $Df(0) = f_1$, it follows that $Tf \in E$, as asserted. Whence it follows that $Tf \in E$ for every $f \in \overline{A}$, the closure of A in $H^\infty(U_{E'})$. Thus we have a commutative diagram

$$\begin{array}{ccccc} E'' & \xrightarrow{S} & H^\infty(U_{E'}) & \xrightarrow{T} & E'' \\ \uparrow & & \uparrow & & \uparrow \\ E & \hookrightarrow & \overline{A} & \longrightarrow & E \end{array}$$

and E is isometrically isomorphic to a 1-complemented subspace of \overline{A} . Since \overline{A} may be regarded also as a closed subalgebra of $C(\overline{U}_{E'}, \sigma(E', E))$, the proof is complete in the case of a complex Banach space E .

Finally, if E is a real Banach space, then E is isometrically isomorphic to 1-complemented subspace of its complexification $E_{\mathbb{C}}$. Thus the desired conclusion follows from the first part of the proof.

3. Embeddings of Fréchet Spaces in Uniform Fréchet Algebras

3.1 Theorem. Every Fréchet space E is topologically isomorphic to a complemented subspace of a uniform Fréchet algebra A . If E is, in addition, Montel, Schwartz or nuclear, then we can take an A with the same property.

Proof. Consider first the case of a complex Fréchet space E , and let E'_c denote the dual of E , with the topology of uniform convergence on the compact subsets of E . Then E'_c is a hemicompact k -space (see [11]), and $H(E'_c)$ is a uniform Fréchet algebra. Since $(E'_c)'_c = E$, the method of proof of Theorem 2.1 shows that E is topologically isomorphic to a complemented subspace of $H(E'_c)$ (see [1] or [5]). If E is a Fréchet–Montel space, then $H(E'_c)$ is a Fréchet–Montel space as well, by a result of Dineen [4]. If E is a Fréchet–Schwartz space, then $H(E'_c)$ is a Fréchet–Schwartz space as well, by a result of Colombeau and Perrot [3]. And if E is a Fréchet–nuclear space, then $H(E'_c)$ is a Fréchet–nuclear space as well, by a result of Boland [2]. This completes the proof in the case of a complex Fréchet space E .

Finally, if E is a real Fréchet space, then E is topologically isomorphic to a complemented subspace of its complexification E_C . Thus the desired conclusions follow easily from the first part of the proof.

Since Hogbe–Nlend [8] has constructed a Fréchet–Schwartz space without the approximation property, Theorem 3.1 implies the existence of a uniform Fréchet–Schwartz algebra without the approximation property.

References

- [1] R. ARON and M. SCHOTTENLOHER, Compact holomorphic mappings on Banach spaces and the approximation property. *J. Funct. Anal.* 21 (1976), 7–30.
- [2] P. BOLAND, Holomorphic functions on nuclear spaces. *Trans. Amer. Math. Soc.* 209 (1975), 275–281.
- [3] J. F. COLOMBEAU and B. PERROT, Reflexivity and kernels in infinite dimensional holomorphy. *Portugal. Math.* 36(1977), 291–300.
- [4] S. DINEEN, Holomorphic functions on strong duals of Fréchet–Montel spaces. In: *Infinite Dimensional Holomorphy and Applications*, pp. 147–166. *North–Holland Mathematics Studies*, vol. 12. North–Holland, Amsterdam, 1977.
- [5] S. DINEEN, *Complex Analysis in Locally Convex Spaces*. North–Holland Mathematics Studies, vol. 57. North–Holland, Amsterdam 1981.
- [6] P. ENFLO, A counterexample to the approximation property in Banach spaces. *Acta Math.* 130 (1973), 309–317.
- [7] H. GOLDMANN, *Uniform Fréchet algebras*. North–Holland Mathematics Studies, vol. 162. North–Holland, Amsterdam, 1990.
- [8] H. HOGBE–NLEND, Les espaces de Fréchet–Schwartz et la propriété d’approximation. *C. R. Acad. Sci. Paris* 275 (1972), 1073–1075.
- [9] J. HORVATH, *Topological Vector Spaces and Distributions*, vol. I, Addison–Wesley, Reading, Massachusetts, 1966.
- [10] H. MILNE, Banach space properties of uniform algebras. *Bull. London Math. Soc.* 4(1972), 323–326.
- [11] J. MUJICA, Domains of holomorphy in (DFC)–spaces. In: *Functional Analysis, Holomorphy and Approximation Theory*, pp. 500–533. *Lecture Notes in Mathematics*, vol. 843. Springer, Berlin, 1981.

- [12] J. MUJICA, Complex Analysis in Banach Spaces. North-Holland Mathematics Studies, vol. 120. North-Holland, Amsterdam, 1986.
- [13] J. MUJICA, Linearization of bounded holomorphic mappings on Banach spaces. Trans. Amer. Math. Soc., to appear.
- [14] H. SCHAEFER, Topological Vector Spaces. Graduate Texts in Mathematics, vol. 3. Springer, New York, 1971.

Instituto de Matemática
 Universidade Estadual de Campinas
 Caixa Postal 6065
 13081 Campinas, SP, Brazil

RELATÓRIOS TÉCNICOS — 1990

- 01/90 Harmonic Maps Into Periodic Flag Manifolds and Into Loop Groups — Caio J. C. Negreiros.
- 02/90 On Jacobi Expansions — E. Capelas de Oliveira.
- 03/90 On a Superlinear Sturm–Liouville Equation and a Related Bouncing Problem — D. G. Figueiredo and B. Ruf.
- 04/90 F -Quotients and Envelope of F -Holomorphy — Luiza A. Moraes, Otília W. Paques and M. Carmelina F. Zaine.
- 05/90 S -Rationally Convex Domains and The Approximation of Silva-Holomorphic Functions by S -Rational Functions — Otília W. Paques and M. Carmelina F. Zaine.
- 06/90 Linearization of Holomorphic Mappings On Locally Convex Spaces — Jorge Mujica and Leopoldo Nachbin.
- 07/90 On Kummer Expansions — E. Capelas de Oliveira.
- 08/90 On the Convergence of SOR and JOR Type Methods for Convex Linear Complementarity Problems — Alvaro R. De Pierro and Alfredo N. Iusem.
- 09/90 A Curvilinear Search Using Tridiagonal Secant Updates for Unconstrained Optimization — J. E. Dennis Jr., N. Echebest, M. T. Guardarucci, J. M. Martínez, H. D. Scolnik and C. Vacchino.
- 10/90 The Hypebolic Model of the Mean \times Standard Deviation “Plane” — Sueli I. R. Costa and Sandra A. Santos.
- 11/90 A Condition for Positivity of Curvature — A. Derdzinski and A. Rigas.
- 12/90 On Generating Functions — E. Capelas de Oliveira.
- 13/90 An Introduction to the Conceptual Difficulties in the Foundations of Quantum Mechanics a Personal View — V. Buonomano.
- 14/90 Quasi-Invariance of product measures Under Lie Group Perturbations: Fisher Information And L^2 -Differentiability — Mauro S. de F. Marques and Luiz San Martin.
- 15/90 On Cyclic Quartic Extensions with Normal Basis — Miguel Ferrero, Antonio Paques and Andrzej Solecki.
- 16/90 Semilinear Elliptic Equations with the Primitive of the Nonlinearity Away from the Spectrum — Djairo G. de Figueiredo and Olimpio H. Miyagaki.
- 17/90 On a Conjugate Orbit of G_2 — Lucas M. Chaves and A. Rigas.
- 18/90 Convergence Properties of Iterative Methods for Symmetric Positive Semidefinite Linear Complementarity Problems — Álvaro R. de Pierro and Alfredo N. Iusem.

- 19/90 The Status of the Principle of Relativity — *W. A. Rodrigues Jr. and Q. A. Gomes de Souza.*
- 20/90 Geração de Gerenciadores de Sistemas Reativos — *Antonio G. Figueiredo Filho e Hans K. E. Liesenberg.*
- 21/90 Um Modelo Linear Geral Multivariado Não-Paramétrico — *Belmer Garcia Negrillo.*
- 22/90 A Method to Solve Matricial Equations of the Type $\sum_{i=1}^p A_i X B_i = C$ — *Vera Lúcia Rocha Lopes and José Vitório Zago.*
- 23/90 \mathbb{Z}_2 -Fixed Sets of Stationary Point Free \mathbb{Z}_4 -Actions — *Claudina Izepe Rodrigues.*
- 24/90 The m -Ordered Real Free Pro-2-Group Cohomological Characterizations — *Antonio José Engler.*
- 25/90 On Open Arrays and Variable Number of Parameters — *Claudio Sergio Da Rós de Carvalho and Tomasz Kowaltowski.*
- 26/90 Bordism Ring of Complex Involutions — *J. Carlos S. Kiihl.*
- 27/90 Approximation of Continuous Convex-Cone-Valued Functions by Monotone Operators — *João B. Prolla.*
- 28/90 On Complete Digraphs Which Are Associated to Spheres — *Davide C. Demaria and J. Carlos S. Kiihl.*
- 29/90 Deriving Ampère's Law from Weber's Law — *A. K. T. Assis.*
- 30/90 Fatores Não Paramétricos para Experimentos Completamente Casualizados para Três Fatores, com Interação — *Belmer Garcia Negrillo.*
- 31/90 On the Velocity which appears in Lorentz Force Law: An Illuminating Puzzle — *A. K. T. Assis.*