

LINEARIZATION OF HOLOMORPHIC
MAPPINGS ON INFINITE DIMENSIONAL SPACES

87	Matemática
JEPEN	Matemática

**LINEARIZATION OF HOLOMORPHIC
MAPPINGS ON INFINITE
DIMENSIONAL SPACES**

Jorge Mujica

RELATÓRIO TÉCNICO Nº 54/91

Abstract. After giving a survey of the main results concerning linearization of bounded holomorphic mappings on Banach spaces, these results are applied to the study of interpolating sequences. The connection with the classical Hardy spaces, in the case of the open unit disc, is established.

Universidade Estadual de Campinas
Instituto de Matemática, Estatística e Ciência da Computação
IMECC - UNICAMP
Caixa Postal 6065
13.081 - Campinas - SP
BRASIL

O conteúdo do presente Relatório Técnico é de única responsabilidade do autor.

Outubro - 1991

LINEARIZATION OF HOLOMORPHIC MAPPINGS ON INFINITE DIMENSIONAL SPACES

Jorge Mujica

Abstract. After giving a survey of the main results concerning linearization of bounded holomorphic mappings on Banach spaces, these results are applied to the study of interpolating sequences. The connection with the classical Hardy spaces, in the case of the open unit disc, is established.

Introduction

Several authors have obtained linearization theorems for various classes of holomorphic mappings. It seems that the first general result of this kind is due to Mazet [6], who obtained a linearization theorem for holomorphic mappings on locally convex spaces, thus improving previous results of Schottenloher [13] and Ryan [12]. In a recent paper Nachbin and the author [10] gave a new proof and several applications of the Mazet linearization theorem. Very recently Dineen, Galindo, Garcia and Maestre [1] solved two of the problems left open in [10].

By specializing to smaller classes of mappings, the author [8] obtained a linearization theorem for bounded holomorphic mappings, whereas Galindo, Garcia and Maestre [2] obtained a linearization theorem for holomorphic mappings of bounded type. In a very recent paper the author [9] showed that the last two classes of mappings are intimately connected.

In this lecture we begin with a survey of the main results concerning linearization of bounded holomorphic mappings on Banach spaces, taken mainly from [8]. We next apply these results to the study of interpolating sequences. At the end we restrict our attention to the case of the open unit disc and show the connection of the preceding results with the classical

Hardy spaces.

I wish to thank the organizers of the X Latin American School of Mathematics for their kind invitation and financial support to attend and deliver this lecture.

1. Notation and Terminology

The letters E and F always represent complex Banach spaces, and $L(E; F)$ denotes the vector space of all continuous linear operators from E into F . Unless stated otherwise, $L(E; F)$ is endowed with its natural norm topology. We write E' instead of $L(E; \mathbb{C})$ for the dual of E .

Let U be a nonvoid open subset of E . A mapping $f : U \rightarrow F$ is said to be *holomorphic* if f is continuous and the function $\lambda \rightarrow \psi \circ f(a + \lambda b)$ is holomorphic on an open neighborhood of the origin in \mathbb{C} for every $a \in U, b \in F$ and $\psi \in F'$. Let $H(U; F)$ denote the vector space of all holomorphic mappings from U into F , and let $H^\infty(U; F)$ denote the subspace of all bounded members of $H(U; F)$. Unless stated otherwise, $H^\infty(U; F)$ is equipped with the sup norm topology. When $F = \mathbb{C}$ we write $H(U)$ instead of $H(U; \mathbb{C})$, and $H^\infty(U)$ instead of $H^\infty(U; \mathbb{C})$. We refer to the author's book [7] for the properties of holomorphic mappings on Banach spaces.

2. Linearization of Bounded Holomorphic Mappings

The following linearization theorem is taken from [8].

2.1. Theorem [8]. *Let U be an open subset of a Banach space E . Then there are a Banach space $G^\infty(U)$ and a mapping $\delta_U \in H^\infty(U; G^\infty(U))$ with the following universal property: For each Banach space F and each mapping $f \in H^\infty(U; F)$, there is a unique operator $T_f \in L(G^\infty(U); F)$ such that $T_f \circ \delta_U = f$. The mapping*

$$f \in H^\infty(U; F) \rightarrow T_f \in L(G^\infty(U); F)$$

is an isometric isomorphism. These properties characterize $G^\infty(U)$ uniquely up to an isometric isomorphism.

The following proposition shows the connection between E and $G^\infty(U)$.

2.2. Proposition [8]. *Let U be a bounded open subset of a Banach space E . Then E is topologically isomorphic to a complemented subspace of $G^\infty(U)$.*

The following proposition shows the connection between properties of a mapping $f \in H^\infty(U; F)$ and properties of the corresponding operator $T_f \in L(G^\infty(U); F)$.

2.3. Proposition [8]. *Let E and F be Banach spaces, and let U be an open subset of E .*

(a) *The range of a mapping $f \in H^\infty(U; F)$ is contained in a finite dimensional subspace of F if and only if the corresponding operator $T_f \in L(G^\infty(U); F)$ has finite rank.*

(b) *The range of a mapping $f \in H^\infty(U; F)$ is relatively compact (resp. relatively weakly compact) if and only if the corresponding operator $T_f \in L(G^\infty(U); F)$ is compact (resp. weakly compact).*

We denote by $H_K^\infty(U; F)$ (resp. $H_{wK}^\infty(U; F)$) the subspace of all $f \in H^\infty(U; F)$ which have a relatively compact range (resp. relatively weakly compact range).

3. Another Topology on $H^\infty(U; F)$

Theorem 2.1 tells us that the mapping

$$f \in H^\infty(U; F) \rightarrow T_f \in L(G^\infty(U); F)$$

is an isometric isomorphism. If τ_c denotes the compact-open topology, then it is very useful to consider the unique locally convex topology τ_γ on $H^\infty(U; F)$ such that the mapping

$$f \in (H^\infty(U; F), \tau_\gamma) \rightarrow T_f \in (L(G^\infty(U); F), \tau_c)$$

is a topological isomorphism. The following theorem furnishes an explicit description of τ_γ in terms of seminorms.

3.1. Theorem [8]. Let E and F be Banach spaces, and let U be an open subset of E . Then τ_γ is the locally convex topology on $H^\infty(U; F)$ generated by all the seminorms of the form

$$p(f) = \sup_j \alpha_j \|f(x_j)\|,$$

with $(x_j) \subset U$ and $(\alpha_j) \in c_0, \alpha_j > 0$.

With the aid of Theorem 3.1 we can describe $G^\infty(U)$ as follows.

3.2. Theorem [8][9]. Let U be an open subset of a Banach space E . Then $G^\infty(U)$ consists of all linear functionals $u \in H^\infty(U)'$ of the form

$$u = \sum_{j=1}^{\infty} \beta_j \delta_{x_j},$$

with $(x_j) \subset U$ and $(\beta_j) \in l^1$. Moreover,

$$\|u\| = \inf \sum_{j=1}^{\infty} |\beta_j|,$$

where the infimum is taken over all such representations of u .

4. The Approximation Property

We recall that a Banach space E is said to have the *approximation property*, introduced by Grothendieck [4], if the identity operator on E lies in the τ_c -closure of $E' \otimes E$ in $L(E; E)$. The following results give several characterizations of the approximation property in terms of bounded holomorphic mappings.

4.1. Theorem [8]. Let U be a balanced, bounded, open subset of a Banach space E . Then the following conditions are equivalent:

- (a) E has the approximation property.
- (b) For each Banach space F , $H^\infty(U) \otimes F$ is τ_γ -dense in $H^\infty(U; F)$.
- (c) The inclusion mapping $U \hookrightarrow E$ lies in the τ_γ -closure of $H^\infty(U) \otimes E$ in $H^\infty(U; E)$.

(d) The mapping δ_U lies in the τ_γ -closure of $H^\infty(U) \otimes G^\infty(U)$ in $H^\infty(U; G^\infty(U))$.

(e) $G^\infty(U)$ has the approximation property.

(f) For each Banach space F and each open set $V \subset F$, $H^\infty(V) \otimes E$ is τ_γ -dense in $H^\infty(V; E)$.

(g) For each Banach space F and each open set $V \subset F$, $H^\infty(V) \otimes E$ is norm-dense in $H_K^\infty(V; E)$.

4.2. Proposition [8]. *Let U be an open subset of a Banach space E . Then $H^\infty(U)$ has the approximation property if and only if, for each Banach space F , $H^\infty(U) \otimes F$ is norm-dense in $H_K^\infty(U; F)$.*

Since it is still unknown whether $H^\infty(\Delta)$ has the approximation property, where Δ denotes the open unit disc, Proposition 4.2 may be of some use in this connection.

5. Interpolating Sequences

Let $B(X)$ denote the Banach algebra of all bounded complex-valued functions on a nonvoid set X , with the sup norm. We will need the following result from the book of Garnett [3, Theorem VII.2.2].

5.1. Theorem [3]. *Let A be a subalgebra of $B(X)$ which contains the constants and separates the points of X . Let x_1, \dots, x_n be distinct points of X and let*

$$M = \sup_{\|f\|_\infty \leq 1} \inf\{\|f\| : f \in A, f(x_k) = \eta_k \text{ for } k = 1, \dots, n\}.$$

Then for each $\varepsilon > 0$ there are $f_1, \dots, f_n \in A$ such that $f_j(x_k) = \delta_{jk}$ for $j, k = 1, \dots, n$ and

$$\sup_{x \in X} \sum_{j=1}^n |f_j(x)| \leq M^2 + \varepsilon.$$

Let U be an open subset of a Banach space E . A sequence $(x_k) \subset U$ is said to be an *interpolating sequence* if the mapping $f \in H^\infty(U) \rightarrow (f(x_k)) \in l^\infty$ is surjective. The following theorem gives several characterizations of in-

terpolating sequences.

5.2. Theorem. *Let U be an open subset of a Banach space E . For a sequence $(x_k) \subset U$ the following conditions are equivalent:*

(a) *The mapping*

$$Q : f \in H^\infty(U) \rightarrow (f(x_k)) \in l^\infty$$

is surjective.

(b) *There exists $P \in L(l^\infty; H^\infty(U))$ such that $Q \circ P\eta = \eta$ for every $\eta \in l^\infty$.*

(c) *There exists $P \in L(c_0; H^\infty(U))$ such that $Q \circ P\eta = \eta$ for every $\eta \in c_0$.*

(d) *The mapping*

$$S : (\xi_k) \in l^1 \rightarrow \sum_{k=1}^{\infty} \xi_k \delta_{x_k} \in G^\infty(U)$$

is an embedding.

(e) *There exists $T \in L(G^\infty(U); l^1)$ such that $T \circ S\xi = \xi$ for every $\xi \in l^1$.*

(f) *There exists $\varphi \in H^\infty(U; l^1)$ such that $\varphi(x_k) = e_k$ for every $k \in \mathbb{N}$.*

(g) *For each Banach space F , the mapping*

$$Q_F : f \in H^\infty(U; F) \rightarrow (f(x_k)) \in l^\infty(F)$$

is surjective.

(h) *For each Banach space F , there exists $P_F \in L(l^\infty(F); H^\infty(U; F))$ such that $Q_F \circ P_F y = y$ for every $y \in l^\infty(F)$.*

Proof. (a) \Rightarrow (f): By the open mapping theorem, Q is open and therefore induces a topological isomorphism

$$R : H^\infty(U)/\text{Ker}Q \rightarrow l^\infty.$$

If $M = \|R^{-1}\|$, then

$$M = \sup_{\|\eta\|_\infty \leq 1} \inf\{\|f\| : f \in H^\infty(U), f(x_k) = \eta_k \text{ for every } k \in \mathbb{N}\}.$$

By Theorem 5.1, for each $n \in \mathbb{N}$ there are functions $\varphi_{n1}, \dots, \varphi_{nn} \in H^\infty(U)$ such that $\varphi_{nj}(x_k) = \delta_{jk}$ for $j, k = 1, \dots, n$, and

$$\sup_{x \in U} \sum_{j=1}^n |\varphi_{nj}(x)| \leq M^2 + \frac{1}{n}.$$

Then a compactness argument yields a sequence (φ_j) in $H^\infty(U)$ such that $\varphi_j(x_k) = \delta_{jk}$ for all $j, k \in \mathbb{N}$ and

$$\sup_{x \in U} \sum_{j=1}^{\infty} |\varphi_j(x)| \leq M^2.$$

If we define $\varphi(x) = (\varphi_j(x))$ for $x \in U$, then φ verifies (f).

(f) \Rightarrow (e): By (f) there exists $\varphi \in H^\infty(U; l^1)$ such that $\varphi(x_k) = e_k$ for every $k \in \mathbb{N}$. By Theorem 2.1 there exists $T \in L(G^\infty(U); l^1)$ such that $T \circ \delta_U = \varphi$. Whence T verifies (e).

(e) \Rightarrow (d): This is obvious.

(d) \Rightarrow (a): Since we can readily verify that $S' = Q$, the desired conclusion follows at once (see [14, Theorem 4.7 - A]).

Thus we have shown that conditions (d), (e) and (f) are equivalent to (a).

(e) \Rightarrow (b): Since $T \circ S\xi = \xi$ for every $\xi \in l^1$, it follows that $S' \circ T'\eta = \eta$ for every $\eta \in l^\infty$. We already know that $S' = Q$. If we define $P = T'$, then P verifies (b).

(b) \Rightarrow (c): This is obvious.

(c) \Rightarrow (e): By (c) there exists $P \in L(c_0; H^\infty(U))$ such that $P\eta(x_k) = \eta_k$ for every $\eta = (\eta_k) \in c_0$ and every $k \in \mathbb{N}$. Whence it follows that $P'\delta_{x_k} = e_k$ for every $k \in \mathbb{N}$. If we define $T = P'|G^\infty U$, then T verifies (e).

Thus we have shown that conditions (b) and (c) are also equivalent to (a).

(f) \Rightarrow (h): By (f) there exists $(\varphi_j) \subset H^\infty(U)$ such that $\varphi_j(x_k) = \delta_{jk}$ for all $j, k \in \mathbb{N}$ and $\sup_{x \in U} \sum_j |\varphi_j(x)| < \infty$. Given $y = (y_j) \in l^\infty(F)$, let $P_F y \in H^\infty(U; F)$ be defined by $P_F y(x) = \sum_j \varphi_j(x) y_j$ for $x \in U$. Then P_F verifies (h).

Since the implications (h) \Rightarrow (g) and (g) \Rightarrow (f) are obvious, the proof of the theorem is complete.

6. Connection between $G^\infty(\Delta)$ and $L^1(\partial\Delta)/H_0^1(\Delta)$.

For $1 \leq p < \infty$ let $H^p(\Delta)$ denote the Banach space of all $f \in H(\Delta)$ such that

$$\|f\|_p = \lim_{r \rightarrow 1} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p} < \infty.$$

Let $1 < p \leq \infty$ and let $f \in H^p(\Delta)$. By a theorem of Fatou, the radial limits $f(e^{i\theta}) = \lim_{r \rightarrow 1} f(re^{i\theta})$ exist almost everywhere and

$$f(z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(e^{i\theta})}{1 - ze^{-i\theta}} d\theta$$

for every $z \in \Delta$. The same conclusion is true for $p = 1$, by a theorem of *F.* and *M. Riesz*. Thus for $1 \leq p \leq \infty$ we may identify $H^p(\Delta)$ with a closed subspace of $L^p(\partial\Delta)$, namely

$$H^p(\Delta) = \{f \in L^p(\partial\Delta) : \int_{-\pi}^{\pi} f(e^{i\theta}) e^{in\theta} d\theta = 0 \text{ for } n = 1, 2, 3, \dots\}.$$

If we set $H_0^1(\Delta) = \{f \in H^1(\Delta) : f(0) = 0\}$, then

$$H_0^1(\Delta) = \{f \in L^1(\partial\Delta) : \int_{-\pi}^{\pi} f(e^{i\theta}) e^{in\theta} d\theta = 0 \text{ for } n = 0, 1, 2, \dots\}.$$

Whence it follows that the canonical isometric isomorphism between $L^\infty(\partial\Delta)$ and the dual of $L^1(\partial\Delta)$ induces an isometric isomorphism between $H^\infty(\Delta)$ and the dual of $L^1(\partial\Delta)/H_0^1(\Delta)$. All this is well known and can be found in the book of *Hoffman* [5].

Since $L^1(\partial\Delta)/H_0^1(\Delta)$ is the unique predual of $H^\infty(\Delta)$, up to an isometric isomorphism (see [3, Theorem V.5.4]), it follows that $G^\infty(\Delta)$ is isometrically isomorphic to $L^1(\partial\Delta)/H_0^1(\Delta)$. But it is illustrative to give a

direct proof of this fact.

6.1 Theorem. *The spaces $G^\infty(\Delta)$ and $L^1(\partial\Delta)/H_0^1(\Delta)$ are isometrically isomorphic.*

Proof. For every $f \in H^\infty(\Delta)$ and $ze\Delta$ we have that

$$(6.1) \quad f(z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(e^{i\theta})}{1 - ze^{-i\theta}} d\theta.$$

Let $g : z \in \Delta \rightarrow g_z \in L^1(\partial\Delta)$ be defined by $g_z(e^{i\theta}) = (1 - ze^{-i\theta})^{-1}$, and let $\pi : L^1(\partial\Delta) \rightarrow L^1(\partial\Delta)/H_0^1(\Delta)$ denote the quotient mapping. If we identify $H^\infty(\Delta)$ with the dual of $L^1(\partial\Delta)/H_0^1(\Delta)$, then (6.1) tells us that $\langle f, \pi g_z \rangle = f(z)$ for every $f \in H^\infty(\Delta)$ and $z \in \Delta$. By using [7, Theorem 8.12] we easily see that $\pi \circ g \in H^\infty(\Delta; L^1(\partial\Delta)/H_0^1(\Delta))$. Then by Theorem 2.1 there exists $T \in L(G^\infty(\Delta); L^1(\partial\Delta)/H_0^1(\Delta))$ such that $T\delta_z = \pi g_z$ for every $z \in \Delta$. Since we can readily verify that the dual mapping T' is the identity on $H^\infty(\Delta)$, we can conclude that T is an isometric isomorphism, as we wanted.

Since the space $L^1(\partial\Delta)/H_0^1(\Delta)$ is weakly sequentially complete and has the Dunford-Pettis property (see [3, Theorem V.5.2] and [11, Corollary 8.1]), it is natural to pose the following problems.

6.2. Problem. Let U be a bounded open subset of a weakly sequentially complete Banach space E . Is $G^\infty(U)$ weakly sequentially complete?

6.3. Problem. Let U be a bounded open subset of a Banach space E with the Dunford-Pettis property. Does $G^\infty(U)$ have the Dunford-Pettis property?

Observe that in each of these problems the hypotheses on E are necessary, in view of Proposition 2.2.

References

- [1] S. Dineen, P. Galindo, D. Garcia and M. Maestre, *Linearization of holomorphic mappings on fully nuclear spaces with a basis*. To appear.
- [2] P. Galindo, D. Garcia and M. Maestre, *Holomorphic mappings of bounded type*. J. Math. Anal. Appl., to appear.
- [3] J. Garnett, *Bounded Analytic Functions*. Academic Press, New York, 1981.
- [4] A. Grothendieck, *Produits Tensoriels Topologiques et Espaces Nucleaires*. Memoirs American Mathematical Society N^o 16, 1955.
- [5] K. Hoffman, *Banach Spaces of Analytic Functions*. Dover, New York, 1988.
- [6] P. Mazet, *Analytic Sets in Locally Convex Spaces*. North-Holland Mathematics Studies, vol. 89, North-Holland, Amsterdam, 1984.
- [7] J. Mujica, *Complex Analysis in Banach Spaces*. North-Holland Mathematics Studies, vol. 120, North-Holland, Amsterdam, 1986.
- [8] J. Mujica, *Linearization of bounded holomorphic mappings on Banach spaces*. Trans. Amer. Math. Soc. 324 (1991), 867-887.
- [9] J. Mujica, *Linearization of holomorphic mappings of bounded type*. In: Progress in Functional Analysis. North-Holland, Amsterdam, to appear.
- [10] J. Mujica and L. Nachbin, *Linearization of holomorphic mappings on locally convex spaces*. J. Math. Pures Appl., to appear.
- [11] A. Pelczynski, *Banach Spaces of Analytic Mappings and Absolutely Summing Operators*. Regional Conference Series in Mathematics N^o 30, American Mathematical Society, 1977.
- [12] R. Ryan, *Applications of topological tensor products to infinite dimensional holomorphy*. Ph.D. thesis, Trinity College Dublin, 1980.

- [13] **M. Schottenloher**, *ε -products and continuation of analytic mappings*.
In: *Analyse Fonctionnelle et Applications*, edité par L. Nachbin, pp.
261-270. Hermann, Paris, 1975.
- [14] **A. Taylor**, *Introduction to Functional Analysis*. John Wiley, New
York, 1958.

Instituto de Matematica
Universidade Estadual de Campinas
Caixa Postal 6065
13081 Campinas, SP, Brazil

RELATÓRIOS TÉCNICOS — 1991

- 01/91 Um Método Numérico para Resolver Equações de Silvester e de Ricatti — Vera Lucia da Rocha Lopes and José Vitório Zago.
- 02/91 “Regge-Like” Relations for (Non-Evaporating) Black Holes and Cosmological Models — Vilson Tonin-Zanchin and Erasmo Recami.
- 03/91 The Exponential of the Generators of the Lorentz Group and the Solution of the Lorentz Force Equation — J. R. Zeni and Waldyr A. Rodrigues Jr.
- 04/91 Tensornorm Techniques for the (DF)-Space Problem — Andreas Defant and Klaus Floret.
- 05/91 Nonreversibility of Subsemigroups of Semi-Simple Lie Groups — Luiz San Martin.
- 06/91 Towards a General Theory of Convolutional Sets (With Applications to Fractals) — Jayme Vaz Jr.
- 07/91 Linearization of Holomorphic Mappings of Bounded Type — Jorge Mujica.
- 08/91 Topological Equivalence of Diffeomorphisms and Curves — M. A. Teixeira.
- 09/91 Applications of Finite Automata Representing Large Vocabularies — Cláudio L. Lucchesi and Tomasz Kowaltowski.
- 10/91 Torsion, Superconductivity and Massive Electrodynamics
Cartan’s Torsion Vector and Spin-0 Fields — L. C. Garcia de Andrade.
- 11/91 On The Continuity of Fuzzy Integrals — G. H. Greco and R. C. Bassanezi.
- 12/91 Optimal Chemical Control of Populations Developing Drug Resistance — M. I. S. Costa, J. L. Boldrini and R. C. Bassanezi.
- 13/91 Strict Monotonicity of Eigenvalues and Unique Continuation — Djairo G. de Figueiredo and Jean-Pierre Gossez.
- 14/91 Continuity of Tensor Product Operators Between Spaces of Bochner Integrable Functions — Andreas Defant and Klaus Floret.
- 15/91 Some Remarks on the Join of Spheres and their Particular Triangulations — Davide C. Demaria and J. Carlos S. Kuhl.
- 16/91 Sobre a Equação do Telégrafo e o Método de Riemann — L. Prado Jr. and E. Capelas de Oliveira.
- 17/91 Positive Solutions of Semilinear Elliptic Systems — Ph. Clément, D. G. de Figueiredo and E. Mitidieri.
- 18/91 The Strong Coupling Constant: Its Theoretical Derivation from a Geometric Approach to Hadron Structure — Erasmo Recami and Vilson Tonin-Zanchin.

- 19/91 Time Analysis of Tunnelling Processes, and Possible Applications in Nuclear Physics — *Vladislav S. Olkhovskiy and Erasmo Recami.*
- 20/91 Procedimento, Função, Objeto ou Lógica? — *M. Cecília Calani Baranauskas.*
- 21/91 The Relation Between 2-Spinors and Rotations — *W. A. Rodrigues Jr. and J. R. Zeni.*
- 22/91 Boundaries for Algebras of Analytic Functions on Infinite Dimensional Banach Spaces — *R. M. Aron, Y. S. Choi, M. L. Lourenço and O. W. Paques.*
- 23/91 Factorization of Uniformly Holomorphic Functions — *Luiza A. Moraes, Otilia W. Paques and M. Carmelina F. Zaine.*
- 24/91 Métrica de Prohorov e Robustez — *Mario Antonio Gneri.*
- 25/91 Cálculo de Funções de Green para a Equação de Schrödinger pelo Método de Expansão Tipo Sturm-Liouville — *L. Prado Jr. and E. Capelas de Oliveira.*
- 26/91 On the Weierstrass-Stone Theorem — *João B. Prolla.*
- 27/91 Sull'Equazione di Laplace nell'Universo di De Sitter — *E. Capelas de Oliveira and G. Arcidiacono.*
- 28/91 The Generalized Laplace Equation in Special Projective Relativity — *E. Capelas de Oliveira and G. Arcidiacono.*
- 29/91 The Projective D'Alembert Equation — *E. Capelas de Oliveira and G. Arcidiacono.*
- 30/91 The Generalized D'Alembert Equation in Special Projective Relativity — *E. Capelas de Oliveira and G. Arcidiacono.*
- 31/91 A General Algorithm for Finding the Minimal Angle between Subspaces — *Alvaro R. De Pierro and Alfredo N. Iusem.*
- 32/91 Scalar Curvature on Fibre Bundles — *Maria Alice B. Grou.*
- 33/91 Sur la Dimension des Algèbres Symétriques — *Rachid Chibloun, Artibano Micali et Jean Pierre Olivier.*
- 34/91 An Inverse Column-Updating Method for Solving Large-Scale Nonlinear Systems of Equations — *José M. Martínez em Mário C. Zambaldi.*
- 35/91 Parallel Implementations of Broyden's Method — *Francisco A. M. Gomes and José M. Martínez.*
- 36/91 Equivalência Elementar entre Feixes — *A. M. Sette and X. Caicedo.*
- 37/91 Unique Ergodicity for Degenerate Diffusions and the Accessibility Property of Control Systems — *Luiz San Martin.*
- 38/91 Unobservability of the Sign Change of Spinors Under a 2π Rotation in Neutron Interferometric Experiments — *J. E. Maiorino, J. R. R. Zeni and W. A. Rodrigues Jr.*
- 39/91 Disappearance of the Numerically irrelevant Solutions (NIS) in Non-Linear Elliptic Eigenvalue problems — *Pedro C. Espinoza.*
- 40/91 Positive Ordered Solutions of a Analogue of Non-Linear Elliptic Eigenvalue Problems — *Pedro C. Espinoza.*
- 41/91 On von Neumann's Variation of the Weierstrass-Stone Theorem — *João B. Prolla.*

- 42/91 Representable Operators and the Dunford-Pettis Theorem — *Klaus Floret.*
- 43/91 Simultaneous Approximation and Interpolation for Vector-Valued Continuous Functions — *João B. Prolla.*
- 44/91 On Applied General Equilibrium Analysis — *José A. Scaramucci.*
- 45/91 Global Solutions to the Equations for the Motion of Stratified Incompressible Fluids — *José Luiz Boldrini and Marko Antonio Rojas-Medar.*
- 46/91 A characterization of the set of fixed points of some smoothed operators — *Alfredo N. Iusem and Alvaro R. De Pierro.*
- 47/91 Lyapunov Graphs and Flows on Surfaces — *K. A. de Rezende and R. D. Franzosa.*
- 48/91 On the Multiplicative Generators of Semi-Free Circle Actions — *J. Carlos S. Kiihl and Claudina Izepe Rodrigues.*
- 49/91 A Priori Estimate and Existence of Positive Solutions of Nonlinear Cooperative Elliptic Equations Systems — *Marco Aurelio S. Souto.*
- 50/91 On a Class of Theories of Mechanics - Part I — *Jayme Vaz Jr.*
- 51/91 Complexification of Operators Between L_p -Spaces — *Klaus Floret.*
- 52/91 Function Spaces and Tensor Product — *Raymundo Alencar.*
- 53/91 The Weierstrass-Stone Theorem in Absolute Valued Division Rings — *João B. Prolla.*