

R 2805

A NEW EXPANSION FOR JACOBI FUNCTION

E. Capelas de Oliveira


Outubro

RP 47/92

RT-IMECC
IM/4134

Relatório de Pesquisa

**Instituto de Matemática
Estatística e Ciência da Computação**



**UNIVERSIDADE ESTADUAL DE CAMPINAS
Campinas - São Paulo - Brasil**

R.P.
IM/47/92

ABSTRACT - We obtain an expansion for the product of two hypergeometric functions with different parameters and different arguments. As a by product we obtain a new expansion for the Jacobi function of second kind. Particular cases are discussed.

IMECC - UNICAMP
Universidade Estadual de Campinas
CP 6065
13081 Campinas SP
Brasil

O conteúdo do presente Relatório de Pesquisa é de única responsabilidade do autor.

Outubro - 1992

A NEW EXPANSION FOR JACOBI FUNCTION

E. Capelas de Oliveira

Departamento de Matemática Aplicada

Universidade Estadual de Campinas

IMECC – UNICAMP

13083 – Campinas, SP, Brasil

ABSTRACT: We obtain an expansion for the product of two hypergeometric functions with different parameters and different arguments. As a by product we obtain a new expansion for the Jacobi function of second kind. Particular cases are discussed.

INTRODUCTION

In a recent paper [1] we presented a way to obtain many general expansions involving a product of two hypergeometric functions with different arguments. In another recent paper we obtained many new generating functions for the hypergeometric and confluent hypergeometric functions using a known expansion for the confluent hypergeometric function [2]. In both papers we obtain general expansions which generalize many other results [3], (Brafman, Osscini, Toscano, Montaldi et al).

In this paper we use an expression for a product of two hypergeometric functions with different arguments and different parameters which is interpreted as generating function for the product of two hypergeometric functions with different arguments (or alternatively, an expansion for a hypergeometric function where the argument is a double product) to obtain an expansion for a product of two Jacobi polynomials. A product of two Jacobi polynomials appear, for example, when we solve the Schrödinger differential equation for the symmetrical top parametrized in terms of Euler angles [4], using a Green's function technique.

The paper is organized as follows: In section 1 we present a brief review to obtain the expansion for the product of two hypergeometric function with different arguments.

In the section 2 we discuss the new expansion for the Jacobi function of second kind and some other cases and discussions we present in the last section.

1. A REVIEW OF THE METHOD

In this section we present an algebraic derivation of an expansion for a product of two hypergeometric functions by means of an expansion for the confluent hypergeometric function. We consider the following expansion

$$(1) \quad {}_1F_1\left[\beta; 1 + \alpha; \frac{vxz}{(1-z)(1-z+vx)}\right] = (1-z)^{-\beta+\alpha+1}(1-z+vx)^\beta \exp\left(\frac{xz}{1-z}\right) \cdot \sum_{n=0}^{\infty} \frac{z^n}{n!} \frac{\Gamma(n+\alpha+1)}{\Gamma(\alpha+1)} {}_2F_1(-n, \beta; 1 + \alpha; v) {}_1F_1(-n; 1 + \alpha; x)$$

where $|z| < 1$ and ${}_2F_1(a, b; c; x)$ is a hypergeometric function and ${}_1F_1(a; c; x)$ is a confluent hypergeometric function.

Introducing the above expansion in the following integral representation for the hypergeometric function

$$(2) \quad \Gamma(b)s^{-b} {}_2F_1(a, b; c; k/s) = \int_0^\infty \exp(-st)t^{b-1} {}_1F_1(a; c; kt)dt$$

which is valid for $|s| > |k|$ with $\text{Re } b > 0$ and making the integral (Laplace transform of the confluent hypergeometric function) for $s > z/(1-z)$ we obtain

$$(3) \quad (1-z)^{1+\alpha-\beta-b}(1-z+vx)^\beta(1-z+uv)^b \sum_{n=0}^{\infty} \frac{z^n}{n!} \frac{\Gamma(n+\alpha+1)}{\Gamma(\alpha+1)} {}_2F_1(-n, \beta+1; 1 + \alpha; v) \cdot {}_2F_1(-n, b; 1 + \alpha; u) = {}_2F_1\left[\beta, b; 1 + \alpha; \frac{zuv}{(1-z+vx)(1-z+uz)}\right]$$

where $|u| < 1$, $|v| < 1$, $|z| < 1$ and we have defined the variable u by $1/u = s - z/(1-z)$.

The above expression is an expansion for the product of two hypergeometric functions with different arguments which can be interpreted as a generating function for the product of two hypergeometric functions with different arguments or an expansion for a hypergeometric function. We note that this expansion is a symmetrical expansion; i.e. the expansion does not change when we interchange $\beta \rightleftharpoons b$ and $v \rightleftharpoons u$. We also note that

a formal relation among the parameters does not exist. This fact is important because it permits us to obtain a serie of interesting results.

2. JACOBI POLYNOMIALS

In this section we obtain a new expansion for the Jacobi polynomial with different arguments and different orders. Firstly, we introduce the following variables

$$(4) \quad \frac{v}{x} = \frac{1-z}{1-zx} \quad \text{and} \quad \frac{u}{y} = \frac{1-z}{1-zy}$$

in eq.(3) and we obtain

$$(5) \quad (1-z)^{\alpha+1}(1-xz)^{-\beta}(1-zy)^{-b} \sum_{n=0}^{\infty} \frac{z^n \Gamma(n+\alpha+1)}{n! \Gamma(\alpha+1)} \cdot {}_2F_1\left(-n, \beta+1; \alpha+1; \frac{x-xz}{1-xz}\right) \cdot {}_2F_1\left(-n, b; \alpha+1; \frac{y-yz}{1-yz}\right) = {}_2F_1(\beta, b; \alpha+1; xyz).$$

This expression can be interpreted as an expansion for a hypergeometric function where the argument is a double product or a bilinear formula for the hypergeometric function with different arguments and different parameters.

In consequence of the freedom of the parameters and using the following relation for [5] the Jacobi polynomials, $P_m^{(\mu, \nu)}(x)$,

$$(6) \quad P_m^{(\mu, \nu)}(x) = \frac{\Gamma(m+\mu+1)}{m! \Gamma(\mu+1)} {}_2F_1\left(-m, \mu+\nu+m+1; \mu+1; \frac{1-x}{2}\right)$$

we obtain

$$(7) \quad (1-z)^{\alpha+1}(1-xz)^{-\beta}(1-zy)^{-b} \sum_{n=0}^{\infty} z^n \frac{n! \Gamma(\alpha+1)}{\Gamma(n+\alpha+1)} \cdot P_n^{(\alpha; b-n-\alpha-1)}\left(\frac{1+yz-2y}{1-yz}\right) P_n^{(\alpha; \beta-n-\alpha)}\left(\frac{1+xz-2x}{1-xz}\right) = {}_2F_1(\beta, b; \alpha+1; xyz)$$

where $(\alpha)_n = \Gamma(\alpha + n)/\Gamma(\alpha)$.

In consequence of the freedom of the parameters we can obtain many other results, e.g. when we have $\nu = -\mu$ we obtain an expansion for Legendre function of second kind, and so on. We believe that this results are importants because when we discuss the Laplace or D'Alembert generalized differential equation [6], the Jacobi function appear almost naturally but this an object of the another paper.

Acknowledgements: The author express is grateful to Professor W.A. Rodrigues Jr. for constant stimulus and many other discussions and H. Assis Jr. for stimulus and CNPq for research grants.

REFERENCES

- [1] Capelas de Oliveira E 1991 *Il Nuovo Cimento* **106B**, 689
- [2] Capelas de Oliveira E 1992 *Il Nuovo Cimento* **107B**, 59
- [3] See References in the Papers [1] and [2]
- [4] Capelas de Oliveira E 1990 *Int. J. Math. Educ. Sci. Technol.* **21**, 701
- [5] Gradshteyn IS and Ryshik I M 1980 *Table of Integral, Series and Products* (Academic Press, Inc., New York, N.Y.)
- [6] Capelas de Oliveira E 1992 "On the Solution of Homogeneous Generalized wave Equation" to appear in *Journ. Math. Phys.*

RELATÓRIOS DE PESQUISA — 1992

- 01/92 Uniform Approximation the: Non-locally Convex Case — *João B. Prolla.*
- 02/92 Compactificação de $L_{\omega\omega}(Q)$ com τ Finito — *A. M. Sette and J. C. Cifuentes.*
- 03/92 Um Modelo para Aquisição da Especificação — *Cecilia Inés Sosa Arias and Ariadne Carvalho.*
- 04/92 Convergence Estimates for the Wavelet Galerkin Method — *Sônia M. Gomes and Elsa Cortina.*
- 05/92 Optimal Chemotherapy: A Case Study with Drug Resistance, Saturation Effect and Toxicity — *M. I. S. Costa, J. L. Boldrini and R. C. Bassanezi.*
- 06/92 On the Paper “Cauchy Completeness of Elementary Logic” of D. Mundici and A. M. Sette — *J. C. Cifuentes.*
- 07/92 What is the EM Algorithm for Maximum Likelihood Estimation in PET and How to Accelerate it — *Alvaro R. De Pierro.*
- 08/92 Bifurcation from infinity and multiple solutions for an elliptic system — *Raffaele Chiappinelli and Djairo G. de Figueiredo.*
- 09/92 Approximation Processes for Vector-Valued Continuous Functions — *João B. Prolla.*
- 10/92 Aplicação do Método de Fraissé à Compactificação de Lógicas com Quantificadores Co-filtro — *A. M. Sette and J. C. Cifuentes.*
- 11/92 Absolutely Summing Holomorphic Mappings — *Mário C. Matos.*
- 12/92 The Feynman-Dyson Proof of Maxwell Equations and Magnetic Monopoles — *Adolfo A. Jr. and Waldyr A. R. Jr.*
- 13/92 A Generalized Dirac’s Quantization Condition for Phenomenological Non-abelian Magnetic Monopoles — *Adolfo M. Jr. and Waldyr A. R. Jr.*
- 14/92 Multiplicity Results for the 1-Dimensional Generalized p -Laplacian — *Pedro Ubilla.*
- 15/92 Nowhere Vanishing Torsion Closed Curves Always Hide Twice — *Sueli R. Costa. and Maria Del Carmen R. Fuster.*
- 16/92 Uniform Approximation of Continuous Convex-Cone-Valued Functions — *João B. Prolla.*
- 17/92 Monotonically Dominated Operators on Convex Cones — *A. O. Chiacchio, J. B. Prolla, M. L. B. Queiroz and M. S. M. Roversi.*
- 18/92 Testing the Concept of a Photon as an Extended Object in a Variation of Franson’s Experiment — *V. Buonomano, A. J. R. Madureira and L. C. B. Ryff.*
- 19/92 A New Trust Region Algorithm for Bound Constrained Minimization — *Ana Friedlander, José Mario Martínez and Sandra A. Santos.*

- 43/92** **Periodic Motions of Sucker Rod Pumping Systems** — *Aloisio Freiria Neves and José Luiz Boldrini.*
- 44/92** **Algorithms for Solving Nonlinear Systems of Equations** — *José Mario Martínez.*
- 45/92** **Spin and Electron Structure** — *Matej Pavšič, Erasmo Recami, Waldyr A. Rodrigues Jr. and G. Daniele Maccarrone, Fabio Raciti, Giovanni Salesi.*
- 46/92** **The Initial Value Problem For the Boussinesq Equations in a Time-Dependent Domain** — *Carlos Couca Rozende and Marko Rojas Medar.*