

2175

HOLOMORPHIC GERMS ON TSIRELSON'S SPACE


*Jorge Mujica*  
and  
*Manuel Valdivia*

Janeiro

RP 04/93

## Relatório de Pesquisa

Instituto de Matemática  
Estatística e Ciência da Computação



UNIVERSIDADE ESTADUAL DE CAMPINAS  
Campinas - São Paulo - Brasil

R.P.  
IM/04/93

RT - BIMECC  
3068

**ABSTRACT** - We show that if  $K$  is an arbitrary compact subset of the Banach space constructed by Tsirelson, then the space  $\mathcal{H}(K)$  of all holomorphic germs on  $K$ , with its natural inductive limit topology, is totally reflexive.

IMECC - UNICAMP  
Universidade Estadual de Campinas  
CP 6065  
13081 Campinas SP  
Brasil

O conteúdo do presente Relatório de Pesquisa é de única responsabilidade dos autores.

Janeiro - 1993

# HOLOMORPHIC GERMS ON TSIRELSON'S SPACE

*Jorge Mujica\** and *Manuel Valdivia*

## ABSTRACT

We show that if  $K$  is an arbitrary compact subset of the Banach space constructed by Tsirelson, then the space  $\mathcal{H}(K)$  of all holomorphic germs on  $K$ , with its natural inductive limit topology, is totally reflexive.

## INTRODUCTION

If  $K$  is a compact subset of a complex Fréchet space  $E$ , then the space of holomorphic germs on  $K$  is defined as the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$ , where  $(U_n)$  is any decreasing fundamental sequence of open neighborhoods of  $K$ , and  $\mathcal{H}^\infty(U_n)$  is the Banach space of all bounded holomorphic functions on  $U_n$ . In [6] Bierstedt and Meise proved that the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$  is compact if and only if  $E$  is a Fréchet-Schwartz space, and in [7] they posed the problem of characterizing those Fréchet spaces for which the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$  is weakly compact. Not only this problem still remains open, but until now no example of a Fréchet space was known, outside Fréchet-Schwartz space, for which the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$  is weakly compact.

In this paper we remedy this situation by proving that the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$  is weakly compact for each compact subset  $K$  of the Banach space  $X$  constructed by Tsirelson in [20]. In particular  $\mathcal{H}(K)$  is totally reflexive for each compact subset  $K$  of  $X$ . Boyd [8] had previously proved that  $\mathcal{H}(K)$  is reflexive for each balanced, compact subset  $K$  of  $X$ .

As an application of our methods we show that, for each open subset  $U$  of  $X$ , the space  $\mathcal{H}(U)$  of all holomorphic functions on  $U$ , with the Nachbin topology  $\tau_\omega$ , is semi-reflexive. In the case of balanced open sets this result is due to Alencar, Aron and Dineen [2].

As a further application of our methods we show that the space  $\mathcal{H}_b(X)$  of all entire functions of bounded type on  $X$  is a totally reflexive Fréchet space.

---

\* Supported in part by DGICYT (Spain), GMD (Germany) and FAPESP (Brazil)  
Mathematics Subject Classification: 46G20, 46E50, 46B09



We refer to the books of Dineen [11] or Mujica [16] for the terminology from infinite dimensional complex analysis.

This paper was written when the first author was visiting the University of Valencia (Spain) and the University of Paderborn (Germany) during the year 1992. The authors are indebted to Richard Aron, Klaus Bierstedt and José Bonet for several helpful discussions.

## 1. SPACES OF HOLOMORPHIC GERMS

Let  $\mathcal{P}(E)$  denote the algebra of all continuous polynomials on  $E$ , let  $\mathcal{P}_w(E)$  denote the subalgebra of all  $P \in \mathcal{P}(E)$  which are weakly continuous on bounded sets, and let  $\mathcal{P}_f(E)$  denote the subalgebra generated by the dual  $E'$  of  $E$ . We always have that  $\mathcal{P}_f(E) \subset \mathcal{P}_w(E) \subset \mathcal{P}(E)$  and the relations among these algebras have been extensively studied by Aron and Prolla [4] and by Aron, Hervés and Valdivia [3].

The main tool in the proof of the next theorem is the following result of Mujica [17]: For each open subset  $U$  of a Banach space  $E$  there are a Banach space  $G^\infty(U)$  and a mapping  $\delta_U \in \mathcal{H}^\infty(U; G^\infty(U))$  with the following universal property: For each Banach space  $F$  and each mapping  $f \in \mathcal{H}^\infty(U; F)$  there is a unique operator  $T_f \in \mathcal{L}(G^\infty(U); F)$  such that  $T_f \circ \delta_U = f$ .

**1.1. THEOREM.** Let  $E$  be a reflexive Banach space such that  $\mathcal{R}(E) = \mathcal{P}_w(E)$ . Then the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$  is weakly compact, and in particular  $\mathcal{H}(K)$  is totally reflexive, for each compact subset  $K$  of  $E$ .

**PROOF.** Let  $U_n = K + V_n$ , where  $(V_n)$  is a sequence of open balls in  $E$ , centered at the origin, with  $V_{n+1} \subset \frac{1}{3}V_n$  for every  $n \in \mathbb{N}$ . By the aforementioned result of Mujica, there is an operator  $S_n : G^\infty(K + V_{n+1}) \rightarrow G^\infty(K + V_n)$  such that the following diagram is commutative:

$$\begin{array}{ccc} K + V_{n+1} & \hookrightarrow & K + V_n \\ \downarrow \delta_{K+V_{n+1}} & & \downarrow \delta_{K+V_n} \\ G^\infty(K + V_{n+1}) & \xrightarrow{S_n} & G^\infty(K + V_n). \end{array}$$

Since  $K$  is compact, for each  $n \in \mathbb{N}$  there is a finite set  $A_n \subset K$  such that  $K \subset A_n + V_n$ . Hence

$$K + V_{n+1} \subset \bigcup_{a \in A_{n+1}} (a + 2V_{n+1}) \subset \bigcup_{a \in A_{n+1}} (a + \frac{2}{3}V_n).$$

If  $f \in \mathcal{H}^\infty(a + V_n; F)$ , then it follows from the hypotheses on  $E$  that the restriction of  $f$  to the set  $a + \frac{2}{3}V_n$  is continuous for the topologies  $\sigma(E, E')$  and  $\sigma(F, F')$ . By applying this remark to the mapping  $\delta_{K+V_n}$ , we conclude that the set  $\delta_{K+V_n}(K + V_{n+1})$

is a relatively weakly compact subset of  $G^\infty(K + V_n)$ . Then [17, Proposition 3.4] guarantees that the operator  $S_n$  is weakly compact. Since the restriction mapping  $R_n : \mathcal{H}^\infty(K + V_n) \rightarrow \mathcal{H}^\infty(K + V_{n+1})$  is nothing but the transpose of  $S_n$ , we conclude that  $R_n$  is weakly compact as well. That  $\mathcal{H}(K)$  is totally reflexive follows then from results of Komatsu [13].

**1.2. EXAMPLES.** (a) Let  $X$  be the reflexive Banach space constructed by Tsirelson [20]. Then it follows from the proof of a result of Alencar, Aron and Dineen [2] that  $\mathcal{P}(X) = \mathcal{P}_w(X)$ .

(b) By using the notion of spreading models, Farmer [12] has given other examples of reflexive Banach spaces  $E$  such that  $\mathcal{P}(E) = \mathcal{P}_w(E)$ .

## 2. SPACES OF POLYNOMIALS

Let  $\mathcal{P}^{(m)}(E)$  denote the Banach space of all  $m$ -homogeneous members of  $\mathcal{P}(E)$ , let  $\mathcal{P}_w^{(m)}(E) = \mathcal{P}_w(E) \cap \mathcal{P}^{(m)}(E)$  and let  $\mathcal{P}_f^{(m)}(E) = \mathcal{P}_f(E) \cap \mathcal{P}^{(m)}(E)$ . If  $\mathcal{H}(K)$  is reflexive, then each  $\mathcal{P}^{(m)}(E)$  is reflexive as well, and it follows from a result of Alencar [1] that the sufficient conditions in Theorem 1.1 are also necessary within the class of Banach spaces with the approximation property. To improve this result we begin with the following observation, where  $\tau_c$  denotes the compact-open topology.

**2.1. PROPOSITION.** If  $E$  is a Banach space, then the following conditions are equivalent:

- (a)  $\mathcal{P}_w(E)$  is  $\tau_c$ -dense in  $\mathcal{P}(E)$ .
- (b)  $\mathcal{P}_w^{(m)}(E)$  is  $\tau_c$ -dense in  $\mathcal{P}^{(m)}(E)$  for every  $m \in \mathbb{N}$ .

The implication (b)  $\Rightarrow$  (a) is obvious, whereas the converse implication follows at once from the Cauchy integral formula (see [16, Corollary 7.3.]).

Aron and Schottenloher [5] (see [16, Theorem 28.1]) proved that if  $E$  is a Banach space with the approximation property, then  $\mathcal{P}_f(E)$  is  $\tau_c$ -dense in  $\mathcal{P}(E)$ . Similarly we have the following result.

**2.2. PROPOSITION.** If  $E$  is a Banach space with the compact approximation property, then  $\mathcal{P}_w(E)$  is  $\tau_c$ -dense in  $\mathcal{P}(E)$ .

**PROOF.** Let  $P \in \mathcal{P}(E)$ , let  $K$  be a compact subset of  $E$ , and let  $\varepsilon > 0$ . Then there is  $\delta > 0$  such that  $|P(y) - P(x)| < \varepsilon$  whenever  $x \in K$  and  $y \in E$ , with  $\|y - x\| < \delta$ . Then  $|P \circ T(x) - P(x)| < \varepsilon$  for all  $x \in K$ . Since  $T$  maps bounded, weakly convergent nets into norm convergent nets, we see that the polynomial  $P \circ T$  is weakly continuous on bounded sets.



**2.3. EXAMPLES.** (a) Pisier [18] has given an example of a Banach space  $E$ , without the approximation property, such that  $E \otimes_r E = E \otimes_r E$ . Whence  $\mathcal{P}_f(^2 E)$  is  $\tau_c$ -dense in  $\mathcal{P}(^2 E)$ .

(b) Willis [22] has given an example of a reflexive Banach space  $E$  which has the compact approximation property, but does not have the approximation property. In particular  $\mathcal{P}_w(E)$  is  $\tau_c$ -dense in  $\mathcal{P}(E)$ .

(c) As pointed out by Lindenstrauss and Tzafriri [14, p. 94], Davie [9] has given examples of reflexive Banach spaces without the compact approximation property.

However we have been unable to answer the following questions.

**2.4. QUESTIONS.** Is there a Banach space  $E$  such that  $\mathcal{P}_f(E)$  is not  $\tau_c$ -dense in  $\mathcal{P}(E)$ ? Is there a Banach space  $E$  such that  $\mathcal{P}_w(E)$  is not  $\tau_c$ -dense in  $\mathcal{P}(E)$ ?

The main tool in the proof of the next theorem is the following result of Ryan [19] (see also [17]): For each Banach space  $E$  and each  $m \in \mathbb{N}$  there are a Banach space  $Q(^m E)$  and a mapping  $\delta_m \in \mathcal{P}(^m E; Q(^m E))$  with the following universal property; For each Banach space  $F$  and each polynomial  $P \in \mathcal{P}(^m E; F)$  there is a unique operator  $T_P \in \mathcal{L}(Q(^m E); F)$  such that  $T_P \circ \delta_m = P$ . Furthermore  $Q(^m E) = (\mathcal{P}(^m E), \tau_c)'$ .

**2.5. THEOREM.** Let  $E$  be a Banach space and let  $m \in \mathbb{N}$ . Then the following conditions are equivalent:

- (a)  $E$  is reflexive and  $\mathcal{P}(^m E) = \mathcal{P}_w(^m E)$ .
- (b)  $\mathcal{P}(^m E)$  is reflexive and  $\mathcal{P}_w(^m E)$  is  $\tau_c$ -dense in  $\mathcal{P}(^m E)$ .

**PROOF.** (a)  $\Rightarrow$  (b): It follows from (a) that the polynomial  $\delta_m$  maps bounded subsets of  $E$  into relatively weakly compact subsets of  $Q(^m E)$ . Then [17, Proposition 3.4] guarantees that the identity mapping on  $Q(^m E)$  is weakly compact, and hence  $Q(^m E)$  is reflexive. Thus  $\mathcal{P}(^m E) = Q(^m E)'$  is reflexive too.

(b)  $\Rightarrow$  (a): Since  $\mathcal{P}(^m E)$  is reflexive, we have that  $\mathcal{P}(^m E)' = Q(^m E) = (\mathcal{P}(^m E), \tau_c)'$ , and therefore

$$\mathcal{P}(^m E) = \overline{\mathcal{P}_w(^m E)}^{\tau_c} = \overline{\mathcal{P}_w(^m E)}^{\|\cdot\|} = \mathcal{P}_w(^m E).$$

### 3. SPACES OF HOLOMORPHIC FUNCTIONS

**3.1. THEOREM.** Let  $E$  be a reflexive Banach space such that  $\mathcal{P}(E) = \mathcal{P}_w(E)$ . Then the space  $(\mathcal{H}(U), \tau_w)$  is semi-reflexive for each open subset  $U$  of  $E$ .

**PROOF.** Let  $K$  be a compact subset of  $U$ , and let  $(U_n)$  be a decreasing fundamental sequence of open neighborhoods of  $K$ . Following [15] let  $\mathcal{H}^K(U)$  denote the image of the

natural mapping  $\mathcal{H}(U) \rightarrow \mathcal{H}(K)$ , and let  $\hat{\mathcal{H}}^K(U)$  denote the inductive limit

$$(3.1) \quad \hat{\mathcal{H}}^K(U) = \text{ind } \overline{\mathcal{H}^K(U) \cap \mathcal{H}^\infty(U_n)},$$

where  $\overline{\mathcal{H}^K(U) \cap \mathcal{H}^\infty(U_n)}$  denotes the closure of  $\mathcal{H}^K(U) \cap \mathcal{H}^\infty(U_n)$  in  $\mathcal{H}^\infty(U_n)$ , with the induced norm. Since the inductive limit  $\mathcal{H}(K) = \text{ind } \mathcal{H}^\infty(U_n)$  is weakly compact, by Theorem 1.1, it is clear that the inductive limit (3.1) is weakly compact too. In particular the space  $\hat{\mathcal{H}}^K(U)$  is totally reflexive. Now by [15, Lemma 5.6] the space  $(\mathcal{H}(U), \tau_w)$  is topologically isomorphic to the projective limit of the spaces  $\hat{\mathcal{H}}^K(U)$ , with  $K \subset U$ . Whence the space  $(\mathcal{H}(U), \tau_w)$  is semi-reflexive.

#### 4. SPACES OF HOLOMORPHIC FUNCTIONS OF BOUNDED TYPE

If  $U$  is an open subset of a Banach space  $E$ , then  $\mathcal{H}_b(U) = \text{proj } \mathcal{H}^\infty(U_n)$ , where

$$U_n = \{x \in U : \|x\| < n \quad \text{and} \quad d_U(x) > \frac{1}{n}\}.$$

**4.1. THEOREM.** Let  $E$  be a reflexive Banach space such that  $\mathcal{P}(E) = \mathcal{P}_w(E)$ . Then the projective limit  $\mathcal{H}_b(U) = \text{proj } \mathcal{H}^\infty(U_n)$  is weakly compact, and in particular  $\mathcal{H}_b(U)$  is totally reflexive, for each balanced open subset  $U$  of  $E$ .

**PROOF.** If  $U$  is balanced, then each  $U_n$  is balanced as well, and there is  $\theta_n$ ,  $0 < \theta_n < 1$ , such that  $U_n \subset \theta_n U_{n+1}$ . Then we can show, as in the proof of Theorem 1.1, that the natural mapping  $S_n : G^\infty(U_n) \rightarrow G^\infty(U_{n+1})$  is weakly compact. Whence it follows that the restriction mapping  $R_n : \mathcal{H}^\infty(U_{n+1}) \rightarrow \mathcal{H}^\infty(U_n)$  is weakly compact as well. That  $\mathcal{H}_b(U)$  is totally reflexive follows then, either from results of Komatsu [13], or else from results of Davis et al. [10] and Valdivia [21].

From Theorems 1.1, 2.5, 3.1 and 4.1 we get the following corollary.

**4.2. COROLLARY.** Let  $E$  be a Banach space such that  $\mathcal{P}_w(E)$  is  $\tau_c$ -dense in  $\mathcal{P}(E)$ . Then the following conditions are equivalent:

- (a)  $\mathcal{H}(K)$  is totally reflexive for each (for some) compact subset  $K$  of  $E$ .
- (b)  $\mathcal{H}(K)$  is reflexive for each (for some) compact subset  $K$  of  $E$ .
- (c)  $(\mathcal{H}(U), \tau_w)$  is semi-reflexive for each (for some) open subset  $U$  of  $E$ .
- (d)  $\mathcal{H}_b(U)$  is totally reflexive for each (for some) balanced open subset  $U$  of  $E$ .
- (e)  $\mathcal{H}_b(U)$  is reflexive for each (for some) balanced open subset  $U$  of  $E$ .
- (f)  $\mathcal{P}^{(m)}(E)$  is reflexive for every  $m \in \mathbb{N}$ .
- (g)  $E$  is reflexive and  $\mathcal{P}(E) = \mathcal{P}_w(E)$ .



## References

- [1] R. ALENCAR, On reflexivity and basis for  $\mathcal{P}(\infty E)$ . Proc. Roy. Irish Acad. 85 (1985), 131-138.
- [2] R. ALENCAR, R. ARON and S. DINEEN, A reflexive space of holomorphic functions in infinitely many variables. Proc. Amer. Math. Soc. 90 (1984), 407-411.
- [3] R. ARON, C. HERVES and M. VALDIVIA, Weakly continuous mappings on Banach spaces. J. Funct. Anal. 52 (1983), 189-204.
- [4] R. ARON and J. PROLLA, Polynomial approximation of differentiable functions on Banach spaces. J. Reine Angew. Math. 313 (1980), 195-216.
- [5] R. ARON and M. SCHOTTENLOHER, Compact holomorphic mappings on Banach spaces and the approximation property. J. Funct. Anal. 21 (1976), 7-30.
- [6] K. BIERSTEDT and R. MEISE, Nuclearity and the Schwartz property in the theory of holomorphic functions on metrizable locally convex spaces. In: Infinite Dimensional Holomorphic and Applications, edited by M. Matos, pp. 93-129. North-Holland, Amsterdam, 1977.
- [7] K. BIERSTEDT and R. MEISE, Aspects of inductive limits in spaces of germs of holomorphic functions on locally convex spaces and applications to a study of  $(\mathcal{H}(U), \tau_w)$ . In: Advances in Holomorphy, edited by J.A. Barroso, pp. 111-178. North-Holland, Amsterdam, 1979.
- [8] C. BOYD, Preduals of the space of holomorphic functions on a Fréchet space. Ph.D. thesis, University College Dublin, 1992.
- [9] A. M. DAVIE, The approximation problem for Banach spaces. Bull. London Math. Soc. 5 (1973), 261-266.
- [10] W. DAVIS, T. FIGIEL, W. JOHNSON and A. PELCZYNSKI, Factoring weakly compact operators. J. Funct. Anal. 17 (1974), 311-327.
- [11] S. DINEEN, Complex Analysis in Locally Convex Spaces. North-Holland Mathematics Studies, vol. 57. North-Holland, Amsterdam 1981.
- [12] J. FARMER, polynomial reflexivity in Banach spaces. Preprint, 1992.
- [13] H. KOMATSU, Projective and injective limits of weakly compact sequences of locally convex spaces. J. Math. Soc. Japan 19 (1967), 366-383.
- [14] J. LINDENSTRAUSS and L. TZAFRIRI, Classical Banach Spaces I. Springer, Berlin, 1977.
- [15] J. MUJICA, Spaces of germs of holomorphic functions. In: Studies in Analysis, edited by G. C. Rota, pp. 1-41. Academic Press, New York, 1979.



- [16] J. MUJICA, Complex Analysis in Banach Spaces. North-Holland Mathematics Studies, vol. 120. North-Holland, Amsterdam, 1986.
- [17] J. MUJICA, Linearization of bounded holomorphic mappings on Banach spaces. Trans. Amer. Math. Soc. 324 (1991), 867-887.
- [18] G. PISIER, Counterexamples to a conjecture of Grothendieck. Acta Math, 151 (1983), 181-208.
- [19] R. A. RYAN, Applications of topological tensor products to infinite dimensional holomorphy. Ph.D. thesis, Trinity College Dublin, 1980.
- [20] B. TSIRELSON, Not every Banach space contains an imbedding of  $\ell_p$  or  $c_0$ . Functional Anal. Appl. 8 (1974), 138-141.
- [21] M. VALDIVIA, A characterization of totally reflexive Fréchet spaces. Math. Z. 200 (1989), 327-346.
- [22] G. WILLIS, The compact approximation property does not imply the approximation property. Studia Math. 103 (1992), 99-108.

*Jorge Mujica*  
 Instituto de Matemática  
 Universidade Estadual de Campinas  
 Caixa Postal 6065  
 13.081 Campinas, SP  
 Brasil

*Manuel Valdivia*  
 Facultad de Matemáticas  
 Universidad de Valencia  
 Doctor Moliner 50  
 46100 Burjasot (Valencia)  
 Spain

## RELATÓRIOS DE PESQUISA — 1993

- 01/93 On the Convergence Rate of Spectral Approximation for the Equations for Nonhomogeneous Asymmetric Fluids — *José Luis Boldrini and Marko Rojas-Medar.*
- 02/93 On Fraïssé's Proof of Compactness — *Xavier Caicedo.*
- 03/93 Non Finite Axiomatizability of Finitely Generated Quasivarieties of Graphs — *Xavier Caicedo.*