

Tightness and efficiency of irreducible automorphisms of handlebodies.

Leonardo Navarro Carvalho*
leonardo@ime.unicamp.br

June 2004

Abstract

Among (isotopy classes of) automorphisms of handlebodies those called irreducible (or generic) are the most interesting, analogues of pseudo-Anosov automorphisms of surfaces. We consider the problem of isotoping an irreducible automorphism so that it is most efficient (has minimal growth rate) in its isotopy class. We describe a property, called tightness, of certain invariant laminations, which we conjecture characterizes this efficiency. We obtain partial results towards proving the conjecture. For example, we prove it for genus two handlebodies. We also show that tightness always implies efficiency.

In addition, partly in order to provide counterexamples in our study of properties of invariant laminations, we develop a method for generating a class of irreducible automorphisms of handlebodies.

1 Introduction.

1.1 Some history and background

The classification of *automorphisms* (i.e. self-diffeomorphisms) of a manifold, up to isotopy, is a fundamental problem. Nielsen addressed the case where the manifold is a compact and connected surface and his results were later substantially improved by Thurston (see [Nie86a, Nie86b, Nie86c, Thu88, HT85, FLP79]). We briefly state their main result: An automorphism of a surface is, up to isotopy, either *periodic* (i.e., has finite order),

*Partially supported by CNPq–Brazil, CAPES–Brazil, FAPESP–São Paulo, Brazil, 03/06914-2.

reducible (i.e., preserves an essential codimension-1 submanifold) or *pseudo-Anosov*. We refer the reader to any of [FLP79, HT85, Thu88, CB88] for details, including the definition of a pseudo-Anosov automorphism. The Nielsen-Thurston theory also shows that the reducible case may be reduced to the other two cases. Since periodic automorphisms are relatively easy to understand, the remaining *irreducible* case, i.e., the pseudo-Anosov case, is the most interesting and rich one. Indeed, pseudo-Anosov automorphisms of surfaces are the subject of intense and wide research (see [Thu88]).

In [Oer02], Oertel undertakes a similar classification for a certain class of three-dimensional manifolds. Suppose that a three-dimensional manifold M is compact, connected, orientable and *irreducible* (i.e., every embedded sphere bounds a ball). Assume further that $\partial M \neq \emptyset$. By use of canonical decompositions of M due to Bonahon (determined by his *characteristic compression body* [Bon83a]) and Jaco, Shalen and Johanson (the *JSJ-decomposition* [JS79, Joh79]), the study of automorphisms of M is reduced to the study of automorphisms of *compression bodies* and *handlebodies* (see [Oer02]). We define these types of manifolds.

A *compression body* is a manifold pair (Q, F) obtained from a surface F in the following way. Consider the disjoint union of the product $F \times I$ and finitely many balls (three dimensional 0-handles) B . Attach 1-handles to $(F \times \{1\}) \cup \partial B$, obtaining Q . Identifying F with $F \times \{0\} \subseteq Q$, we obtain the compression body (Q, F) . Then $F \subseteq Q$ is the *interior boundary* of (Q, F) , denoted by $\partial_i Q$. The *exterior boundary* $\partial_e Q$ of (Q, F) is the closure $\overline{\partial Q - \partial_i Q}$. We allow empty or non-empty ∂F , but F cannot have sphere components. If Q is homeomorphic to the disjoint union of $F \times I$ with balls then (Q, F) is said to be *trivial*.

A *handlebody* H is a connected compression body whose interior boundary is empty, i.e. H is obtained from attaching 1-handles to balls. The *genus* of H is the rank of $\pi_1(H)$.

The following definition is due to Oertel:

Definition 1.1. An automorphism $f: H \rightarrow H$ of a handlebody H is said *reducible* if any of the following holds:

- there exists an f -invariant (up to isotopy) non-trivial compression body (Q, F) with $Q \subseteq H$, $\partial_e Q \subseteq \partial H$ and $F = \partial_i Q \neq \emptyset$ not containing ∂ -parallel disc components,
- there exists an f -invariant (up to isotopy) collection of pairwise disjoint, incompressible, non- ∂ -parallel and properly embedded annuli,
or

- H admits an f -invariant (up to isotopy) I -bundle structure.

The automorphism f is called *irreducible* (or *generic*, as in [Oer02]) if both of the following conditions hold:

1. $\partial f = f|_{\partial H}$ is pseudo-Anosov, and
2. there exists no *closed reducing surface* F : a closed reducing surface is a surface $F \neq \emptyset$ which is the interior boundary $\partial_i Q$ of a non-trivial compression body (Q, F) such that $Q \subseteq H$, (Q, F) is f -invariant (up to isotopy) and $\partial_e Q = \partial H$.

An obvious remark is that this definition of irreducible automorphism excludes the periodic case.

Theorem 1.2 (Oertel, [Oer02]). *An automorphism of a handlebody is either:*

1. *periodic,*
2. *reducible, or*
3. *irreducible.*

We note that the theorem above is not entirely obvious. For example, one must show that if an automorphism $f : H \rightarrow H$ of a handlebody does not restrict to a pseudo-Anosov ∂f on ∂H , then f is actually reducible according to Definition 1.1, or periodic.

Our interest is precisely in the irreducible case, which is in many ways analogous to the pseudo-Anosov case for surfaces. An important similarity is related to the existence of certain invariant projective measured laminations (see [Oer02] and Theorem 1.3 below). A good part of the original article is dedicated to the construction of these laminations, which depends on many choices. Among these, one has to choose a handlebody $H_0 \subseteq \mathring{H}$ “concentric” with H , in the sense that the complement $H - \mathring{H}_0$ is a product. Also, the automorphism $f : H \rightarrow H$ must be isotoped so that $H_1 = f(H_0)$ contains H_0 in its interior and $\bigcup_{i \in \mathbb{Z}} f^i(H_0) = \mathring{H}$. Next, a handle decomposition \mathcal{H} of H_0 as union of 0 and 1-handles is also needed (alternatively, one can choose a complete system of discs, as in [Oer02]).

We refer the reader to [Oer02] for details on the construction of the laminations. Important properties are summarized in the following. See the remark after the theorem for a comment on the nature of the singularities of the 1-dimensional lamination.

Theorem 1.3. (Oertel) *Let $f: H \rightarrow H$ be a generic automorphism of a handlebody. Then there exist a 2-dimensional measured lamination (Λ, μ) of \mathring{H} , a 1-dimensional measured singular lamination (Ω, ν) in H_0 which is transverse to Λ , \hat{f} isotopic to f and $\lambda > 1$ such that*

1. $\hat{f}(\Lambda, \mu) = (\Lambda, \lambda\mu)$,
2. $\hat{f}(\Omega, \nu) = (\Omega, \lambda^{-1}\nu)$,
3. *the leaves of Λ are open discs and fill H_0 , in the sense that the components of $H_0 - \Lambda$ are contractible,*
4. $\Lambda \cup \partial H$ *is closed in H ,*
5. $\Lambda \cap \Omega$ *is disjoint from the singular set of Ω .*

Remark. The lamination Ω contains a *singular set* $S(\Omega) = \bigcap_{i \in \mathbb{Z}} f^i(h^0)$, where h^0 is the union of 0-handles of H_0 . One can choose \hat{f} so that $S(\Omega)$ is finite. The complement $\Omega - S(\Omega)$ is a lamination of $H_0 - S(\Omega)$, where the measure ν is defined. Also, the notion of tangency to Ω is not defined at $S(\Omega)$. Therefore, in general, by saying that a surface F is *transverse* to Ω we assume, in particular, that $F \cap S(\Omega) = \emptyset$.

The problem with this theorem is that these laminations, whose construction depends on many choices, are not unique in any reasonable sense. For example, the *growth rate* λ , which is a measure of the complexity of the automorphism, is not unique. This phenomenon is not unlike what happens with automorphisms of surfaces, if one allows the invariant laminations to have monogons for complementary components.

The main problem we shall address here, though not solve, is the following.

Problem. *Characterize canonical invariant laminations for a given irreducible automorphism.*

In the case of surfaces, a solution is to fix a hyperbolic metric in its interior and work with certain geodesic laminations which, among other properties, realize minimal growth. For automorphisms of handlebodies there is no such solution. Still, there is a minimum in the set of possible growth rates. Naturally, canonical laminations must yield minimal growth, i.e. the corresponding automorphism must be most *efficient* in its isotopy class. A step (a big step, we believe) in the direction of solving the problem above would then be to characterize minimal growth. Oertel gives a necessary condition.

Theorem 1.4. [Oer02] *If λ is minimal then Λ has the incompressibility property: for each leaf L of Λ the complement $L - \mathring{H}_0$ is incompressible in $\mathring{H} - \mathring{H}_0$.*

A clear sufficient condition, much stronger than incompressibility, is that the leaves of Ω do not “back-track” (with respect to Λ) in H_0 (see [BH92]). But it cannot always be realized (see remark below).

We will also consider the problem of constructing examples of irreducible automorphisms of handlebodies. As in any field of mathematics, examples provide a useful investigative tool. The construction of irreducible automorphisms of handlebodies is not an obvious task. The main difficulty resides in proving that a given example does not admit closed reducing surfaces (the other property, that the restriction to the boundary is pseudo-Anosov, can be achieved with the help of some well established tools, e.g. [Pen88, BH95]).

A result of Bonahon [Bon83b] implies that any automorphism of a genus two handlebody whose restriction to the boundary is a pseudo-Anosov automorphism is then irreducible (see also [Lon88] and [Car03]). This is not true for higher genus handlebodies.

Remark. This may be used to generate interesting examples. For instance, in [FL80] the authors build an automorphism of a genus two handlebody which 1) restricts to the boundary as a pseudo-Anosov automorphism — thus, as mentioned before, is irreducible — and 2) induces the identity on the fundamental group. In particular, the leaves of Ω have to “back-track”. Such an example illustrates the richness of irreducible automorphisms of handlebodies when compared with pseudo-Anosov automorphisms of surfaces, whose complexity is captured on the level of the fundamental group.

1.2 Summary of Results

Our main results address the problem of characterizing minimal growth of a given irreducible automorphism of a handlebody. We will identify a certain condition on the two-dimensional lamination Λ , which we call “tightness”¹. Essentially, Λ is “tight” if the weighted intersection of its leaves with (Ω, ν) is minimal. The property of being tight is (strictly) stronger than that of Oertel’s incompressibility and (strictly) weaker than that of having “no back-tracking”. We conjecture that it characterizes minimal growth.

(Conjecture 3.4). *The growth rate is minimal if and only if the lamination is tight.*

¹In fact, being tight is a property of the pair of measured laminations.

This work will prove one direction:

(Theorem 3.5). *If Λ is tight then λ is minimal.*

As for necessity, the problem is harder. We will prove it only under some technical hypotheses (Theorem 3.6). These hypotheses are useful: we will show that they can be assumed for genus two handlebodies. In this case tightness characterizes minimal growth:

(Corollary 3.7). *Conjecture 3.4 is true for genus two handlebodies.*

Moreover, tightness yields results concerning the growth rates. We prove:

(Corollary 3.17). *If Λ is tight, then the growth (which is minimal) is less than or equal to the growth of the restriction of the automorphism $f: H \rightarrow H$ to the boundary ∂H (which is pseudo-Anosov).*

The following is a corollary of Theorem 3.5.

(Corollary 3.16). *If Λ is tight then the minimal growth $\lambda_{\min}(f^n)$ of any power f^n is $(\lambda_{\min}(f))^n$.*

From our point of view, the measures ν , μ on the invariant laminations Ω , Λ and the corresponding growth rate λ come from eigenvectors and eigenvalue of certain incidence matrices of a handle decomposition \mathcal{H} of the handlebody (or complete disc system \mathcal{E}). Such a matrix M is non-negative and *irreducible*, having many nice spectral properties. For instance, the spectral radius of M , which we denote by $\lambda(M)$, is realized by a positive real eigenvalue. We call it the *Perron-Frobenius eigenvalue* of M . The corresponding positive eigenvector, which is well-defined (up to scaling, naturally), is called the *Perron-Frobenius eigenvector*. The following is useful (see e.g. [Sen73, BH92]):

Proposition 1.5. *Let M be a non-negative and irreducible $n \times n$ matrix and $v \in \mathbb{R}^n$ with $v_i \geq 0$ and $v \neq 0$. If*

$$(Mv)_i \leq \lambda v_i, \quad \text{for all } i$$

then $\lambda(M) \leq \lambda$ and $v_i > 0$. If, moreover, $(Mx)_i < \lambda x_i$ for some i , then $\lambda(M) < \lambda$.

We will also present a method for generating a certain class of irreducible automorphisms. This method produces examples on higher genus handlebodies. The fact that our techniques fail to prove Conjecture 3.4 for a general

automorphism of a handlebody of genus greater than two makes it especially important to study examples in higher genus cases. Our method will depend on the two following results. We refer the reader to [Pen88] (see also Definition 2.4) and [BH92] respectively for details and precise definitions.

Theorem 1.6 (Penner). *Let \mathcal{C}, \mathcal{D} be two systems of closed curves in an oriented surface S with $\chi(S) < 0$. Assume that \mathcal{C} and \mathcal{D} intersect efficiently, do not have parallel components and fill S . Let $f: S \rightarrow S$ be a composition of Dehn twists: right twists along curves of \mathcal{C} and left twist along curves of \mathcal{D} . If a twist along each curve appears at least once in the composition, then f is isotopic to a pseudo-Anosov automorphism of S .*

Theorem 1.7. *Let S be a compact surface with $\chi(S) < 0$ and precisely one boundary component. An automorphism $f: S \rightarrow S$ is pseudo-Anosov if and only if f_*^n is irreducible for all $n > 0$.*

The results of this article are divided in two following sections. In Section 2 we will describe our method for generating examples of irreducible automorphisms. We will develop a particular case and then generalize it in Theorem 2.5. Its statement depends on some technical constructions unsuited for this introduction. We will then use it to build a certain irreducible automorphism of a genus four handlebody (Example 2.8). It will help in motivating the relevance of the property of tightness. For this reason, we shall determine a certain pair of invariant laminations for this automorphism and estimate the corresponding growth rate. The two-dimensional lamination will have Oertel’s incompressibility property.

Section 3 is dedicated to the tightness property. We shall see that the example built in the previous section does not realize minimal growth. The lack of tightness, which we will define then, will appear naturally there. This will be done through the existence of “tightening discs”, which will be our main objects in dealing with lack of tightness. In the remainder of the section we shall prove the theorems and corollaries on tightness already mentioned.

We will adopt the following notations and conventions. Given a topological space A (typically a manifold or sub-manifold), \overline{A} will denote its topological closure, $\overset{\circ}{A}$ its interior and $|A|$ its number of connected components. If H is a handlebody we denote a handle decomposition of H by \mathcal{H} . By considering co-cores of 1-handles (which we may also call *dual discs*) it is clear that a handle decomposition \mathcal{H} of H corresponds to a complete system of discs $\mathcal{E} \subseteq H$. In fact, the set of handle decompositions and the set of complete disc systems are, up to isotopy, in 1 – 1 correspondence.

This remark is relevant for while [Oer02] uses discs systems — because the author focuses on the two-dimensional lamination —, we shall use handle decompositions — because we focus more on the one-dimensional lamination. The incidence matrix associated to a \mathcal{H} corresponds to the transpose of the incidence matrix associated to the corresponding \mathcal{E} . There is also an embedded graph $\Gamma \subseteq H$ dual to \mathcal{E} , with vertices corresponding to 0-handles of \mathcal{H} and edges corresponding to 1-handles. We can then regard H as a fibered neighborhood of Γ . Such embedded graphs are also, up to isotopy, in 1 – 1 correspondence with handle decompositions and complete disc systems.

I thank Ulrich Oertel for many enlightening meetings and helpful suggestions in his role as dissertation advisor, and also for laying the foundations on which the research in this paper is built.

2 Examples.

2.1 An example.

We will develop a simple particular case of the method that will be obtained in the following subsection. Let H be a genus 2 handlebody. An automorphism of H will be described as a composition of Dehn twists along two annuli and a disc. We shall prove that it is irreducible by showing that its restriction to ∂H is pseudo-Anosov and that, for an algebraic reason, there can be no closed reducing surface. The argument that proves this last part is distinct from that of [Bon83b] for genus 2 handlebodies and, with the right hypotheses, generalizes to higher (even) genus handlebodies.

Example 2.1. We start with a pseudo-Anosov automorphism $\varphi: S \rightarrow S$ of the once punctured oriented torus S . Such a φ will be defined as a composition of Dehn twists along two curves.

We represent S as a cross with pairs of opposite sides identified as shown in Figure 1. Fixing a base point in S we note that $\pi_1(S)$ is the free group on two generators.

Let α_0, α_1 be simple closed curves as in the figure. It is easy to verify that the systems $\mathcal{C} = \{\alpha_0\}$ e $\mathcal{D} = \{\alpha_1\}$ satisfy the hypotheses of Theorem 1.6 (Penner). Let T_0^- be the left Dehn twist along α_0 and T_1^+ the right twist along α_1 . We define:

$$\varphi = T_1^+ \circ T_0^-.$$

By Theorem 1.6, φ is pseudo-Anosov. Then, by Theorem 1.7, any positive




Figure 1: The oriented surface S and the curves α_0, α_1 .

power φ_*^n , of the induced isomorphism $\varphi_*: \pi_1(S) \rightarrow \pi_1(S)$ is *irreducible*². We note this fact for future use.

We now consider the handlebody $H = S \times I$, identifying S with $S \times \{1\}$. The orientation of S then determines an orientation on H through inclusion. Now lift φ to H , obtaining $\phi: H \rightarrow H$, a composition of twists along the annuli $A_0 = \alpha_0 \times I$, $A_1 = \alpha_1 \times I$ as in Figure 2. Identifying $\pi_1(H)$ with $\pi_1(S)$ yields $\phi_* = \varphi_*$.




Figure 2: The automorphism f is defined as a composition of Dehn twists along the annuli A_0, A_1 and the disc Δ .

Finally, we will obtain the desired irreducible automorphism $f: H \rightarrow H$ by composing ϕ with a twist along a disc Δ , shown in Figure 2. Let T_Δ^+ be

²We refer the reader to [BH92] for the definition of an *irreducible automorphism of a free group*.

the right Dehn twist along Δ . We define:

$$f = T_{\Delta}^+ \circ \phi.$$

Proposition 2.2. *The automorphism $f: H \rightarrow H$ is irreducible.*

The lack of closed reducing surfaces will come from the following general lemma.

Lemma 2.3. *Let $g: H \rightarrow H$ be an automorphism of a handlebody H such that ∂g is pseudo-Anosov. If g is reducible then, for some $n \in \mathbb{N}$, $g_*^n: \pi_1(H) \rightarrow \pi_1(H)$ is a reducible automorphism of a free group.*

Remark. It is clear that the result does not depend on the choice of base point.

Proof. Let Q be a compression body invariant under g . Let $F \subseteq \partial_i Q$ be a component of the closed reducing surface. It is easy to see that it bounds a handlebody $J \subseteq \overset{\circ}{H}$. Choosing a base point in J (see the remark above) and omitting the obvious inclusion homomorphisms, we claim that

$$\pi_1(H) = \pi_1(J) * G,$$

where G is not trivial. To see this, first consider the connected and nontrivial compression body $Q' = \overline{H - J}$, whose boundary decomposes as $\partial_i Q' = F$ and $\partial_e Q' = \partial H$. A compression body structure of Q' gives it as a product $F \times I$ to which 1-handles are attached. Regarding $F \times I \subseteq Q' \subseteq H$, we see that the handlebody $J' = (F \times I) \cup J$ deformation retracts to J (so $\pi_1(J') = \pi_1(J)$ through inclusion). Using the compression body structure of Q' we can regard H as J' with 1-handles attached to $\partial J'$. Since $\partial J'$ is connected, we can moreover assume that these 1-handles are attached to a disc in $\partial J'$, which gives $\pi_1(H) = \pi_1(J') * G = \pi_1(J) * G$, where G is a free group (whose rank equals the number of 1-handles of Q'). But Q' is not trivial, so G is not trivial, proving the claim. Therefore $\pi_1(J)$ is a proper free factor of $\pi_1(H)$.

Let g^n be the first power of g preserving J . Isotoping g we assume moreover that the base point is fixed by g^n . From

$$g^n(J) = J$$

it follows that $g_*^n(\pi_1(J))$ is conjugate to $\pi_1(J)$, hence g_*^n is reducible. \square

Proof of Proposition 2.2. We need to prove that $\partial f = f|_{\partial H}$ is pseudo-Anosov and that f does not admit closed reducing surfaces.

We start by verifying that ∂f is pseudo-Anosov. It is given as composition of Dehn twists: left twists along curves of

$$\mathcal{C} = \{(\alpha_0 \times \{1\}), (\alpha_1 \times \{0\})\},$$

(see Figure 2) and right twists along curves of

$$\mathcal{D} = \{(\alpha_0 \times \{0\}), (\alpha_1 \times \{1\}), \partial\Delta\}.$$

We now note that \mathcal{C}, \mathcal{D} satisfy the hypotheses of Theorem 1.6 (also see Definition 2.4), hence ∂f is pseudo-Anosov.

Now suppose by contradiction that there exists a closed reducing surface. Recalling that Lemma 2.3 does not depend of the choice of base point, there exists n such that f_*^n is reducible. But $f = (T_{\Delta}^+) \circ \phi$ and the twist (T_{Δ}^+) (along a disc) induces the identity in $\pi_1(H)$. Therefore, recalling that $\pi_1(H)$ is identified with $\pi_1(S)$, we have that $f_*^n = \phi_*^n = \varphi_*^n$, which was seen before to be irreducible for any n , a contradiction. This shows that there are no closed reducing surfaces, completing the proof. \square

2.2 A method for generating irreducible automorphisms.

The construction of Example 2.1 may be generalized to provide a method for generating a larger class of irreducible automorphisms of handlebodies (Theorems 2.5 and 2.7). It partially solves a problem proposed in [Oer02].

Definition 2.4. Let $(\mathcal{C}, \mathcal{D})$ be a pair of curve systems in a compact, connected and orientable surface S with $\chi(S) < 0$. It is called a *Penner pair in S* if \mathcal{C}, \mathcal{D} satisfy the hypotheses of Penner's Theorem 1.6 i.e.,

1. each \mathcal{C}, \mathcal{D} is a finite collection of simple, closed and pairwise disjoint essential curves without parallel copies,
2. \mathcal{C} and \mathcal{D} intersect efficiently, do not have parallel components and *fill* S (i.e., the components of $S - (\mathcal{C} \cup \mathcal{D})$ are either contractible or deformation retract to a component of ∂S).

Suppose that $(\mathcal{C}, \mathcal{D})$ is a Penner pair. An automorphism φ of S obtained from \mathcal{C}, \mathcal{D} as in Theorem 1.6 is called a *Penner automorphism subordinate to $(\mathcal{C}, \mathcal{D})$* .

If $\partial S \neq \emptyset$ then a properly embedded and essential arc θ is called *dual to $(\mathcal{C}, \mathcal{D})$* if θ intersects $\mathcal{C} \cup \mathcal{D}$ transversely and in exactly one point $p \notin \mathcal{C} \cap \mathcal{D}$.

We constructed the irreducible automorphism in Example 2.1 by lifting a pseudo-Anosov automorphism of a surface to a product and composing it with a twist on a disc. The general method will be similar. Our interest in dual arcs is that we can use them to construct discs that will yield irreducible automorphisms.

Throughout this subsection we fix a compact, connected and oriented surface S with $\partial S \neq \emptyset$ and define $H = S \times I$, which is a handlebody. We identify S with $S \times \{1\} \subseteq H$, inducing orientation in H . We also fix a base point in $S \times \{1\}$ for both S and H and identify $\pi_1(H)$ with $\pi_1(S)$.

Given a Penner pair $(\mathcal{C}, \mathcal{D})$ in S and a dual arc θ , we build a disc Δ_θ in H in the following way. Let γ be the curve of $(\mathcal{C}, \mathcal{D})$ that θ intersects and assume without loss of generality that $\gamma \subseteq \mathcal{C}$. Let $D = \theta \times I \subseteq H$. Then ∂D intersects $\gamma_1 = \gamma \times \{1\}$ in a point. Now let Δ_θ be the *band sum* of D with itself along γ_1 . This means that Δ_θ is obtained from D and γ_1 by the following construction. Consider a regular neighborhood $N = N(D \cup \gamma_1)$. Then $\Delta_\theta = \overline{\partial N - \partial H}$ is a properly embedded disc.

Theorem 2.5. *Suppose that $\partial S \neq \emptyset$ has exactly one component. Let $(\mathcal{C}, \mathcal{D})$ be a Penner pair in S with dual arc θ and $\varphi: S \rightarrow S$ a Penner automorphism subordinate to $(\mathcal{C}, \mathcal{D})$. Let $\hat{\varphi}: H \rightarrow H$ be the lift of φ to the product $H = S \times I$ and $\Delta_\theta \subseteq H$ the disc constructed from the arc θ as above. Then there exists a simple Dehn twist $T_{\Delta_\theta}: H \rightarrow H$ along Δ_θ such that the composition*

$$\hat{\varphi} \circ T_{\Delta_\theta}: H \rightarrow H$$

is an irreducible automorphism of H .

The key to the proof is the verification that \mathcal{C} , \mathcal{D} and $\partial\Delta_\theta$ determine a Penner pair in ∂H .

Lemma 2.6. *Let S , $(\mathcal{C}, \mathcal{D})$, θ , $H = S \times I$ and Δ_θ be as in the statement of Theorem 2.5. Let $\mathcal{C}_i = \mathcal{C} \times \{i\} \subseteq S_i = S \times \{i\}$ and $\mathcal{D}_i = \mathcal{D} \times \{i\} \subseteq S_i = S \times \{i\}$, defining $\mathcal{C}_0, \mathcal{D}_0 \subseteq S_0$ and $\mathcal{C}_1, \mathcal{D}_1 \subseteq S_1$. Under these conditions the following system of curves in ∂H :*

$$\begin{aligned} \mathcal{Q} &= \mathcal{D}_0 \cup \mathcal{C}_1 \cup \{\partial\Delta_\theta\}, \\ \mathcal{R} &= \mathcal{C}_0 \cup \mathcal{D}_1, \end{aligned}$$

determine a Penner pair $(\mathcal{Q}, \mathcal{R})$ in ∂H .

Proof. We start by making the obvious remarks that $\mathcal{C}_0, \mathcal{D}_0, \mathcal{C}_1, \mathcal{D}_1 \subseteq \partial H$ and $\mathcal{C}_0 \cap \mathcal{D}_1 = \emptyset, \mathcal{D}_0 \cap \mathcal{C}_1 = \emptyset$. Recall that we are assuming that $\theta \cap (\mathcal{C} \cup \mathcal{D}) \subseteq \gamma \subseteq \mathcal{C}$. We verify that:

- $\partial\Delta_\theta \cap \mathcal{D}_0 = \emptyset$, because $(\theta \times \{0\}) \cap \mathcal{D}_0 = \emptyset$ and $\partial\Delta_\theta \cap S_0$ consists of two arcs in S_0 parallel to $\theta \times \{0\}$,
- $\partial\Delta_\theta \cap \mathcal{C}_1 = \emptyset$, because $\partial\Delta_\theta \cap \gamma_1 = \emptyset$ by construction.

Therefore each $\mathcal{Q} = \mathcal{D}_0 \cup \mathcal{C}_1 \cup \{\partial\Delta\}$ and $\mathcal{R} = \mathcal{C}_0 \cup \mathcal{D}_1$ is a system of simple closed curves essential in ∂H . To conclude that $(\mathcal{Q}, \mathcal{R})$ is indeed a Penner pair it remains to verify that $\mathcal{Q} \cup \mathcal{R}$ fills ∂H .

A component of $S - (\mathcal{C} \cup \mathcal{D})$ either is a disc or an annulus that retracts to ∂S . Therefore a component of $\partial H - (\mathcal{C}_0 \cup \mathcal{D}_0 \cup \mathcal{C}_1 \cup \mathcal{D}_1)$ either is a disc or an annulus A (that retracts to $\partial S \times I$). But $A \cap \partial\Delta_\theta$ is a union of four arcs essential in A , hence each component of $\partial H - (\mathcal{Q} \cup \mathcal{R})$ is a disc, showing that $\mathcal{Q} \cup \mathcal{R}$ fills ∂H , completing the proof. \square

Instead of proving Theorem 2.5 we will prove the more general result below, which clearly implies the other. We note that twists on curves of \mathcal{C} , \mathcal{D} in S lift to twists along annuli in H . We call these systems of annuli $\hat{\mathcal{C}}$, $\hat{\mathcal{D}}$ respectively. The “direction of a twist” along these vertical annuli should be understood as the direction of its restriction to $S \times \{1\} \subseteq \partial H$.

Theorem 2.7. *Let $(\mathcal{C}, \mathcal{D})$, S , θ , H and Δ_θ be as in Theorem 2.5. Let f be a composition $f: H \rightarrow H$ of twists along the annuli of $\hat{\mathcal{C}}$, $\hat{\mathcal{D}}$ and the disc Δ_θ : in one direction along the annuli in $\hat{\mathcal{D}}$ and in the opposite direction along the annuli in $\hat{\mathcal{C}}$ and the disc Δ_θ . If each of these twists appear in the composition at least once f is irreducible.*

Proof. We first show that $f_*^n: \pi_1(H) \rightarrow \pi_1(H)$ is an irreducible automorphism of a free group for any $n \geq 0$ (hence there can be no closed reducing surface by Lemma 2.3) and then that $\partial f = f|_{\partial H}$ is pseudo-Anosov, thus completing the proof that f is irreducible.

Recall that S is identified with $S \times \{1\} \subseteq H$ and $\pi_1(S)$ with $\pi_1(H)$. Let T_{Δ_θ} be a twist along Δ_θ . Since $(T_{\Delta_\theta})_*: \pi_1(H) \rightarrow \pi_1(H)$ is the identity (Δ_θ is a disc) the hypotheses on f imply that $f_* = \varphi_*$ for some Penner automorphism $\varphi: S \rightarrow S$ subordinate to $(\mathcal{C}, \mathcal{D})$. Penner automorphisms are pseudo-Anosov so, given that ∂S has a single component, it follows from Theorem 1.7 that φ_*^n is an irreducible automorphism of $\pi_1(S)$ for any $n \geq 0$. Therefore $f_*^n: \pi_1(H) \rightarrow \pi_1(H)$ is irreducible, and then f does not admit closed reducing surfaces (Lemma 2.3).

To see that ∂f is pseudo-Anosov, let $(\mathcal{Q}, \mathcal{R})$ be as in Lemma 2.6, therefore a Penner pair. By construction the twists that compose f restrict to ∂H as twists along curves of \mathcal{Q} or \mathcal{R} . It is then straightforward to verify that

∂f is a Penner automorphism subordinate to $(\mathcal{Q}, \mathcal{R})$, hence pseudo-Anosov, completing the proof that f is irreducible. □

Remark. Note that the condition that $|\partial S| = 1$ implies that H has even genus.

Example 2.8. Consider S a genus 2 surface minus a disc, represented in Figure 3 as an octagon whose sides are identified according to the arrows.




Figure 3: A Penner pair in S , with dual arc θ .

In the picture there are represented four further curves: α , β , γ and δ . Defining

$$\begin{aligned} \mathcal{C} &= \{\beta, \delta\}, \\ \mathcal{D} &= \{\alpha, \gamma\}, \end{aligned}$$

it is easy to check that $(\mathcal{C}, \mathcal{D})$ is a Penner pair in S . The automorphism $\varphi: S \rightarrow S$ defined by

$$\varphi = T_{\beta}^{-} \circ T_{\delta}^{-} \circ T_{\alpha}^{+} \circ T_{\gamma}^{+}$$

is, therefore, a Penner automorphism subordinate to the pair $(\mathcal{C}, \mathcal{D})$.

The pair $(\mathcal{C}, \mathcal{D})$ admits dual arcs. The picture shows one, labelled as θ . We consider the corresponding disc Δ_{θ} . Figure 4 shows $S_0 = S \times \{0\}$, $S_1 = S \times \{1\} \subseteq \partial H$ and how $\partial\Delta_{\theta}$ intersects them³.

Figure 4 also shows the pair $(\mathcal{Q}, \mathcal{R})$ obtained by Lemma 2.6: \mathcal{Q} consists of the “solid” curves, including $\partial\Delta_{\theta}$, while the “dashed” lines form \mathcal{R} .

³In fact, one can picture the whole $\partial\Delta_{\theta} \cap \partial H$ in the figure. The only portion of ∂H not represented is the vertical annulus $\partial S \times I$, which $\partial\Delta_{\theta}$ intersects in four vertical arcs.




Figure 4: The curve $\partial\Delta_\theta$ in ∂H .

Let $\hat{\varphi}: H \rightarrow H$ be the lift of φ to H . By Theorem 2.7

$$\hat{\varphi} \circ T_{\Delta_\theta}^-: H \rightarrow H$$

is an irreducible automorphism, where $T_{\Delta_\theta}^-$ is the left twist along Δ_θ .

2.3 The incompressibility condition.

Example 2.9. We continue working with Example 2.8. Now we will determine certain f -invariant laminations and estimate their corresponding growth rate λ . We refer the reader to [Oer02] for details on this construction.

Recall the oriented surface S , a once punctured genus two surface. Here it will be convenient to regard $S \times I$ as $H_0 \subseteq H$. We now choose a complete system of discs \mathcal{E}_0 in H_0 , as follows. Consider the labeled arcs A, B, C and $D \subseteq S$ as in the figure (the sides of the octagon), and define the discs $A \times I, B \times I, C \times I, D \times I \subseteq H_0$. Abusing notation, we use the same labels A, B, C and D to represent these discs. Let $\mathcal{E}_0 = \{A, B, C, D\}$. From \mathcal{E}_0 we can consider the dual graph Γ . In fact, we will regard Γ as an oriented labeled graph (represented as a spine in Figure 5, on the left). To avoid ambiguities, we require that $\Gamma \subseteq S \times \{\frac{1}{2}\}$. We also consider the corresponding handle decomposition \mathcal{H}_0 in H_0 .

Together with the handle decomposition we will now choose a representative in the class of f (which will be called f) and study the associated laminations. These are determined by how the handles of H_0 intersect the handles of $H_1 = f(H_0)$. But it is equivalent to consider $H_{-1} = f^{-1}(H_0) \subseteq H_0$ (apply the diffeomorphism f^{-1}), which is easier to picture.

Regarding H_0 as a neighborhood of Γ , we consider $f^{-1}(\Gamma) = T_{\Delta_\theta}^+ \circ (\hat{\varphi})^{-1}(\Gamma) \subseteq H_0$ (the disc Δ_θ and the automorphism $\hat{\varphi}$ are defined in Example 2.8). Figure 5, right, shows this image $f^{-1}(\Gamma)$, determining how Γ , and hence H_0 , should be pictured in H_1 .




Figure 5: The graph $\Gamma \subseteq H_0$ and $\Gamma \subseteq H_1$.

This intersection pattern determines an f -invariant measured lamination Λ with full support provided that the incidence matrix corresponding to \mathcal{E}_0 is irreducible. We verify this using the transpose of that matrix, which is the incidence matrix $M(\mathcal{H}_0)$ for the 1-handles of \mathcal{H}_0 . To do that, consider the handle decomposition \mathcal{H}_1 of H_1 induced by \mathcal{H}_0 through f . The incidence matrix $M(\mathcal{H}_0) = \{m_{ij}\}$ is given by $m_{ij} = |f(e_i) \cap e_j|$, where $f(e_i)$ is a 1-handle of \mathcal{H}_1 and e_j is a 1-handle of \mathcal{H}_0 . Therefore,

$$M(\mathcal{H}_0) = \begin{pmatrix} 3 & 1 & 1 & 0 \\ 4 & 1 & 3 & 2 \\ 1 & 0 & 2 & 1 \\ 1 & 0 & 1 & 1 \end{pmatrix},$$

whose Perron-Frobenius eigenvalue is, up to three decimals,

$$\lambda = \lambda(\mathcal{H}_0) \approx 4.987.$$

We now ask the question of whether λ is the minimal growth in the isotopy class of f or not. The “no back-tracking” sufficient condition does not apply to this case (for instance, there is an unremovable “back-tracking”,

linked on the circled part of Figure 5, right). Oertel’s incompressibility property, a necessary condition, holds here, as methodic, though tedious, computation reveals⁴.

We see then that, up to this point, we have no tools to decide whether $\lambda \approx 4.987$ is minimal or not. In the next section, Section 3, a simple argument shows that this growth rate is not minimal. This will be related to the fact that this lamination is not “tight”.

3 Tightness.

From now on we assume that $f: H \rightarrow H$ is an irreducible automorphism of a handlebody H . We consider a handle decomposition \mathcal{H}_0 of H_0 and the corresponding: disc system \mathcal{E}_0 , f -invariant measured laminations (Λ, μ) , (Ω, ν) and growth rate λ . The 1-dimensional lamination Ω and its measure ν will play an important role throughout this section. Given an immersed surface $F \subseteq H$ transverse to Ω we will denote $\int_F \nu$ by either $\nu(F)$ or $F \bullet (\Omega, \nu)$. The advantage of the first notation is in its simplicity and will be preferred whenever there are no ambiguities. The advantage of the second is that it emphasizes the object Ω supporting ν , which will be helpful in certain contexts. We call $\nu(F) = F \bullet (\Omega, \nu)$ the *weighted intersection* (or just the *intersection*) of F with (Ω, ν) .

Recall that the goal is to characterize minimal growth.

3.1 Tightening discs.

In Example 2.9 we left unproven the claim that Oertel’s incompressibility property does not imply minimal λ .

So assume that Λ has the incompressibility property and consider the associated growth rate λ . The next schematic example suggests a reason for the fact that this incompressibility does not imply minimality of λ . Recall that a handle decomposition \mathcal{H}_0 of H_0 determines, through f^i , a handle decomposition \mathcal{H}_i of $H_i = f^i(H_0)$, $i \in \mathbb{Z}$.

Example 3.1. Let V be a 0-handle of \mathcal{H}_1 and suppose that $H_0 \cap V$ is as in Figure 6 a), with V intersecting 1-handles e_p and e_q of \mathcal{H}_0 , and $f(E)$ the image of a disc $E \in \mathcal{E}_0 = \{E_1, \dots, E_k\}$.

⁴One can carry on this computation in a manner not unlike those in [BH92], [BH95]. Here one just has to be careful with certain back-trackings, which are allowed because of linkings (e.g., the circled section in Figure 5).




Figure 6: a) A disc $f(E)$ dual to a 1-handle of H_1 , intersecting the 1-handle e_i of \mathcal{H}_0 ; b) the move that reduces λ .

Now consider $M = M(\mathcal{H}_0) = \{m_{ij}\}$, the incidence matrix for the 1-handles of \mathcal{H}_0 : $m_{ij} = |f(e_i) \cap e_j|$ counts how many times the 1-handle e_j (of \mathcal{H}_0) crosses the 1-handle $f(e_i)$ of \mathcal{H}_1 . We can assume that M is irreducible and recall that its Perron-Frobenius eigenvalue is λ , the growth rate of f (with respect to \mathcal{H}_0).

Now note that E_i is transverse to Ω , so it makes sense to consider

$$\hat{\nu}_i = \nu(E_i),$$

where we recall that ν is the transverse measure on Ω . That determines a vector $\hat{\nu} = (\hat{\nu}_1, \dots, \hat{\nu}_k)$, which is precisely a Perron-Frobenius eigenvector of M :

$$M\hat{\nu} = \lambda\hat{\nu}.$$

Suppose that $\hat{\nu}_q < \hat{\nu}_p$. We can isotope f to replace intersections of $f(E)$ with e_p by intersections with e_q (see Figure 6 b)).

This operation does not change the handle decomposition. Now the new incidence matrix $M' = \{m'_{ij}\}$ is given by:

$$\begin{aligned} m'_{ij} &= m_{ij} & \text{if } ij \neq 1p, 1q; \\ m'_{ip} &= m_{ip} - 2; \\ m'_{iq} &= m_{iq} + 2. \end{aligned}$$

Assume that M' is irreducible. We recall that $\hat{\nu}_q < \hat{\nu}_p$ and consider $M'\hat{\nu}$:

$$\begin{aligned} (M'\hat{\nu})_i &= \lambda\hat{\nu}_i & \text{if } i \neq 1; \\ (M'\hat{\nu})_1 &= \lambda\hat{\nu}_1 - 2\hat{\nu}_p + 2\hat{\nu}_q < \lambda\hat{\nu}_1. \end{aligned}$$

By Proposition 1.5

$$\lambda(M') < \lambda = \lambda(M),$$

therefore the isotopy reduces the growth rate.

The situation described in the example above indeed happens, see Example 3.3. It not only shows that Oertel's incompressibility property does not imply minimality of the growth rate but also suggests that the weighted intersection $E \bullet (\Omega, \nu) = \nu(E)$ (where $E \in \mathcal{E}_0$) should be relevant in the search for the minimal growth. We introduce, then, the following definition:

Definition 3.2. Let $(\Delta, \partial\Delta) \rightarrow (H, \Lambda)$ be an embedded disc transverse to Ω . Consider $\Delta' \subseteq \Lambda$ such that $\partial\Delta' = \partial\Delta$. We say that Δ is a *tightening disc* for the triple (Λ, Ω, ν) if

$$\nu(\Delta) < \nu(\Delta').$$

The triple (Λ, Ω, ν) is said *tight* if there is no tightening disc. We will often abuse notation and say that Λ is tight or not, leaving (Ω, ν) implicit. Accordingly, we may say that a tightening disc for (Λ, Ω, ν) is a tightening disc for Λ only.

Remark. The requirement that the tightening disc Δ is transverse to Ω implies that it does not intersect the singular set $S(\Omega)$.

We also note that Λ being tight implies that it has the incompressibility property: a compressing disc for $\Lambda - \mathring{H}_0$ is a tightening disc.

Now we can say that the original lamination in Example 3.1 is not tight, with a tightening disc represented in Figure 6 b). As previously mentioned, that is a hypothetical situation. The following is a specific example.

Example 3.3. We refer to Example 2.9 and consider the automorphism $f: H \rightarrow H$ and the handle decomposition \mathcal{H}_0 of H_0 defined then. We consider the disc $f(B) \subseteq H_1$, the co-core of a handle of \mathcal{H}_1 .

One can see in Figure 7 a tightening disc Δ (represented at the right by a dashed line), with its boundary in a leaf of $\Lambda \cap H_1$ parallel to $f(B)$. Indeed, let $\Delta' \subseteq f(B)$ be the disc such that $\partial\Delta' = \partial\Delta$ (represented by a thick line). It is easy to check that

$$\nu(\Delta) = 2\nu(D) < 2(\nu(A) + \nu(C) + \nu(D)) = \nu(\Delta'),$$

for ν has full support on Ω . We now can change f through an isotopy taking Δ' to Δ . Figure 8 shows the result of such an isotopy.




Figure 7: A tightening disc Δ in H_1 .

We verify that the new incidence matrix is:

$$M(\mathcal{H}) = \begin{pmatrix} 3 & 1 & 1 & 0 \\ 2 & 1 & 1 & 2 \\ 1 & 0 & 2 & 1 \\ 1 & 0 & 1 & 1 \end{pmatrix},$$

whose Perron-Frobenius eigenvalue is $\lambda(\mathcal{H}) \approx 4.542$ (precise up to three decimals), showing that the previous lamination did not have minimal growth rate.

We are interested in the problem of characterizing minimal growth rate. Considering these examples one should expect tightness to play a role in the problem.

Conjecture 3.4. *The growth rate is minimal if and only if the lamination Λ is tight.*

At this point we note that tightness is strictly stronger than Oertel's incompressibility — which is too weak — and strictly weaker than “no back-tracking” — which is too strong.

In the direction of proving the conjecture we will show that:

Theorem 3.5. *If Λ is tight then λ is minimal.*

Concerning the converse we will prove:




Figure 8: After the growth-reducing isotopy.

Proposition 3.6. *Suppose that all 0-handles of the handle decomposition \mathcal{H} have valence 2 or 3. If λ is minimal then Λ is tight.*

By *valence* of a 0-handle of \mathcal{H}_0 we mean the number of ends of 1-handles that are attached to it. In other words, if Γ is the graph corresponding to \mathcal{H}_0 , consider the valence of the vertex corresponding to the 0-handle. As an application of the technical proposition above we will get:

Corollary 3.7. *Conjecture 3.4 is true if the handlebody has genus 2.*

The proofs will be given in the following subsections.

3.2 Strong tightening discs.

In this subsection we introduce some technical constructions and results.

The definition of tightening disc that we gave was quite general. We will work with tightening discs having some special properties. These will be called *strong tightening disc* (see Definition 3.8 below). We will show that there is no loss of generality in working with them (Proposition 3.9).

Consider the handle decomposition \mathcal{H}_0 of H_0 , with 1-handles e_1, \dots, e_k . We will give weights to these 1-handles, i.e. assign a positive number v_j to each 1-handle e_j . More precisely, given a positive vector $v = (v_1, \dots, v_k)$, ($v_j > 0$), we say that the pair (\mathcal{H}_0, v) is a *weighted handle decomposition* of H_0 . In this case we say that v is a *system of weights in \mathcal{H}_0* . We extend

these notions for any level i , so it makes sense to refer to a weighted handle decomposition (\mathcal{H}_i, v) for any $H_i = f^i(H_0)$.

Consider a weighted handle decomposition (\mathcal{H}_i, v) for H_i and let S be an embedded surface intersecting H_i just in its 1-handles. Suppose further that $S \cap H_i$ consists of discs dual to the 1-handles e_j of \mathcal{H}_i . We define

$$S \bullet (\mathcal{H}_i, v) = \sum_{1 \leq j \leq k} |S \cap e_j| \cdot v_j.$$

If E_j is a disc dual to the 1-handle e_j of H_0 we consider

$$\hat{\nu}_j = E_j \bullet (\Omega, \nu) = \nu(E_j).$$

Note that the vector $\hat{\nu} = (\hat{\nu}_1, \dots, \hat{\nu}_k)$ is a Perron-Frobenius eigenvector of the incidence matrix associated to \mathcal{H}_0 and f . We regard it as a system of weights in \mathcal{H}_0 , defining $(\mathcal{H}_0, \hat{\nu})$, which we call the *standard weighted handle decomposition of H_0* .

It is clear by the equalities above that

$$E_j \bullet (\mathcal{H}_0, \hat{\nu}) = E_j \bullet (\Omega, \nu).$$

Hence, for a general surface S intersecting H_0 in dual discs,

$$S \bullet (\mathcal{H}_0, \hat{\nu}) = S \bullet (\Omega, \nu). \tag{1}$$

In the definition below $(\mathcal{H}_0, \hat{\nu})$ is standard, as constructed above.

Definition 3.8. Let $(\Delta, \partial\Delta) \subseteq (H, \Lambda)$ be an embedded disc and $\Delta' \subseteq \Lambda$ be such that $\partial\Delta' = \partial\Delta$. We say that Δ is a *strong tightening disc* if there exists n such that the following hold:

1. $\Delta \subseteq H_n$ and $\partial\Delta \subseteq \partial H_n$;
2. for any $0 \leq i \leq n$, $\Delta \cap H_i$ consists of essential discs (in H_i) and $\Delta \cap H_0$ consists, moreover, of discs dual to the 1-handles;
3. $\Delta \cap \Lambda = \partial\Delta$;
4. $\Delta \bullet (\mathcal{H}_0, \hat{\nu}) < \Delta' \bullet (\mathcal{H}_0, \hat{\nu})$.

Proposition 3.9. *There exists a tightening disc if and only if there exists a strong tightening disc.*

Proof. One direction is immediate: a strong tightening disc is a tightening disc by property 4 and the equation (1) above.

We prove the other direction. Let Δ be a tightening disc. We will build another one with the required properties. These properties will be realized progressively (not necessarily in the order specified in Definition 3.8 above). For simplicity of notation, we will also label these intermediate discs as Δ .

Part 1: property 3

Let n be such that $\Delta \subseteq H_n$. For each 1-handle e_i of H_n , choose a dual disc $E_i \subseteq H_n$. This choice is arbitrary but for the handle e_i containing $\partial\Delta$, where we ask that $\partial\Delta \subseteq E_i$. We assume that Δ is transverse to $\bigcup_i E_i$.

The idea is to perform surgeries that reduce the complexity $|\Delta \cap \bigcup_i E_i|$ preserving the property of being a tightening disc, eventually yielding $\Delta \cap \bigcup_i E_i = \partial\Delta$. At that point we can then isotope Δ along the I -fibers of the 1-handles so that $\Delta \cap \Lambda = \partial\Delta$.

Consider $\Delta \cap \bigcup_i E_i$ (which consists just of closed curves) and choose a curve $\gamma \subseteq E_i$ that is innermost in E_i . Let $D \subseteq E_i$ and $D' \subseteq \Delta$ be the discs bounded by γ . There are two cases to consider:

1. $\nu(D) \leq \nu(D')$.

In this case we perform a surgery in Δ , replacing $D' \subseteq \Delta$ by D and pushing it a bit away from E_i (this pushing should be vertical, i.e., along the I -fibers of the product structure $D^2 \times I$ of the 1-handle e_i , and “to the side opposite to” D'). This surgery does not increase $\nu(\Delta)$ and reduces $|\Delta \cap \bigcup_i E_i|$.

2. $\nu(D) > \nu(D')$.

Here D' is a tightening disc by definition. If $\gamma \neq \partial\Delta$ then $|D' \cap \bigcup_i E_i| < |\Delta \cap \bigcup_i E_i|$ and we replace Δ with D' , reducing complexity. If $\gamma = \partial\Delta$ then there are several possibilities:

- there exists another curve $\gamma' \subseteq (\Delta \cap \bigcup_i E_i)$, $\gamma' \neq \gamma$, innermost in some E_j . In this case $\gamma' \neq \partial\Delta$ and we apply the procedure described above for γ' , reducing $|\Delta \cap \bigcup_i E_i|$.
- $\Delta \cap \bigcup_i E_i = \gamma = \partial\Delta$ and we have achieved what was desired.
- $|\Delta \cap \bigcup_i E_i| \geq 2$ and no curve $\gamma' \subseteq (\Delta \cap \bigcup_i E_i)$, $\gamma' \neq \gamma$, other than $\partial\Delta$ is innermost in the dual disc E_j that contains it. Therefore $\partial\Delta \subseteq E_j$ and, if $i \neq j$, $\Delta \cap E_i = \emptyset$. Let then $\gamma' \subseteq \Delta \cap E_j$ be *second innermost* in E_j , in the sense that the (interior of the) disc $D'' \subseteq E_j$ it bounds

contains just innermost curves. Since we are assuming that $\partial\Delta$ is the single innermost curve, $D'' \cap \Delta = \gamma' \cup \partial\Delta$. On the other hand, $\gamma' \subseteq \Delta$ bounds a disc $\Delta'' \subseteq \Delta$. We shall prove that Δ'' is a tightening disc. To see that, recall that $\Delta'' \subseteq \Delta$ and $\Delta' \subseteq D''$ and, therefore:

$$\nu(\Delta'') \leq \nu(\Delta) \quad \text{and} \quad \nu(\Delta') \leq \nu(D'').$$

But Δ is a tightening disc, i.e. $\nu(\Delta) < \nu(\Delta')$. Combining these inequalities one gets that

$$\nu(\Delta'') < \nu(D''),$$

i.e., Δ'' is a tightening disc. It is also clear that $|\Delta'' \cap \bigcup_i E_i| < |\Delta \cap \bigcup_i E_i|$, reducing complexity. We relabel Δ'' as Δ .

In any case, complexity is reduced, so eventually $\Delta \cap \bigcup_i E_i = \partial\Delta$. Since we can regard the 1-handles of \mathcal{H}_n as neighborhoods of the E_i 's, we may assume that those do not intersect Δ . An exceptional case is for E_i containing $\partial\Delta$. Here we assume that E_i is an extreme boundary leaf (there are two choices, one works) of $\Lambda \cap e_i$, where e_i is the corresponding 1-handle. Finally the resulting disc Δ has the property that $\Delta \cap \Lambda = \partial\Delta$.

Part 2: property 1

Again let n be such that $\Delta \subseteq H_n$. Let L be the leaf of $\Lambda \cap H_n$ containing $\partial\Delta$. Hence $\partial\Delta$ bounds a disc $\Delta' \subseteq L$. Consider the annulus $A = \overline{L - \Delta'}$ (Figure 9).




Figure 9: The curve $\partial\Delta$ separates a disc Δ' from an annulus A in L .

The desired disc D (satisfying property 1) will be, essentially, $\Delta \cup A$. Just push $\Delta \cup A$ slightly away from L , leaving $\partial(\Delta \cup A)$ unchanged. Since $\Delta \cup A$ is transverse to Ω we can do that preserving $\nu(\Delta \cup A)$. Therefore the resulting disc D has the property that $\nu(D) = \nu(\Delta \cup A)$.

It is easy to see that D still is a tightening disc because the whole operation increased intersection with (Ω, ν) by the same amount $\nu(A)$ both on the side of the disc and that of the leaf. More precisely,

$$\nu(D) = \nu(\Delta) + \nu(A) < \nu(\Delta') + \nu(A) = \nu(L).$$

We relabel D as Δ , realizing property 1 (and preserving property 3).

Part 3: property 4

At this point we have a tightening disc Δ and n such that, $(\Delta, \partial\Delta) \subseteq (H_n, \partial H_n)$ and $\Delta \cap \Lambda = \partial\Delta$.

It is clear that, for a sufficiently large j , Δ does not intersect the 0-handles of H_{-j} , otherwise Δ would intersect the singular set $S(\Omega)$. In fact, for a sufficiently large j , Δ is transverse to the I -fibers of the 1-handles $D^2 \times I$ of H_{-j} , otherwise Δ would not be transverse to Ω (we recall that, for a 1-handle e_i , $\Omega \cap e_i$ consists of I -fibers). By taking a sufficiently large N , $f^N(\Delta)$ will intersect H_0 only in its 1-handles and transverse to its I -fibers. We relabel $f^N(\Delta)$ as Δ and $n + N$ as n .

Now note that, since Δ is transverse to the fibers of the 1-handles e_i of H_0 , $\Delta \cap e_i$ consists of discs parallel to dual discs. We can then isotope Δ so that $\Delta \cap e_i$ consists of dual discs. This isotopy does not change $\Delta \bullet (\Omega, \nu)$ and Δ still is a tightening disc. From (1) $\Delta \bullet (\mathcal{H}_0, \hat{\nu}) = \Delta \bullet (\Omega, \nu)$, so it follows that Δ satisfies property 4 of a strong tightening disc.

Properties 3 and 1 of Δ were preserved, so it only remains also to realize property 2.

Part 4: property 2

We assume the following technical lemma, which will be proved later:

Lemma 3.10. *Let $(E, \partial E) \subseteq (H_m, \partial H_m)$ be an embedded disc intersecting H_0 in discs dual to the 1-handles. If $E \cap \Lambda$ (which may be empty) is contained in ∂E then there exists a disc $E'' \subseteq H_m$ such that:*

- $E'' \cap \Lambda \subseteq \partial E'' = \partial E$,
- $E'' \cap H_{m-1}$ consists of discs essential in H_{m-1} ,
- $E'' \cap H_0$ consists of dual discs, and
- $E'' \bullet (\mathcal{H}_0, \hat{\nu}) \leq E \bullet (\mathcal{H}_0, \hat{\nu})$.

So suppose that, for some $0 \leq j \leq n - 1$, $\Delta \cap H_j$ does not consist of essential discs. Let $m - 1$ be the biggest such value. Therefore the components of $\Delta \cap H_m$ are essential discs. Apply the lemma for each such component E , replacing $E \subseteq \Delta$ by E'' . Now $\Delta \cap H_j$ consists of essential discs for all $m - 1 \leq j \leq n$ and we proceed by induction. In the end Δ will satisfy property 2 of the definition of strong tightening disc.

We note that the other assertions of the lemma assure that the other properties 1, 3 and 4 of Δ are preserved.

This completes the proof of the proposition up to the proof of Lemma 3.10. \square

Corollary 3.11. *The lamination is tight if and only if there exists no strong tightening disc.*

Proof of Lemma 3.10. Perturb E in a neighborhood of ∂H_{m-1} so that it is transverse to ∂H_{m-1} . We want to change E so that $E \cap H_{m-1}$, which is a planar surface, consists just of essential discs. A natural strategy would be the following: isotope any inessential component of $E \cap H_{m-1}$ away from H_{m-1} , so we would have to deal only with essential components. These would be either discs, which we want, or compressible in H_{m-1} . Simplify the compressible components by compressing them. The problem with this last step is that these compressions, being performed in H_{m-1} , could introduce undesired intersections with H_0 , increasing $E \bullet (\mathcal{H}_0, \hat{\nu})$. Instead, we will perform surgeries in $H_m - \mathring{H}_{m-1}$, where they will not be subject to this problem.

We start by making some quite general comments. ‘‘Cut H_m open along Λ ’’ (i.e., take the completion of $H_m - \Lambda$ under a path-metric inherited from a metric in H_m) and let C be the component that contains E . We claim that $\partial H_{m-1} \cap C$ is incompressible in $C - \mathring{H}_{m-1}$. Indeed, let F be a component of $\partial H_{m-1} \cap C$ and consider the following commutative diagram:

$$\begin{array}{ccc} \pi_1(F) & \xrightarrow{i_1} & \pi_1(C - \mathring{H}_{m-1}) \\ i_2 \downarrow 1-1 & & \downarrow i_3 \\ \pi_1(\partial H_{m-1}) & \xrightarrow[1-1]{i_4} & \pi_1(H_m - \mathring{H}_{m-1}) \end{array}$$

where all homomorphisms are induced by the corresponding inclusion. The inclusion i_2 is injective (because $\Lambda \cap \partial H_{m-1}$ consists of essential curves in ∂H_{m-1}), as is i_4 (because $H_m - \mathring{H}_{m-1} \simeq \partial H_{m-1} \times I$). Therefore $i_4 \circ i_2$, and

thus $i_3 \circ i_1$, is injective. Then i_1 is injective and hence F is incompressible in $C - \mathring{H}_{m-1}$.

From $\partial E \subseteq \partial H_m$ follows that $E \cap (\partial H_{m-1} \cap C)$ consists of simple closed curves. Therefore, if $S = E \cap (C - \mathring{H}_{m-1})$, then ∂S has exactly one component in $\partial H_m \cap C$ (namely, ∂E) and all the others in $\partial H_{m-1} \cap C$.

We now describe the operations that change E to yield the desired final E'' . They will be performed in the interior of $H_m \cap C$, therefore will not introduce intersections with Λ or with ∂H_m . For simplicity of notation, we also label the modified discs by E .

- Process 1

Let γ be a component of $E \cap (\partial H_{m-1} \cap C)$. If $\gamma \subseteq (\partial H_{m-1} \cap C)$ is not essential then it bounds a disc $D \subseteq (\partial H_{m-1} \cap C)$. But γ bounds a disc $D' \subseteq E$, so we can isotope D' to D (also pushing any part of E that is on the way).

Since $D \cap H_0 = \emptyset$ this operation does not introduce intersections with H_0 , hence it does not increase $E \bullet (\mathcal{H}_0, \hat{\nu})$. It is also clear that it reduces $|E \cap (\partial H_{m-1} \cap C)|$. By repeating the process we can assume that $E \cap (\partial H_{m-1} \cap C)$ consists only of curves that are essential in $\partial H_{m-1} \cap C$.

Consider F a component of $S = E \cap (C - \mathring{H}_{m-1})$. If F is a disc then, by the previous paragraph, $\partial F \subseteq (\partial H_{m-1} \cap C)$ is an essential curve. Hence F is a compressing disc for $\partial H_{m-1} \cap C$ in $C - \mathring{H}_{m-1}$, a contradiction. So S cannot contain discs.

Now let γ be a component of $E \cap (\partial H_{m-1} \cap C)$ which is innermost in E . It bounds a disc $E' \subseteq E$. From the paragraph above $E' \subseteq (H_{m-1} \cap C)$, which will be essential (by previous considerations). We conclude that when all components of $E \cap (\partial H_{m-1} \cap C)$ are innermost in E (i.e., when S is connected) the proof is complete.

Suppose that that is not the case. The following procedure allows us to assume that S is incompressible (in $C - \mathring{H}_{m-1}$).

- Process 2

If there exists a compressing disc for S in $C - \mathring{H}_{m-1}$ we compress S along it.

Irreducibility says that the compression is realized by an isotopy. The isotopy does not increase $E \bullet (\mathcal{H}_0, \hat{\nu})$ because the compressing disc does not intersect H_0 . Moreover, it reduces $|E \cap (\partial H_{m-1} \cap C)| = |\partial S|$, so we can repeat the process and assume that S is incompressible in $C - \mathring{H}_{m-1}$.

The next step makes S connected and hence, as was argued before, $E \cap H_{m-1}$ consisting of essential discs.

- Process 3

Supposing that S is not connected, let S' be a component such that $\partial S' \subseteq (\partial H_{m-1} \cap C)$. Now consider S' in $H_m - \mathring{H}_{m-1}$. We verify that S' , being incompressible in $C - \mathring{H}_{m-1}$, is incompressible in $H_m - \mathring{H}_{m-1}$. Indeed, if D is a compressing disc for S' (in $H_m - \mathring{H}_{m-1}$) we can simplify $D \cap \Lambda$ by standard “cut and paste” techniques until $D \cap \Lambda = \emptyset$. These surgeries will not change ∂D , so D still is a compressing disc. But now $D \subseteq C$, contradicting incompressibility of S' in $C - \mathring{H}_{m-1}$. So S' is incompressible in $H_m - \mathring{H}_{m-1}$.

But $H_m - \mathring{H}_{m-1}$ has a product structure $(H_m - \mathring{H}_{m-1}) \simeq \partial H_{m-1} \times I$. It is a well known fact that an incompressible surface S' in such a product with $\partial S' \subseteq \partial H_{m-1} \times \{0\}$ is parallel to a surface $S'' \subseteq \partial H_{m-1}$. Let P be the product bounded by $S' \cup S''$. It is easy to see that $P \cap \Lambda = \emptyset$, so we can use the I -fibers of P to isotope $E \cap P$ vertically, through an isotopy taking S' a bit further than S'' .

We note that this operation reduces $|S|$, does not change intersections with H_0 and does not introduce intersections of E with Λ (recall that $P \cap \Lambda = \emptyset$). Repetition of the process yields $|S| = 1$, i.e. S is connected and, therefore, $E \cap H_{m-1}$ consists of essential discs.

We relabel E as E'' . It is clear that it satisfies all the desired conditions in the statement of the lemma, completing the proof. \square

3.3 Minimality and tightness.

We will use strong tightening discs to perform isotopies that reduce the growth rate.

Lemma 3.12. *If there is a strong tightening disc $\Delta \subseteq H_1$ then λ is not minimal.*

Remark. Example 3.1 shows a particular case of the proposition. The proof follows essentially the same argument that we gave in the example.

Proof. Let $(\mathcal{H}_0, \hat{\nu})$ be the standard weighted handle decomposition of H_0 and consider the weighted decomposition $(\mathcal{H}_1, \lambda \hat{\nu})$. Since $\partial \Delta \subseteq H_1 \cap \Lambda$ then

$\partial\Delta = \partial E_{i_0}^1$, where $E_{i_0}^1 = f(E_{i_0})$ is a disc dual to a 1-handle of H_1 . Moreover, from (1):

$$\Delta \bullet (\mathcal{H}_0, \hat{\nu}) < E_{i_0}^1 \bullet (\mathcal{H}_0, \hat{\nu}) = \lambda \hat{\nu}_{i_0}.$$

Since $\Delta \cap H_0$ consists of discs dual to the 1-handles of \mathcal{H}_0 we can change f through an isotopy taking E_{i_0} to Δ , without changing the handle decomposition and preserving admissibility.

Let $M' = (m'_{ij})$ be the new incidence matrix. Then $m'_{ij} = m_{ij}$ if $i \neq i_0$ and $m'_{i_0j} = m_{i_0j} + d_j$, where $d_j \in \mathbb{Z}$ satisfies the following inequality:

$$\sum_{1 \leq j \leq k} d_j \hat{\nu}_j < 0.$$

If M' is irreducible then for $i \neq i_0$,

$$(M' \hat{\nu})_i = (M \hat{\nu})_i = \lambda \hat{\nu}_i$$

but, for the row i_0 ,

$$(M' \hat{\nu})_{i_0} = (M \hat{\nu})_{i_0} + \sum_{1 \leq j \leq k} d_j \hat{\nu}_j < \lambda \hat{\nu}_{i_0},$$

hence, by Proposition 1.5,

$$\lambda' = \lambda(M') < \lambda,$$

completing the proof in this case.

If M' is reducible then, possibly permuting the indexes, M' may be written in the form:

$$M' = \left(\begin{array}{c|c} * & * \\ \hline 0 & M'' \end{array} \right),$$

where, for some $1 \leq r \leq k-1$, M'' is irreducible of dimension $(k-r)$. From the original system of discs $\mathcal{E} = \{E_1, \dots, E_k\}$ we pass to an irreducible subsystem $\mathcal{E}'' = \{E_{r+1}, \dots, E_k\}$. The transpose of the incidence matrix for these discs is precisely M'' . We will show that also in this case $\lambda(M'') < \lambda(M)$.

Since M is irreducible there exist $r+1 \leq i_1 \leq k$ and $1 \leq j_1 \leq r$ such that $m_{i_1 j_1} > 0$. Let $\hat{\hat{\nu}}$ be the $(k-r)$ -vector defined by $\hat{\hat{\nu}}_i = \hat{\nu}_{i+r}$, $1 \leq i \leq k-r$ (i.e., $\hat{\hat{\nu}}$ consists of the last $k-r$ coordinates of $\hat{\nu}$). It follows that, for $i \neq i_0 - r$, $i_1 - r$,

$$(M'' \hat{\hat{\nu}})_i \leq (M \hat{\nu})_{i+r} = \lambda \hat{\nu}_{i+r} = \lambda \hat{\hat{\nu}}_i$$

and, for $i = i_1 - r$,

$$(M''\hat{\nu})_{i_1-r} \leq (M\hat{\nu})_{i_1} - (m_{(i_1j_1)})\hat{\nu}_{j_1} < (M\hat{\nu})_{i_1} = \lambda\hat{\nu}_{i_1} = \lambda\hat{\nu}_{i_1-r}.$$

If $1 \leq i_0 \leq r$ then it follows that $\lambda'' = \lambda(M'') < \lambda$, proving the lemma in this case.

If $r+1 \leq i_0 \leq k$, in addition to the inequalities above, we further have (for $i = i_0 - r$)

$$(M''\hat{\nu})_{i_0-r} = \sum_{r+1 \leq j \leq k} (n'_{i_0j})\hat{\nu}_j \leq \sum_{1 \leq j \leq k} (n'_{i_0j})\hat{\nu}_j = (M'\hat{\nu})_{i_0} < \lambda\hat{\nu}_{i_0} = \lambda\hat{\nu}_{i_0-r},$$

therefore $\lambda'' < \lambda$, completing the proof of the lemma. □

The hypothesis of Δ being contained in H_1 in the statement of Lemma 3.12 is needed because the growth-reducing move has to be performed equivariantly. The difficulty in proving the conjecture is precisely in finding such a disc in H_1 . We can do that under the hypotheses of Proposition 3.6 (i.e., maximum 0-handle valence 3), as stated in Lemma 3.13 below. The reason is, essentially, that under such conditions every properly embedded essential disc in H_0 which is disjoint from the original disc system is parallel to a disc of this system.

Lemma 3.13. *Suppose that all 0-handles of the handle decomposition \mathcal{H}_0 have valence 2 or 3. If $\Lambda(\mathcal{H})$ is not tight then there exists a strong tightening disc in H_1 .*

Proof. Supposing that Λ is not tight we use Proposition 3.9 to get a strong tightening disc. Among all such discs, let Δ have minimal *height*, in the sense that if H_n contains a strong tightening disc then $\Delta \subseteq H_n$. Let n be the smallest n such that $\Delta \subseteq H_n$. The lemma states that $n = 1$, so assume otherwise that $n \geq 2$.

Consider $\Delta \cap H_{n-1}$, which consists of essential discs in H_{n-1} . If such a disc D does not intersect any 0-handles then it is contained in a 1-handle of H_{n-1} . In this case it is clear that D is parallel to a dual disc. If D intersects a 0-handle we can assume it is actually contained in it: using the product structure on the 1-handles that D intersects and the fact that $D \cap \Lambda = \emptyset$, we can isotope D into that 0-handle. Since the handle has valence at most 3, again D is parallel to a disc dual to a 1-handle. So $\Delta \cap H_{n-1}$ consists of discs parallel to the co-cores of the 1-handles.

Let D be such a disc, parallel to the co-core $E_i^{n-1} \subseteq H_{n-1}$. Since $D \subseteq \Delta$ (which is a strong tightening disc) then $D \cap H_0$ consists of co-cores of H_0 , so it makes sense to consider $D \bullet (\mathcal{H}_0, \hat{\nu})$. If

$$D \bullet (\mathcal{H}_0, \hat{\nu}) \geq E_i^{n-1} \bullet (\mathcal{H}_0, \hat{\nu}) \quad (2)$$

for all discs $D \subseteq \Delta \cap H_{n-1}$, then we can alter Δ by an isotopy in such a way that each D is moved to the corresponding dual disc. This operation does not increase $\Delta \bullet (\mathcal{H}_0, \hat{\nu})$ and preserves the other properties of strong tightening discs. Now $E_i^{n-1} \bullet (\mathcal{H}_0, \hat{\nu}) = \lambda^{n-1} \hat{\nu}_i$ is precisely the weight $E_i^{n-1} \bullet (\Omega, \nu)$ on the i -th 1-handle of H_{n-1} . If we apply f^{-n+1} to Δ and E_i^{-n+1} the inequality

$$f^{-n+1}(D) \bullet (\mathcal{H}_0, \hat{\nu}) \geq E_i \bullet (\mathcal{H}_0, \hat{\nu}),$$

is obtained from (2) by multiplying both sides by a factor of λ^{-n+1} , proving that $f^{-n+1}(\Delta) \subseteq H_1$ is a strong tightening disc, a contradiction to the assumption $n \geq 2$.

The argument above then shows that

$$D \bullet (\mathcal{H}_0, \hat{\nu}) < E_i^{n-1} \bullet (\mathcal{H}_0, \hat{\nu})$$

for some $D \subseteq D \cap H_{n-1}$. Modifying D through an isotopy supported in a regular neighborhood of ∂H_{n-1} (hence preserving $D \bullet (\mathcal{H}_0, \hat{\nu})$), we can assume that $\partial D = \partial E_i^{n-1}$. Now $D \subseteq H_{n-1}$ is a strong tightening disc, contradicting minimality of n .

Therefore $n = 1$ and the proof is complete. \square

We recall and prove:

(Proposition 3.6). *Suppose that all 0-handles of the handle decomposition \mathcal{H} have valence 2 or 3. If $\Lambda(\mathcal{H})$ is not tight then $\lambda(\mathcal{H})$ is not minimal.*

Proof. Use Lemma 3.13 and apply Lemma 3.12. \square

Corollary 3.14. *Let $f: H \rightarrow H$ be a generic automorphism of a handlebody of genus 2. If λ is minimal then Λ is tight.*

Proof. Consider the handle decomposition \mathcal{H}_0 of H_0 with co-core \mathcal{E}_0 . If every 0-handle has valence 2 or 3 then Proposition 3.6 completes the proof. So we assume that this is not the case. Since H has genus 2, then it has just one 0-handle with valence 4 and the others (possibly none) with valence 2.

We sketch the proof in the 1-dimensional setting: we consider Γ_0 the graph corresponding to \mathcal{H}_0 , which will have a valence 4 vertex and some

valence 2 vertices. Using the height function in $H_1 - \mathring{H}_0$ (projection on the I coordinate of the product) we obtain graphs Γ_t dual to $\mathcal{E}_1 \cap H_t$ (when t is a regular value of the height function in \mathcal{E}_1). As t increases, Γ_t changes by folds. The first fold will have to happen at the valence 4 vertex (folds at valence 2 vertices do not happen by incompressibility). We want to say that this fold replaces the valence 4 vertex by two of valence 3, reducing the problem to the previous case. It could happen that the fold is done along an edge determining a closed loop and, after that, the fold actually replaces the vertex by another one still with valence 4. We can solve this problem by sufficiently subdividing the edges of Γ (i.e., introducing valence 2 vertices in the interior of the edges).

We now give the more detailed proof in the 2-dimensional setting. The fact that one 0-handle has valence 4 means, in genus 2, that \mathcal{E}_0 consists of just two isotopy classes of discs. We want each of these isotopy classes to contain at least two distinct discs of \mathcal{E}_0 . This can be easily achieved by *splittings* (see [Oer02] or the dual *subdivision* of [BH92]).

Since λ is minimal, Λ has the incompressibility property. In particular $\mathcal{E}_1 - \mathring{H}_0$ is incompressible in $H_1 - \mathring{H}_0$. We then have a height function in $H_1 - \mathring{H}_0$ with respect to which we may suppose that \mathcal{E}_1 is in Morse position having just saddles as critical points, no pair in the same level (see [Oer02]). Let t be a bit greater than the first critical value and consider the intermediate $H_0 \subsetneq H_t \subsetneq H_1$. Then $\mathcal{E}_t = \mathcal{E}_1 \cap H_t$ is a system of discs for H_t . It determines a handle decomposition \mathcal{H}_t . By regarding H_t as a “new H_0 ” and following the construction of the invariant measures laminations, we obtain the same original lamination and growth: $(\Lambda_t, \mu_t) = (\Lambda, \mu)$, $(\Omega_t, \nu_t) = (\Omega, \nu)$ and $\lambda_t = \lambda$ (see [Oer02] for details).

We claim that \mathcal{E}_t contains three distinct isotopy classes of discs. To see this we identify H_t with H_0 through the product structure and note that \mathcal{E}_t is obtained from $\mathcal{E}_0 \subseteq H_0$ by replacing two distinct discs by their band sum. This sum is done along a band contained in a 0-handle. If such a 0-handle had valence 2 then the band sum would join two parallel discs, contradicting incompressibility in $H_1 - \mathring{H}_0$, so the band is contained in the 0-handle with valence 4. But the band sum of discs in a 0-handle with valence greater than 3 yields a disc in a new isotopy class. Now recall that we chose \mathcal{E}_0 to have at least two discs in each class. Therefore a single band sum will preserve at least one disc in each of these original classes, proving the claim that \mathcal{E}_t contains three classes of discs.

Now that \mathcal{E}_t contains three distinct isotopy classes of discs, all its 0-handles have valence 3 or 2. Therefore, by Proposition 3.6, $\Lambda_t = \Lambda$ is

tight. □

Remark. There is another interesting point of view, from which we also sketch the proof here. It uses the disc complex of H . The disc complex $\mathcal{D}(H)$ of a handlebody H of genus 2 has dimension 2. Now consider the path $t \mapsto \Lambda \cap H_t$ on $\mathcal{D}(H)$ (here we use normalized transverse measures as barycentric coordinates). But $\Lambda \cap H_t$ determines a complete system of discs for any t so it has at least two isotopy classes of discs. Therefore the path $t \mapsto \Lambda \cap H_t$ never intersects the 0-skeleton of $\mathcal{D}(H)$. But the path cannot be confined to a single 1-simplex, so it has to intersect the interior of some 2-simplex of $\mathcal{D}(H)$, which corresponds to three distinct isotopy classes of weighted discs. The argument is finished as before.

3.4 Tightness implies minimality.

The next goal is to prove Theorem 3.5. For that, we need a technical result that follows below. If $G, G' \subseteq H$ are embedded graphs we shall say that G' follows G if $G' \subseteq F_G$, where F_G is a fixed fibered neighborhood of G , the vertices of G' are contained in the union of the neighborhoods of vertices of G and the edges of G' are transverse to the fibers of F_G over the edges.

Now suppose that $g: H \rightarrow H$ is an automorphism and that $g(G)$ follows G . In this case we define an incidence matrix $N = N_G(g)$ by $n_{ij} = |g(e_i) \cap F(e_j)|$, where $F(e_j) \subseteq F_G$ is the fibered neighborhood over the edge e_j . If N is irreducible we say that its Perron-Frobenius eigenvalue $\lambda_G = \lambda_G(g)$ is the growth of g on G .

In the following we assume that $f: H \rightarrow H$ and handle decomposition \mathcal{H}_0 of H_0 are fixed. We consider the corresponding disc system \mathcal{E}_0 , dual graph Γ_0 , laminations (Λ, μ) , (Ω, ν) and growth rate λ .

Proposition 3.15. *Suppose that there exists g isotopic to f^{-1} and graph $G \subseteq H$ such that $g(G)$ follows G with $\lambda_G < \lambda$. If Γ_0 is isotopic to a graph Γ'_0 which follows G then Λ is not tight.*

Proof. As usual, the measured lamination (Ω, ν) determines the standard weighted handle decomposition $(\mathcal{H}_0, \hat{\nu})$ of H_0 . We use $(\mathcal{H}_0, \hat{\nu})$ and f^{-n} to induce a weighted handle decomposition $(\mathcal{H}_{-n}, \hat{\nu})$ in $H_{-n} = f^{-n}(H_0)$ (i.e., the weight in the 1-handle $f^{-n}(e_i)$ of H_{-n} is $\hat{\nu}_i$).

Let E_i be the co-core of the 1-handle e_i of \mathcal{H}_0 . By the eigenvalue property of $\hat{\nu}$, for any $n > 0$

$$\frac{E_i \bullet (\mathcal{H}_{-n}, \hat{\nu})}{\lambda^n} = \hat{\nu}_i.$$

The goal of the argument is to find a disc Δ isotopic to E_i (rel ∂E_i) such that, for some $N > 0$,

$$\frac{\Delta \bullet (\mathcal{H}_{-N}, \hat{\nu})}{\lambda^N} < \hat{\nu}_i. \quad (3)$$

Since E_i may be chosen as a leaf of $\Lambda \cap H_0$ and $\Delta \bullet (\Omega, \nu) = \frac{\Delta \bullet (\mathcal{H}_{-N}, \hat{\nu})}{\lambda^N}$, such a Δ will then be a tightening disc.

The rough strategy is to isotope H_0 into the fibered neighborhood F_G of G and iterate g . Since its growth is smaller, the number of components of the intersection with the disc E_i will grow more slowly than originally, what will yield a tightening disc. We shall develop this idea more precisely.

A big part of the proof consists of certain constructions, as follows. We choose an isotopy h taking Γ_0 to Γ'_0 . We now consider H_0 as a neighborhood of Γ_0 and can obtain an isotopic $H'_0 \subseteq F_G$. The weighted handle decomposition of H_0 determines a handle decomposition $(\mathcal{H}'_0, \hat{\nu})$ of H'_0 . Also, the fibered structure of F_G determines a handle decomposition \mathcal{G} of F_G in the natural way (i.e., neighborhoods of vertices correspond to 0-handles and fibered neighborhoods over the edges to 1-handles). By adjusting h we can assume further that \mathcal{H}'_0 and \mathcal{G} are *compatible* in the following sense. Any dual disc of a 1-handle of \mathcal{G} intersects H'_0 in dual discs of \mathcal{H}'_0 .

Now use $(\mathcal{H}'_0, \hat{\nu})$ to induce a system of weights on \mathcal{G} in the following way. Let e_0, \dots, e_l be the 1-handles of \mathcal{G} and, for each i , let D_i be a dual disc of e_i . Define

$$(\hat{\nu}_G)_i = D'_i \bullet (\mathcal{H}'_0, \hat{\nu}). \quad (4)$$

Such a $\hat{\nu}_G$ is well defined: the way the handles of \mathcal{G} intersect those of \mathcal{H}'_0 assures that (4) above makes sense and does not depend on the choice of disc dual to e_i . This defines a weighted decomposition $(\mathcal{G}, \hat{\nu}_G)$.

We recall g from the hypotheses of the lemma. The decomposition \mathcal{G} determines a decomposition \mathcal{G}_{-1} of $g(F_G)$. We can adjust g so that \mathcal{G} and \mathcal{G}_{-1} are compatible (in particular, $g(F_G) \subseteq F_G$). Through g^n we can define decompositions \mathcal{G}_{-n} of $g^n(F_G)$. It is clear that these decompositions are automatically compatible.

We now define $H'_{-n} = g^n(H'_0)$, also with weighted handle structure $(\mathcal{H}'_{-n}, \hat{\nu})$ defined through g^n . Clearly $(\mathcal{H}'_{-n}, \hat{\nu})$ is isotopic to $(\mathcal{H}_{-n}, \hat{\nu})$. It is also clear that if $-m \leq -n$ then \mathcal{G}_{-n} and \mathcal{H}'_{-m} are compatible.

Recall the weight system $\hat{\nu}_G$ defined in (4) above. Now define weighted handle decompositions $(\mathcal{G}_{-n}, \hat{\nu}_G)$. For a general surface S the construction implies that (assuming that both sides makes sense):

$$S \bullet (\mathcal{H}'_{-n}, \hat{\nu}) = S \bullet (\mathcal{G}_{-n}, \hat{\nu}_G). \quad (5)$$

To complete the constructions necessary in this proof, let E_i be a disc in the original system \mathcal{E}_0 . For a technical reason we extend it through the product structure in $H - \mathring{H}_0$ to a disc $(E, \partial E) \subseteq (H, \partial H)$. It is clear that $E \bullet (\Omega, \nu) = E_i \bullet (\Omega, \nu) = \hat{\nu}_i$. We isotope E and also assume that $E \cap F_G$ consists of dual discs of \mathcal{G} .

Recall that $\lambda_G = \lambda_G(g)$ is the growth of g on G . Clearly the sequence

$$n \mapsto \frac{E \bullet (\mathcal{G}_{-n}, v_G)}{(\lambda_G)^n}$$

is bounded. But $\lambda > \lambda_G$, therefore

$$\frac{E \bullet (\mathcal{G}_{-n}, v_G)}{\lambda^n} \rightarrow 0$$

and then, for some N ,

$$\frac{E \bullet (\mathcal{G}_{-N}, v_G)}{\lambda^N} < \hat{\nu}_i. \quad (6)$$

By (5) and (6):

$$\frac{E \bullet (\mathcal{H}'_{-N}, \hat{\nu})}{\lambda^N} < \hat{\nu}_i. \quad (7)$$

Now note that there is an ambient isotopy $h': H \rightarrow H$ such that $h'(\mathcal{H}'_{-N}, \hat{\nu}) = (\mathcal{H}_{-N}, \hat{\nu})$. By applying h' to $(\mathcal{H}'_{-N}, \hat{\nu})$ follows from (7) above that

$$\frac{h'(E) \bullet (\mathcal{H}_{-N}, \hat{\nu})}{\lambda^N} < \hat{\nu}_i. \quad (8)$$

We can choose h' restricting to the identity at ∂H , so $\Delta' = h'(E)$ has the property that $\partial \Delta' = \partial E$.

Now recall that E is the extension to H of the co-core E_i of a handle of \mathcal{H}_0 . We can use the product structure in $H - \mathring{H}_0$ to obtain a disc $\Delta \subseteq H_0$ from Δ' . Clearly $\partial \Delta = \partial E_i$, E_i may be chosen as a leaf of $\Lambda \cap H_0$ and $\Delta \bullet (\mathcal{H}_{-N}, \hat{\nu}) = \Delta' \bullet (\mathcal{H}_{-N}, \hat{\nu})$. Therefore (3) follows from (8), showing Δ as a tightening disc. \square

(Theorem 3.5). *If Λ is tight then λ is minimal.*

Proof. It is a corollary of Proposition 3.15 above. We prove the contrapositive, so assume that λ is not minimal. Then there exists another structure $\hat{\mathcal{H}}_0$ for some \hat{H}_0 and representative \hat{f} for which the growth rate $\hat{\lambda}$ is less than λ . We consider the graph $\hat{\Gamma}_0$ corresponding to $\hat{\mathcal{H}}_0$. It is direct that 1) $(\hat{f})^{-1}(\hat{\Gamma}_0)$ follows Γ_0 (with $\lambda_{\hat{\Gamma}_0} = \hat{\lambda} < \lambda$) and 2) that Γ_0 is isotopic to a $\Gamma'_0 = \hat{\Gamma}_0$. By Proposition 3.15 Λ is not tight, completing the proof. \square

Remark. Theorem 3.5 can be used to find the minimal growth of some actual examples (see [Car03]).

3.5 Applications: comparing growth rates.

In this subsection we prove some results on the growth rate of an irreducible automorphism with respect to a tight lamination.

The following is a direct corollary of Theorem 3.5.

Corollary 3.16. *If Λ is tight and $n > 0$ then $\lambda_{\min}(f^n) = \lambda^n$.*

Proof. Since $f^n(\Lambda, \mu) = (\Lambda, \lambda^n \mu)$ and $f^n(\Omega, \nu) = (\Omega, \lambda^{-n} \nu)$, the growth of f^n with respect to the handle decomposition that determine Λ and Ω is λ^n . But Λ is tight, therefore λ^n is minimal. \square

In [Oer02] the question is posed whether there is any relation between the growth rate $\lambda = \lambda(f)$ of a generic automorphism and the growth rate λ_∂ of the pseudo-Anosov restriction $\partial f = f|_{\partial H}$ to the boundary. The following is another corollary of Proposition 3.15.

Corollary 3.17. *Let $f: H \rightarrow H$ be a generic automorphism. If Λ is tight then $\lambda \leq \lambda_\partial$.*

Proof. We prove the contrapositive, so suppose that $\lambda_\partial < \lambda$.

Let $g = f^{-1}$ and let τ be a stable train-track of ∂g (we regard τ simply as a graph, with switches for vertices and branches for edges). Let $F_\tau \subseteq H$ be a fibered neighborhood of τ and isotope g so that $g(\tau)$ follows τ . It is clear the Γ_0 is isotopic to a graph Γ'_0 which follows τ . Indeed, Γ_0 is boundary parallel and τ fills ∂H . Proposition 3.15 completes the proof. \square

Corollary 3.18. *The (minimal) growth rate of a generic automorphism $f: H \rightarrow H$ of a handlebody of genus 2 is less than or equal to the growth rate of the pseudo-Anosov $\partial f = f|_{\partial H}$.*

Proof. Use Corollary 3.17 and Corollary 3.14. \square

References

- [BH92] M. Bestvina and M. Handel. Train tracks and automorphisms of free groups. *Ann. of Math.*, 135:1–51, 1992.

- [BH95] M. Bestvina and M. Handel. Train-tracks for surface homeomorphisms. *Topology*, 34(1):109–140, 1995.
- [Bon83a] F. Bonahon. Cobordism of automorphisms of surfaces. *Ann. Sci. École Norm. Sup. (4)*, 16(2):237–270, 1983.
- [Bon83b] F. Bonahon. Ribbon fibered knots, cobordism of surface diffeomorphisms and pseudo-Anosov diffeomorphisms. *Math. Proc. Cambridge Philos. Soc.*, 94(2):235–251, 1983.
- [Car03] Leonardo N. Carvalho. *Generic Automorphisms of Handlebodies*. PhD thesis, Rutgers University–Newark, 2003.
- [CB88] Andrew J. Casson and Steven A. Bleiler. *Automorphisms of surfaces after Nielsen and Thurston*, volume 9 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1988.
- [FL80] A. Fathi and F. Laudenbach. Difféomorphismes pseudo-Anosov et décomposition de Heegaard. *C. R. Acad. Sc. Paris*, 291:423–425, 1980. Série A.
- [FLP79] A. Fathi, F. Laudenbach, and V. Poenaru. *Travaux de Thurston sur les Surfaces*, volume 66 of *Astérisque*. Soc. Math. France, 1979.
- [HT85] M. Handel and W. Thurston. New proofs of some results of Nielsen. *Adv. in Math.*, 56(2):173–191, 1985.
- [Joh79] K. Johannson. *Homotopy equivalences of 3-manifolds with boundaries*. Number 761 in *Lecture Notes in Mathematics*. Springer-Verlag, 1979.
- [JS79] W. Jaco and P. Shalen. Seifert fibered spaces in 3-manifolds. *Mem. Amer. Math. Soc.*, 220:1–192, 1979.
- [Lon88] D. D. Long. Bounding laminations. *Duke Math. J.*, 56(1):1–16, 1988.
- [Nie86a] J. Nielsen. *Investigations in the topology of closed orientable surfaces, I*, pages 223–341. In *Contemporary Mathematicians* [Nie86d], 1986. Translation by John Stillwell of: Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen. [Acta Math., 50 (1927), 189–358].

- [Nie86b] J. Nielsen. *Investigations in the topology of closed orientable surfaces, II*, pages 348–400. In *Contemporary Mathematicians* [Nie86d], 1986. Translation by John Stillwell of: Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen, II. [Acta Math., 53 (1929), 1–76].
- [Nie86c] J. Nielsen. *Investigations in the topology of closed orientable surfaces, III*, pages 401–459. In *Contemporary Mathematicians* [Nie86d], 1986. Translation by John Stillwell of: Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen, III. [Acta Math., 58 (1932), 87–167].
- [Nie86d] J. Nielsen. *Jakob Nielsen: collected mathematical papers. Vol. 1*. Contemporary Mathematicians. Birkhäuser Boston Inc., Boston, MA, 1986. Edited and with a preface by Vagn Lundsgaard Hansen.
- [Oer02] U. Oertel. Automorphisms of 3-dimensional handlebodies. *Topology*, 41:363–410, 2002.
- [Pen88] R. Penner. A construction of pseudo-Anosov homeomorphisms. *Trans. Amer. Math. Soc.*, 310(1):179–197, 1988.
- [Sen73] E. Seneta. *Non-negative Matrices*. Halsted Press, 1973.
- [Thu88] W. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc.*, 19:417–431, 1988.