

The effective potential and transshipment in thermodynamic formalism at temperature zero

Eduardo Garibaldi*

Departamento de Matemática
Universidade Estadual de Campinas
13083-859 Campinas - SP, Brasil
garibaldi@ime.unicamp.br

Artur O. Lopes†

Instituto de Matemática
Universidade Federal do Rio Grande do Sul
91509-900 Porto Alegre - RS, Brasil
arturoscar.lopes@gmail.com

August 5, 2010

Abstract

Denote the points in $\{1, 2, \dots, r\}^{\mathbb{Z}} = \{1, 2, \dots, r\}^{\mathbb{N}} \times \{1, 2, \dots, r\}^{\mathbb{N}}$ by $(\mathbf{y}^*, \mathbf{x})$. Given a Lipschitz continuous observable $A : \{1, 2, \dots, r\}^{\mathbb{Z}} \rightarrow \mathbb{R}$, we define the map $\mathcal{G}^+ : \mathcal{H} \rightarrow \mathcal{H}$ by

$$\mathcal{G}^+(\phi)(\mathbf{y}^*) = \sup_{\mu \in \mathcal{M}_\sigma} \left[\int_{\{1, 2, \dots, r\}^{\mathbb{N}}} (A(\mathbf{y}^*, \mathbf{x}) + \phi(\mathbf{x})) d\mu(\mathbf{x}) + h_\mu(\sigma) \right],$$

where:

- σ is the left shift map acting on $\{1, 2, \dots, r\}^{\mathbb{N}}$;
- \mathcal{M}_σ denotes the set of σ -invariant Borel probabilities;
- $h_\mu(\sigma)$ indicates the Kolmogorov-Sinai entropy;
- \mathcal{H} is the Banach space of Lipschitz real-valued functions on $\{1, 2, \dots, r\}^{\mathbb{N}}$.

We show there exist a unique $\phi^+ \in \mathcal{H}$ and a unique $\lambda^+ \in \mathbb{R}$ such that

$$\mathcal{G}^+(\phi^+) = \phi^+ + \lambda^+.$$

We say that ϕ^+ is the effective potential associated to A . This also defines a family of σ -invariant Borel probabilities $\mu_{\mathbf{y}^*}$ on $\{1, 2, \dots, r\}^{\mathbb{N}}$, indexed by the points $\mathbf{y}^* \in \{1, 2, \dots, r\}^{\mathbb{N}}$. Finally, for A fixed and for variable positive real values β , we consider the same problem for the Lipschitz observable βA . We investigate then the asymptotic limit when $\beta \rightarrow \infty$ of the effective potential (which depends now on β) as well as the above family of probabilities. We relate the limit objects with an ergodic version of Kantorovich transshipment problem. In statistical mechanics $\beta \propto 1/T$, where T is the absolute temperature. In this way, we are also analyzing the problem related to the effective potential at temperature zero.

Keywords: thermodynamic formalism, effective potential, transshipment, Gibbs state, additive eigenvalue, maximizing probabilities.

MSC: 37A05, 37A35, 37A60, 37D35, 49K27, 49Q20, 82B05, 90C05, 90C50.

*Supported by PROCAD UNICAMP-UFRGS 162-2007.

†Partially supported by CNPq, PRONEX – Sistemas Dinâmicos, INCT em Matemática, and beneficiary of CAPES financial support (PROCAD UFRGS-IMPACT and PROCAD UNICAMP-UFRGS).

1 Introduction

Our purpose is to propose, in a rigorous mathematical way, a description of the main features of what could be called in statistical mechanics the *effective potential formalism* for long range interactions. In this way, we are able to present a family of *effective probabilities*, each one corresponding to a Gibbs state in the sense of Ruelle's thermodynamic setting [13]. We also consider the limit behavior of this family of probabilities when the temperature goes to zero. In this case, we relate our analysis with a kind of ergodic Kantorovich transshipment problem. We point out that in the classical transport theory [14, 16, 17] there is no assumptions about invariant probabilities playing a role in the problem.

Actually our framework will be more general than Bernoulli shifts. We will develop the theory using one-sided topologically transitive subshifts of finite type defined by symmetric transition matrices.

Hence let $\mathbf{M} : \{1, \dots, r\} \times \{1, \dots, r\} \rightarrow \{0, 1\}$ be an irreducible transition matrix. One has naturally two subshifts associated to such a matrix. We can introduce the standard subshift of finite type

$$\Sigma_{\mathbf{M}} = \left\{ (x_0, x_1, \dots) \in \{1, \dots, r\}^{\mathbb{Z}^+} : \mathbf{M}(x_j, x_{j+1}) = 1 \right\},$$

as well as the dual subshift of finite type

$$\Sigma_{\mathbf{M}^T}^* = \left\{ (\dots, x_1, x_0) \in \{1, \dots, r\}^{\mathbb{Z}^-} : \mathbf{M}^T(x_j, x_{j+1}) = 1 \right\}.$$

As topological spaces, both subshifts are always compact metrizable spaces. We suppose henceforth that the matrix \mathbf{M} is symmetric. So we have a canonical homeomorphism $\mathbf{x} = (x_0, x_1, \dots) \in \Sigma_{\mathbf{M}} \mapsto \mathbf{x}^* = (\dots, x_1, x_0) \in \Sigma_{\mathbf{M}}^*$.

Given $\Lambda \in (0, 1)$, we equip as usual $\Sigma_{\mathbf{M}}$ with the metric $d(\mathbf{x}, \mathbf{y}) = \Lambda^k$, where $\mathbf{x} = (x_0, x_1, \dots), \mathbf{y} = (y_0, y_1, \dots) \in \Sigma_{\mathbf{M}}$ and $k = \min\{j : x_j \neq y_j\}$. Hence, for $\mathbf{x}^*, \mathbf{y}^* \in \Sigma_{\mathbf{M}}^*$, we just set $d^*(\mathbf{x}^*, \mathbf{y}^*) := d(\mathbf{x}, \mathbf{y})$.

Let σ be the left shift map acting on $\Sigma_{\mathbf{M}}$ and let σ^* be the right shift map acting on $\Sigma_{\mathbf{M}}^*$, namely,

$$\sigma(x_0, x_1, x_2, \dots) = (x_1, x_2, \dots) \quad \text{and} \quad \sigma^*(\dots, x_2, x_1, x_0) = (\dots, x_2, x_1).$$

Clearly, $* \circ \sigma = \sigma^* \circ *$. Furthermore, since \mathbf{M} is irreducible, notice that the dynamics $(\Sigma_{\mathbf{M}}, \sigma)$ is transitive – and consequently the conjugated dynamical system $(\Sigma_{\mathbf{M}}^*, \sigma^*)$ too.

Let $C^0(\Sigma_{\mathbf{M}})$ and $C^0(\Sigma_{\mathbf{M}}^*)$ denote the spaces of continuous real-valued functions on respectively $\Sigma_{\mathbf{M}}$ and $\Sigma_{\mathbf{M}}^*$, both equipped with the topology of uniform convergence. Thus, we can obtain from the previous homeomorphism an isometry $* : C^0(\Sigma_{\mathbf{M}}) \rightarrow C^0(\Sigma_{\mathbf{M}}^*)$ writing $f^*(\mathbf{x}^*) := f(\mathbf{x})$ for every function $f \in C^0(\Sigma_{\mathbf{M}})$. This fact allows us to make the identification $C^0(\Sigma_{\mathbf{M}}) \simeq C^0(\Sigma_{\mathbf{M}}^*)$.

The same isometric property is verified for either Hölder or Lipschitz continuous real-valued functions. Since one can simply incorporate the Hölder exponent into the distance, we remark that to work with the Lipschitz class does not lead to loss of generality. Therefore, \mathcal{H} will denote in this article the Banach space of

Lipschitz continuous real-valued functions on either $\Sigma_{\mathbf{M}}$ or $\Sigma_{\mathbf{M}}^*$, equipped with the norm $\|\cdot\|_{\mathcal{H}} := \|\cdot\|_0 + \text{Lip}(\cdot)$, where $\|\cdot\|_0$ denotes the uniform norm and

$$\text{Lip}(\phi) = \sup_{d(\mathbf{x}, \mathbf{y}) > 0} \frac{|\phi(\mathbf{x}) - \phi(\mathbf{y})|}{d(\mathbf{x}, \mathbf{y})} = \sup_{d^*(\mathbf{x}^*, \mathbf{y}^*) > 0} \frac{|\phi^*(\mathbf{x}^*) - \phi^*(\mathbf{y}^*)|}{d^*(\mathbf{x}^*, \mathbf{y}^*)} = \text{Lip}(\phi^*).$$

Using the standard subshift $\Sigma_{\mathbf{M}}$ and its dual $\Sigma_{\mathbf{M}}^*$, one may easily introduce its natural invertible extension $(\hat{\Sigma}_{\mathbf{M}}, \hat{\sigma})$:

$$\hat{\Sigma}_{\mathbf{M}} = \{(\mathbf{y}^*, \mathbf{x}) \in \Sigma_{\mathbf{M}}^* \times \Sigma_{\mathbf{M}} : \mathbf{M}(y_0, x_0) = 1\},$$

$$\hat{\sigma}(\dots, y_1, y_0 | x_0, x_1, \dots) = (\dots, y_0, x_0 | x_1, x_2, \dots).$$

Denote by \mathcal{M}_{σ} the weak* compact and convex set of σ -invariant Borel probability measures. For any $\mu \in \mathcal{M}_{\sigma}$, let $h_{\mu}(\sigma)$ indicate the Kolmogorov-Sinai entropy.

Definition 1. Given a Lipschitz continuous function $A : \hat{\Sigma}_{\mathbf{M}} \rightarrow \mathbb{R}$, we consider then the map $\mathcal{G}^+ = \mathcal{G}_A^+ : \mathcal{H} \rightarrow \mathcal{H}$ defined¹ by

$$\mathcal{G}^+(\phi)(\mathbf{y}^*) = \sup_{\mu \in \mathcal{M}_{\sigma}} \left[\int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}^*, \mathbf{x}) + \phi(\mathbf{x})) d\mu(\mathbf{x}) + h_{\mu}(\sigma) \right]$$

It is not difficult to see that $\text{Lip}(\mathcal{G}^+(\phi)) \leq \|A\|_0 + \text{Lip}(A)$ for all $\phi \in \mathcal{H}$. Furthermore, thanks to the characterization via variational principle of the topological pressure $P_{TOP} : \mathcal{H} \rightarrow \mathbb{R}$, that is,

$$P_{TOP}(\phi) = \max_{\mu \in \mathcal{M}_{\sigma}} \left[\int_{\Sigma_{\mathbf{M}}} \phi(\mathbf{x}) d\mu(\mathbf{x}) + h_{\mu}(\sigma) \right] \quad \forall \phi \in \mathcal{H},$$

we immediately get

$$\mathcal{G}^+(\phi)(\mathbf{y}^*) = P_{TOP}(A(\mathbf{y}^*, \cdot) + \phi).$$

In particular, thanks to the Ruelle-Perron-Frobenius Theorem, for each $\mathbf{y}^* \in \Sigma_{\mathbf{M}}^*$, there exists a unique probability $\mu_{\mathbf{y}^*} \in \mathcal{M}_{\sigma}$ (the equilibrium state associated to $A(\mathbf{y}^*, \cdot) + \phi \in \mathcal{H}$) achieving the supremum in the definition of the value $\mathcal{G}^+(\phi)(\mathbf{y}^*)$.

As a physical motivation to analyze the above problem, we mention the paper by W. Chou and R. Griffiths [3]. They study ground states of one-dimensional systems, in particular of a very common model in solid state physics: the Frenkel-Kontorova model (specific applications are presented in section VI). They consider a certain model which depends on temperature and which has a natural potential. But due to interaction and temperature, there exists another potential, called the *effective potential*, which plays the essential role in the problem. The limit when temperature goes to zero is considered in section III B. Their expression (3.16)

¹Notice that a more rigorous definition would consider $\int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}^*, \mathbf{x})\mathbf{M}(\mathbf{y}^*, \mathbf{x}) + \phi(\mathbf{x})) d\mu(\mathbf{x})$, where $\mathbf{M}(\mathbf{y}^*, \mathbf{x}) := \mathbf{M}(y_0, x_0)$ for any point $(\mathbf{y}^*, \mathbf{x}) = (\dots, y_1, y_0 | x_0, x_1, \dots)$. We prefer to simplify the notation.

may be seen as a min-plus version of the one in our main theorem. See also [1, 2, 4] for more details on additive eigenvalue problems.

The entropy penalization method was considered in [8] and [7] (see the main properties on these references) in the setting of Aubry-Mather theory. In [10], questions also related to the article of Chou and Griffiths were analyzed in the context of Markov chains on the interval. The problem we consider here has similarities. Nevertheless, we point out that our entropy is Kolmogorov-Sinai entropy, which has a dynamical character. As we said before, our setting is the one of thermodynamic formalism [13]. Finally, the relation of the effective action problem with the ergodic Kantorovich transshipment problem (see section 3), as far as we know, is completely new.

As another physical motivation for the study of the above problem, we mention section 2.5 of Salmhofer's book [15]. The function ϕ plays there the role of a chemical potential. In [15], using another notation, the expression (2.103) of the effective action for the interaction $-\lambda V$ and the propagator C

$$\int e^{-\lambda V(\phi) + (C^{-1}\psi, \phi)} d\mu_C(\phi), \quad \text{for a fixed } \psi,$$

should be read, under our notation, as

$$\int e^{\phi(\mathbf{x}) + A(\mathbf{y}^*, \mathbf{x})} d\mu_C(\mathbf{x}), \quad \text{for a fixed } \mathbf{y}^*.$$

Note that the above probability maximizes pressure when one considers, for each fixed \mathbf{y}^* , the observable $A(\mathbf{y}^*, \cdot) + \phi$, and the corresponding variational problem where the entropy $h(\nu|\mu_C)$ of a given ν is considered relative to a fixed initial probability μ_C . As we are in the framework of thermodynamic formalism, we do not consider relative entropy, but Kolmogorov-Sinai entropy.

Our main result can be stated as follows.

Theorem 1. *Suppose $A : \hat{\Sigma}_{\mathbf{M}} \rightarrow \mathbb{R}$ is a Lipschitz continuous observable. Then there exist a unique function $\phi^+ \in \mathcal{H}$ (up to an additive constant) and a unique constant $\lambda^+ \in \mathbb{R}$ such that*

$$\mathcal{G}^+(\phi^+) = \phi^+ + \lambda^+.$$

We point out that [7, 8, 10] consider a similar problem but for the so called entropy penalization method. The proof of this theorem will be presented in the end of the paper. Obviously the function ϕ^+ and the constant λ^+ in the previous statement depend on A .

Definition 2. *Given a Lipschitz continuous observable $A : \hat{\Sigma}_{\mathbf{M}} \rightarrow \mathbb{R}$, we say that a constant $\lambda^+ \in \mathbb{R}$ is the effective constant for A if there exists a function $\phi^+ \in \mathcal{H}$ such that*

$$\mathcal{G}^+(\phi^+) = \phi^+ + \lambda^+.$$

Any such a function ϕ^+ is called a (forward) effective potential for A .

Definition 3. Given a Lipschitz continuous observable $A : \hat{\Sigma}_{\mathbf{M}} \rightarrow \mathbb{R}$ and a point $\mathbf{y}^* \in \Sigma_{\mathbf{M}}^*$, we say that the unique σ -invariant probability $\mu_{\mathbf{y}^*} = \mu_{\mathbf{y}^*, A}$ on $\Sigma_{\mathbf{M}}$ such that

$$\int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}^*, \mathbf{x}) + \phi^+(\mathbf{x})) d\mu_{\mathbf{y}^*}(\mathbf{x}) + h_{\mu_{\mathbf{y}^*}}(\sigma) = \phi^+(\mathbf{y}^*) + \lambda^+$$

is the effective probability for A at \mathbf{y}^* , where ϕ^+ and λ^+ are the effective ones associated to A . In this way, we get a family of Gibbs states on the variable \mathbf{x} indexed by \mathbf{y}^* .

For a fixed A as above, we consider a positive parameter β , the observable βA , and the corresponding ϕ_{β}^+ , λ_{β}^+ and $\{\mu_{\mathbf{y}^*, \beta A}\}_{\mathbf{y}^* \in \Sigma_{\mathbf{M}}^*}$. We investigate then the limit problem when $\beta \rightarrow \infty$, showing the existence (in the uniform topology) of accumulation Lipschitz functions for the family $\{\phi_{\beta}^+/\beta\}_{\beta > 0}$, characterizing the accumulation probabilities of $\{\mu_{\mathbf{y}^*, \beta A}\}_{\beta > 0}$ for each \mathbf{y}^* , and proving that λ_{β}^+/β converges (see section 2).

We remark at last that one could also consider the (backward) transformation $\mathcal{G}^- = \mathcal{G}_A^- : \mathcal{H} \rightarrow \mathcal{H}$ defined by

$$\mathcal{G}^-(\phi)(\mathbf{x}) = \sup_{\mu \in \mathcal{M}_{\sigma^*}} \left[\int_{\Sigma_{\mathbf{M}}^*} (A(\mathbf{y}^*, \mathbf{x}) + \phi(\mathbf{y}^*)) d\mu(\mathbf{y}^*) + h_{\mu}(\sigma^*) \right],$$

and all analogous results could be easily stated and similarly proved.

The structure of the paper is the following: in section 2 we discuss the thermodynamic properties of the effective objects, in section 3 we consider the ergodic Kantorovich transshipment problem (which appears in a natural way when the temperature goes to zero), and finally in section 4 we present the proof of the main theorem.

2 Thermodynamic formalism at temperature zero

The analysis of the thermodynamic formalism for a given observable A at temperature zero is, by definition, the study of the limit of Gibbs probabilities associated to A at temperature T , that is, for $\frac{1}{T} A$, when $T \rightarrow 0$. We introduce a parameter $\beta = \frac{1}{T}$, and we will analyze the Gibbs probabilities for βA , when $\beta \rightarrow \infty$.

From now on, \mathbf{y}^* is simply denoted by \mathbf{y} and we identify the spaces $\Sigma_{\mathbf{M}}$ and $\Sigma_{\mathbf{M}}^*$. For each real value β , we consider the map $\mathcal{G}_{\beta A}^+ : \mathcal{H} \rightarrow \mathcal{H}$ and the corresponding Lipschitz function ϕ_{β}^+ , the forward effective potential for βA , and the corresponding constant $\lambda_{\beta}^+ \in \mathbb{R}$. For each \mathbf{y} , we consider then the effective probability $\mu_{\mathbf{y}, \beta A}$ as before. In order to avoid a heavy notation we will drop the A and the $+$ in this section.

In this way, for each parameter β , we have the equation

$$\mathcal{G}_{\beta}(\phi_{\beta}) = \phi_{\beta} + \lambda_{\beta}.$$

Recall that, for each \mathbf{y} , we have $\mathcal{G}_{\beta}(\phi_{\beta})(\mathbf{y}) = P_{TOP}(\beta A(\mathbf{y}, \cdot) + \phi_{\beta})$, where the pressure is consider for the setting in the variable \mathbf{x} . Therefore, for each \mathbf{y} and β , one verifies

$$\phi_{\beta}(\mathbf{y}) + \lambda_{\beta} = P_{TOP}(\beta A(\mathbf{y}, \cdot) + \phi_{\beta}).$$

Proposition 2. *The family $\frac{\phi_\beta}{\beta}$ is equi-Lipschitz.*

Proof. For each pair of points \mathbf{y} and $\bar{\mathbf{y}}$, one has

$$|\phi_\beta(\mathbf{y}) - \phi_\beta(\bar{\mathbf{y}})| = |P_{TOP}(\beta A(\mathbf{y}, \cdot) + \phi_\beta) - P_{TOP}(\beta A(\bar{\mathbf{y}}, \cdot) + \phi_\beta)|.$$

The effective probability $\mu_{\mathbf{y},\beta}$ satisfies

$$P_{TOP}(\beta A(\mathbf{y}, \cdot) + \phi_\beta) = \int (\beta A(\mathbf{y}, \cdot) + \phi_\beta) d\mu_{\mathbf{y},\beta} + h_{\mu_{\mathbf{y},\beta}}(\sigma).$$

It is clear that $P_{TOP}(\beta A(\bar{\mathbf{y}}, \cdot) + \phi_\beta) \geq \int (\beta A(\bar{\mathbf{y}}, \cdot) + \phi_\beta) d\mu_{\mathbf{y},\beta} + h_{\mu_{\mathbf{y},\beta}}(\sigma)$.

Therefore, the inequality²

$$\begin{aligned} P_{TOP}(\beta A(\mathbf{y}, \cdot) + \phi_\beta) - P_{TOP}(\beta A(\bar{\mathbf{y}}, \cdot) + \phi_\beta) &\leq \\ &\leq \beta \sup_{\mathbf{x} \in \Sigma_{\mathbf{M}}} |A(\mathbf{y}, \mathbf{x})\mathbf{M}(\mathbf{y}, \mathbf{x}) - A(\bar{\mathbf{y}}, \mathbf{x})\mathbf{M}(\bar{\mathbf{y}}, \mathbf{x})| \end{aligned}$$

yields

$$P_{TOP}(\beta A(\mathbf{y}, \cdot) + \phi_\beta) - P_{TOP}(\beta A(\bar{\mathbf{y}}, \cdot) + \phi_\beta) \leq \beta(\|A\|_0 + \text{Lip}(A)) d(\mathbf{y}, \bar{\mathbf{y}}).$$

Interchanging the roles of \mathbf{y} and $\bar{\mathbf{y}}$ in the above reasoning, we get

$$|\phi_\beta(\mathbf{y}) - \phi_\beta(\bar{\mathbf{y}})| \leq \beta(\|A\|_0 + \text{Lip}(A)) d(\mathbf{y}, \bar{\mathbf{y}}),$$

and finally

$$\text{Lip}\left(\frac{\phi_\beta}{\beta}\right) \leq \|A\|_0 + \text{Lip}(A).$$

□

Remember that the effective potential is unique up to an additive constant. So we will consider the following condition: we fix a point $\mathbf{y}^0 \in \Sigma_{\mathbf{M}}^*$ and we assume that $\phi_\beta(\mathbf{y}^0) = 0$ for all β . Via subsequences $\beta_n \rightarrow \infty$, with $n \rightarrow \infty$, using the previous proposition, we get by the Arzela-Ascoli Theorem that there exists a continuous function $V : \Sigma_{\mathbf{M}}^* \rightarrow \mathbb{R}$ such that $V(\mathbf{y}^0) = 0$ and, in the uniform convergence,

$$\frac{\phi_{\beta_n}}{\beta_n} \rightarrow V.$$

Since $\text{Lip}(\phi_\beta/\beta) \leq \|A\|_0 + \text{Lip}(A)$ implies $\text{Lip}(V) \leq \|A\|_0 + \text{Lip}(A)$, the function V is actually Lipschitz continuous. Notice that, in principle, such a limit could depend on the chosen subsequence.

Proposition 3. *Suppose that in the uniform convergence*

$$\frac{\phi_{\beta_n}}{\beta_n} \rightarrow V,$$

²Recall footnote 1.

when $\beta_n \rightarrow \infty$. Let $\mu_{\mathbf{y},\beta_n}$ be the effective probability for the observable $\beta_n A$ at a fixed point \mathbf{y} . Then, any accumulation probability measure $\mu_{\mathbf{y}}^\infty \in \mathcal{M}_\sigma$ of the sequence $\mu_{\mathbf{y},\beta_n}$ is a maximizing probability for $A(\mathbf{y}, \cdot) + V$, that is,

$$\int (A(\mathbf{y}, \cdot) + V) d\mu_{\mathbf{y}}^\infty = \max_{\mu \in \mathcal{M}_\sigma} \int (A(\mathbf{y}, \cdot) + V) d\mu.$$

Proof. Take any σ -invariant probability μ . Thus

$$\begin{aligned} \int (\beta_n A(\mathbf{y}, \cdot) + \phi_{\beta_n}) d\mu + h_\mu(\sigma) &\leq P_{TOP}(\beta_n A(\mathbf{y}, \cdot) + \phi_{\beta_n}) \\ &= \int (\beta_n A(\mathbf{y}, \cdot) + \phi_{\beta_n}) d\mu_{\mathbf{y},\beta_n} + h_{\mu_{\mathbf{y},\beta_n}}(\sigma). \end{aligned}$$

Given an accumulation probability measure $\mu_{\mathbf{y}}^\infty$ of the sequence $\mu_{\mathbf{y},\beta_n}$, from

$$\int \left(A(\mathbf{y}, \cdot) + \frac{\phi_{\beta_n}}{\beta_n} \right) d\mu + \frac{1}{\beta_n} h_\mu(\sigma) \leq \int \left(A(\mathbf{y}, \cdot) + \frac{\phi_{\beta_n}}{\beta_n} \right) d\mu_{\mathbf{y},\beta_n} + \frac{1}{\beta_n} h_{\mu_{\mathbf{y},\beta_n}}(\sigma),$$

we get the inequality

$$\int (A(\mathbf{y}, \cdot) + V) d\mu \leq \int (A(\mathbf{y}, \cdot) + V) d\mu_{\mathbf{y}}^\infty.$$

Therefore, $\mu_{\mathbf{y}}^\infty$ is a maximizing probability for $A(\mathbf{y}, \cdot) + V$. \square

Proposition 4. Assume that in the uniform convergence

$$\frac{\phi_{\beta_n}}{\beta_n} \rightarrow V,$$

when $\beta_n \rightarrow \infty$. Suppose also that $\mu_{\mathbf{y},\beta_n}$, the effective probability for the observable $\beta_n A$ at a fixed point \mathbf{y} , converges in the weak* topology to $\mu_{\mathbf{y}}^\infty \in \mathcal{M}_\sigma$. Then,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\lambda_{\beta_n}}{\beta_n} &= \max_{\mu \in \mathcal{M}_\sigma} \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y})) d\mu(\mathbf{x}) \\ &= \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y})) d\mu_{\mathbf{y}}^\infty(\mathbf{x}). \end{aligned}$$

Proof. As for any given point \mathbf{y}

$$\phi_{\beta_n}(\mathbf{y}) + \lambda_{\beta_n} = \int (\beta_n A(\mathbf{y}, \cdot) + \phi_{\beta_n}) d\mu_{\mathbf{y},\beta_n} + h_{\mu_{\mathbf{y},\beta_n}}(\sigma),$$

then dividing this expression by β_n , taking limit, and using last proposition, we immediately get the claim. \square

We point out that obviously the limit function $V \in \mathcal{H}$ and the limit measure $\mu_{\mathbf{y}}^\infty \in \mathcal{M}_\sigma$ may depend on the particular choice of the sequence β_n . Notice the previous proposition guarantees that the value $\int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \cdot) + V - V(\mathbf{y})) d\mu_{\mathbf{y}}^\infty$ does not depend on the point \mathbf{y} . Actually it does not even depend on the function V .

Proposition 5. *Suppose that in the uniform convergence*

$$\frac{\phi_{\beta_n}}{\beta_n} \rightarrow V \quad \text{and} \quad \frac{\phi_{\bar{\beta}_n}}{\bar{\beta}_n} \rightarrow \bar{V},$$

when $\beta_n, \bar{\beta}_n \rightarrow \infty$. Then, for all point \mathbf{y} ,

$$\begin{aligned} \max_{\mu \in \mathcal{M}_\sigma} \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y})) \, d\mu(\mathbf{x}) &= \\ &= \max_{\mu \in \mathcal{M}_\sigma} \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + \bar{V}(\mathbf{x}) - \bar{V}(\mathbf{y})) \, d\mu(\mathbf{x}). \end{aligned}$$

Proof. Passing to subsequences if necessary, we use the previous proposition to define

$$c := \lim_{n \rightarrow \infty} \frac{\lambda_{\beta_n}}{\beta_n} \quad \text{and} \quad \bar{c} := \lim_{n \rightarrow \infty} \frac{\lambda_{\bar{\beta}_n}}{\bar{\beta}_n}.$$

Notice that, again from proposition 4,

$$\begin{aligned} V(\mathbf{y}) + c &= \max_{\mu \in \mathcal{M}_\sigma} \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x})) \, d\mu(\mathbf{x}) \quad \text{and} \\ \bar{V}(\mathbf{y}) + \bar{c} &= \max_{\mu \in \mathcal{M}_\sigma} \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + \bar{V}(\mathbf{x})) \, d\mu(\mathbf{x}), \end{aligned}$$

for all point \mathbf{y} . Let \mathbf{y}^0 be a global maximum point for $V - \bar{V}$. Consider then a probability $\mu_0 \in \mathcal{M}_\sigma$ such that

$$V(\mathbf{y}^0) + c = \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}^0, \mathbf{x}) + V(\mathbf{x})) \, d\mu_0(\mathbf{x}).$$

It clearly follows that

$$\begin{aligned} V(\mathbf{y}^0) + c - \bar{V}(\mathbf{y}^0) - \bar{c} &\leq \int_{\Sigma_{\mathbf{M}}} [(A(\mathbf{y}^0, \mathbf{x}) + V(\mathbf{x})) - (A(\mathbf{y}^0, \mathbf{x}) + \bar{V}(\mathbf{x}))] \, d\mu_0(\mathbf{x}) \\ &= \int_{\Sigma_{\mathbf{M}}} (V(\mathbf{x}) - \bar{V}(\mathbf{x})) \, d\mu_0(\mathbf{x}) \leq V(\mathbf{y}^0) - \bar{V}(\mathbf{y}^0), \end{aligned}$$

which shows that $c \leq \bar{c}$. We can proceed in the same way changing in the reasoning V and \bar{V} . Therefore $c = \bar{c}$. \square

Theorem 6. *There exists the limit*

$$c_A := \lim_{\beta \rightarrow \infty} \frac{\lambda_\beta}{\beta}.$$

Proof. The previous propositions guarantee that $\{\lambda_\beta/\beta\}_{\beta>0}$ has a unique accumulation point as β goes to infinity. \square

In the next section, we explain how the real constant c_A is related with an ergodic Kantorovich transshipment problem.

3 Ergodic Transshipment

We remark that one may write, for all limit function $V \in \mathcal{H}$ and for any point \mathbf{y} ,

$$c_A = \max_{\mu \in \mathcal{M}_\sigma} \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y})) d\mu(\mathbf{x}). \quad (1)$$

Therefore, from ergodic optimization theory, one obtains that

$$c_A = \inf_{f \in C^0(\Sigma_{\mathbf{M}})} \sup_{\mathbf{x} \in \Sigma_{\mathbf{M}}} [A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y}) + f(\mathbf{x}) - f(\sigma(\mathbf{x}))].$$

Moreover, if we fixed a limit function $V \in \mathcal{H}$, for each point \mathbf{y} , there exists a function $U_{\mathbf{y}} \in \mathcal{H}$ (called a *sub-action with respect to $A(\mathbf{y}, \cdot) + V - V(\mathbf{y})$*) such that

$$A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y}) + U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x})) \leq c_A, \quad \forall \mathbf{x} \in \Sigma_{\mathbf{M}}, \quad (2)$$

and the equality holds on the support of the maximizing measure $\mu_{\mathbf{y}}^\infty$. We refer the reader to [5, 6, 9] for details on ergodic optimization theory.

Notice that equation (2) implies that

$$V(\mathbf{y}) + c_A \geq A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) + U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x})), \quad \forall (\mathbf{y}, \mathbf{x}) \in \hat{\Sigma}_{\mathbf{M}}.$$

Furthermore, since the equality holds at (\mathbf{y}, \mathbf{x}) whenever \mathbf{x} belongs to the support of $\mu_{\mathbf{y}}^\infty$, one has

$$V(\mathbf{y}) + c_A = \sup_{\mathbf{x}} [A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) + U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x}))], \quad \forall \mathbf{y} \in \Sigma_{\mathbf{M}}^*. \quad (3)$$

We get from the above equation (see, for instance, [2]) that V is an additive eigenfunction and c_A is an additive eigenvalue for

$$\mathcal{C}(\mathbf{y}, \mathbf{x}) := A(\mathbf{y}, \mathbf{x}) + U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x})), \quad \forall (\mathbf{y}, \mathbf{x}) \in \hat{\Sigma}_{\mathbf{M}}.$$

The question about uniqueness of the V which is solution of an additive problem is not so simple. For instance, it was considered in section 4 in [10], but it requires some stringent assumptions.

Notice now that, by its very construction, the map $(\mathbf{y}, \mathbf{x}) \mapsto U_{\mathbf{y}}(\mathbf{x})$ may depend on the fixed limit function V . Moreover, we only have information on its Lipschitz regularity on the \mathbf{x} variable. In particular, one cannot say *a priori* how the map $(\mathbf{y}, \mathbf{x}) \mapsto \mathcal{C}(\mathbf{y}, \mathbf{x})$ varies.

However, it is not difficult to provide examples of observables defining a continuous application \mathcal{C} as above. For instance, considering any $A_1, A_2 \in \mathcal{H}$, this is the case for the observable

$$A(\mathbf{y}, \mathbf{x}) = A_1(\mathbf{x}) + A_2(\mathbf{y}), \quad \forall (\mathbf{y}, \mathbf{x}) \in \hat{\Sigma}_{\mathbf{M}}.$$

Indeed, if $V \in \mathcal{H}$ is any possible limit function, let $U \in \mathcal{H}$ be a sub-action with respect to $A_1 + V$, that is:

$$A_1(\mathbf{x}) + V(\mathbf{x}) + U(\mathbf{x}) - U(\sigma(\mathbf{x})) \leq \max_{\mu \in \mathcal{M}_\sigma} \int (A_1 + V) d\mu, \quad \forall \mathbf{x} \in \Sigma_{\mathbf{M}}.$$

From (1), we then get

$$A(\mathbf{y}, \mathbf{x}) + V(\mathbf{x}) - V(\mathbf{y}) + U(\mathbf{x}) - U(\sigma(\mathbf{x})) \leq c_A$$

everywhere on $\hat{\Sigma}_{\mathbf{M}}$. In particular, we may choose $U_{\mathbf{y}} \equiv U$ for all \mathbf{y} in such a situation.

In general, by standard selection arguments (see section 2.1 in [12] and references therein), one may always assure the existence of a family of sub-actions $\{U_{\mathbf{y}}\}_{\mathbf{y}}$ for which the corresponding map $(\mathbf{y}, \mathbf{x}) \mapsto \mathcal{C}(\mathbf{y}, \mathbf{x})$ is Borel measurable. The main point is to consider just those sub-actions obtained as accumulation functions of eigenfunctions of Ruelle transfer operator when the temperature goes to zero through some fixed sequence (see proposition 29 in [5]). Note that these eigenfunctions are continuous on the observable. We leave the details to the reader. Finally, it is well known in ergodic optimization theory that these sub-actions have uniformly bounded oscillation. Hence, for each fixed limit function V , there exists a family $\{U_{\mathbf{y}}\}_{\mathbf{y}}$ of sub-actions with respect to $A(\mathbf{y}, \cdot) + V - V(\mathbf{y})$ such that the map

$$(\mathbf{y}, \mathbf{x}) \in \hat{\Sigma}_{\mathbf{M}} \mapsto \mathcal{C}(\mathbf{y}, \mathbf{x}) = A(\mathbf{y}, \mathbf{x}) + U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x}))$$

is Borel measurable and bounded³.

We consider from now on \mathcal{C} as a bounded measurable cost function in order to introduce a transshipment problem. Let then $\pi : \hat{\Sigma}_{\mathbf{M}} \rightarrow \Sigma_{\mathbf{M}}$ and $\pi^* : \hat{\Sigma}_{\mathbf{M}} \rightarrow \Sigma_{\mathbf{M}}^*$ be the canonical projections. We are specially interested in the set of Borel probabilities $\hat{\eta}(d\mathbf{y}, d\mathbf{x})$ on $\hat{\Sigma}_{\mathbf{M}}$ verifying $(\pi)_*(\hat{\eta}) = (\pi^*)_*(\hat{\eta})$.

Definition 4 (The Ergodic Kantorovich Transshipment Problem). *Given $A : \hat{\Sigma}_{\mathbf{M}} \rightarrow \mathbb{R}$ Lipschitz continuous, we are interested in the maximization problem*

$$\begin{aligned} \kappa_{\text{erg}} &:= \sup_{(\pi)_*(\hat{\eta})=(\pi^*)_*(\hat{\eta})} \iint_{\hat{\Sigma}_{\mathbf{M}}} \mathcal{C}(\mathbf{y}, \mathbf{x}) d\hat{\eta}(\mathbf{y}, \mathbf{x}) \\ &= \sup_{(\pi)_*(\hat{\eta})=(\pi^*)_*(\hat{\eta})} \iint_{\hat{\Sigma}_{\mathbf{M}}} [A(\mathbf{y}, \mathbf{x}) - U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x}))] d\hat{\eta}(\mathbf{y}, \mathbf{x}). \end{aligned}$$

An ergodic transshipment measure for A is a probability $\hat{\eta}$ on $\hat{\Sigma}_{\mathbf{M}}$, with $(\pi)_(\hat{\eta}) = (\pi^*)_*(\hat{\eta})$, that attains such a supremum.*

We point out that the classical transport or transshipment problems do not have an intrinsic ergodic nature. Note that \mathcal{C} has a dynamical character. We refer the reader to [14] for general results (not of ergodic nature) on transshipment. In [11], it is considered an ergodic transport problem.

We claim that $c_A = \kappa_{\text{erg}}$, or in a more self-contained statement:

Theorem 7. *For the Lipschitz observable βA , $\beta > 0$, consider its forward effective potential ϕ_{β}^+ (normalized by $\phi_{\beta}^+(\mathbf{y}^0) = 0$) and its effective constant λ_{β}^+ . Assume that in the uniform convergence $\phi_{\beta_n}^+/\beta_n \rightarrow V$, when $\beta_n \rightarrow \infty$. Then*

³Notice that it obviously follows from (2) that a such map \mathcal{C} is bounded from above.

there exists a family $\{U_{\mathbf{y}}\}_{\mathbf{y}}$ of sub-actions with respect to $A(\mathbf{y}, \cdot) + V - V(\mathbf{y})$ such that

$$\lim_{\beta \rightarrow \infty} \frac{\lambda_{\beta}^+}{\beta} = \sup_{(\pi)_*(\hat{\eta}) = (\pi^*)_*(\hat{\eta})} \iint_{\hat{\Sigma}_{\mathbf{M}}} [A(\mathbf{y}, \mathbf{x}) - U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x}))] d\hat{\eta}(\mathbf{y}, \mathbf{x}).$$

Proof. We remark that inequality (2) implies that $\kappa_{\text{erg}} \leq c_A$. Indeed, given any Borel probability $\hat{\eta}$ on $\hat{\Sigma}_{\mathbf{M}}$ such that $(\pi)_*(\hat{\eta}) = (\pi^*)_*(\hat{\eta})$, one clearly has

$$\begin{aligned} & \iint_{\hat{\Sigma}_{\mathbf{M}}} [A(\mathbf{y}, \mathbf{x}) - U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x}))] d\hat{\eta}(\mathbf{y}, \mathbf{x}) = \\ & = \iint_{\hat{\Sigma}_{\mathbf{M}}} [A(\mathbf{y}, \mathbf{x}) - U_{\mathbf{y}}(\mathbf{x}) - U_{\mathbf{y}}(\sigma(\mathbf{x})) + V(\mathbf{x}) - V(\mathbf{y})] d\hat{\eta}(\mathbf{y}, \mathbf{x}) \leq c_A. \end{aligned}$$

Recall that functional equation (3) shows the limit V is an additive eigenfunction and the constant c_A is an additive eigenvalue for \mathcal{C} . Actually, since \mathcal{C} is bounded, it is easy to obtain that c_A is uniquely determined by

$$c_A = \sup_{\{\mathbf{z}^k\}_{k \geq 1}} \limsup_{k \rightarrow \infty} \frac{\mathcal{C}(\mathbf{z}^1, \mathbf{z}^2) + \mathcal{C}(\mathbf{z}^2, \mathbf{z}^3) + \cdots + \mathcal{C}(\mathbf{z}^k, \mathbf{z}^1)}{k},$$

where the supremum is taken among sequences $\{\mathbf{z}^k\}$ of points of $\Sigma_{\mathbf{M}} \simeq \Sigma_{\mathbf{M}}^*$. See theorem 2.1 in [1] for a general result. Notice now that

$$\frac{\mathcal{C}(\mathbf{z}^1, \mathbf{z}^2) + \mathcal{C}(\mathbf{z}^2, \mathbf{z}^3) + \cdots + \mathcal{C}(\mathbf{z}^k, \mathbf{z}^1)}{k} = \iint_{\hat{\Sigma}_{\mathbf{M}}} \mathcal{C}(\mathbf{y}, \mathbf{x}) d\hat{\eta}_k(\mathbf{y}, \mathbf{x}),$$

where $\hat{\eta}_k$ is the Borel probability on $\hat{\Sigma}_{\mathbf{M}}$ defined by

$$\hat{\eta}_k := \frac{1}{k} \delta_{(\mathbf{z}^1, \mathbf{z}^2)} + \frac{1}{k} \delta_{(\mathbf{z}^2, \mathbf{z}^3)} + \cdots + \frac{1}{k} \delta_{(\mathbf{z}^k, \mathbf{z}^1)}.$$

Since $(\pi)_*(\hat{\eta}_k) = (\pi^*)_*(\hat{\eta}_k)$ for all $k \geq 1$, it obviously follows that $c_A \leq \kappa_{\text{erg}}$. \square

4 Contraction properties of \mathcal{G}^+

We would like to discuss now the proof of Theorem 1. We start pointing out an immediate contraction property of \mathcal{G}^+ .

Proposition 8. *For all $\phi, \psi \in \mathcal{H}$,*

$$\|\mathcal{G}^+(\phi) - \mathcal{G}^+(\psi)\|_0 \leq \|\phi - \psi\|_0.$$

Proof. Given $\mathbf{y} \in \Sigma_{\mathbf{M}}^*$, take $\mu_{\mathbf{y}} \in \mathcal{M}_{\sigma}$ satisfying

$$\mathcal{G}^+(\phi)(\mathbf{y}) = \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + \phi(\mathbf{x})) d\mu_{\mathbf{y}}(\mathbf{x}) + h_{\mu_{\mathbf{y}}}(\sigma).$$

Obviously $\mathcal{G}^+(\psi)(\mathbf{y}) \geq \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + \psi(\mathbf{x})) d\mu_{\mathbf{y}}(\mathbf{x}) + h_{\mu_{\mathbf{y}}}(\sigma)$. Therefore, we have

$$\mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y}) \leq \int_{\Sigma_{\mathbf{M}}} (\phi(\mathbf{x}) - \psi(\mathbf{x})) d\mu_{\mathbf{y}}(\mathbf{x}) \leq \|\phi - \psi\|_0.$$

Since ϕ and ψ play symmetrical roles, the proof is complete. \square

Notice that, for any real number γ , we have $\mathcal{G}^+(\phi + \gamma) = \mathcal{G}^+(\phi) + \gamma$. Let us now identify all functions belonging to \mathcal{H} which are equal up to an additive constant. So if we introduce the norm

$$\|\phi\|_c := \inf_{\gamma \in \mathbb{R}} \|\phi + \gamma\|_0$$

for each equivalence class $\phi \in \mathcal{H}/\text{constants}$, a fine contraction property can be verified.

Theorem 9. *Consider $\phi, \psi \in \mathcal{H}$ satisfying $\text{Lip}(\phi), \text{Lip}(\psi) \leq K$ for some fixed constant $K > 0$. Then, there exist constants $C = C(K) > 0$ and $\alpha = \alpha(K) > 0$ such that*

$$\|\mathcal{G}^+(\phi) - \mathcal{G}^+(\psi)\|_c \leq (1 - C\|\phi - \psi\|_c^\alpha) \|\phi - \psi\|_c$$

We will need the following lemma.

Lemma 10. *Let $A : \hat{\Sigma}_{\mathbf{M}} \rightarrow \mathbb{R}$ be Lipschitz continuous observable. Suppose $\phi \in \mathcal{H}$ satisfies $\text{Lip}(\phi) \leq K$ for a constant $K > 0$. Given a point $\mathbf{y} \in \Sigma_{\mathbf{M}}^*$, let $\mu_{\mathbf{y}} \in \mathcal{M}_\sigma$ be the equilibrium state associated to $A(\mathbf{y}, \cdot) + \phi \in \mathcal{H}$. Then there exist constants $\Gamma = \Gamma(K) > 0$ and $\alpha = \alpha(K) > 0$ such that, if $B_\rho \subset \Sigma_{\mathbf{M}}$ denotes an arbitrary ball of radius $\rho > 0$,*

$$\mu_{\mathbf{y}}(B_\rho) \geq \Gamma \rho^\alpha.$$

Proof. Let $\mu_\Psi \in \mathcal{M}_\sigma$ be the equilibrium measure associated to $\Psi \in \mathcal{H}$. It is well known that μ_Ψ is a Gibbs state. As a matter of fact, if \mathbf{x} is a point belonging to a ball B_{Λ^n} of radius Λ^n , from the very proof of the Gibbs property one can obtain

$$\begin{aligned} \exp[-\text{Lip}(\Psi)R(\Lambda) - I_{\mathbf{M}}(\text{Lip}(\Psi) + h_{TOP}(\sigma))S(\Lambda)] &\leq \\ &\leq \frac{\mu_\Psi(B_{\Lambda^n})}{\exp\left[\sum_{j=0}^{n-1} (\Psi - P_{TOP}(\Psi)) \circ \sigma^j(\mathbf{x})\right]}, \end{aligned}$$

where R and S are rational functions with $R(0, 1), S(0, 1) \subset (0, +\infty)$, $I_{\mathbf{M}}$ is a positive integer depending only on the irreducible transition matrix \mathbf{M} and $h_{TOP}(\sigma)$ denotes the topological entropy. For details we refer the reader to [13].

From the variational principle, one has $\Psi - P_{TOP}(\Psi) \geq -\text{Lip}\Psi - h_{TOP}(\sigma)$. Therefore, we immediately get

$$\exp[-\text{Lip}(\Psi)R(\Lambda) - (\text{Lip}(\Psi) + h_{TOP}(\sigma))(I_{\mathbf{M}}S(\Lambda) + n)] \leq \mu_\Psi(B_{\Lambda^n}).$$

Thus, applying this inequality to $\Psi = A(\mathbf{y}, \cdot) + \phi$, it is straightforward to verify that

$$\Gamma(K)\Lambda^{n\alpha(K)} \leq \mu_{\mathbf{y}}(B_{\Lambda^n}),$$

where

$$\alpha(K) := \frac{\text{Lip}(A) + K + h_{TOP}(\sigma)}{\log \Lambda^{-1}} \quad \text{and}$$

$$\Gamma(K) := \exp[-(\text{Lip}(A) + K)R(\Lambda) - I_{\mathbf{M}}(\text{Lip}(A) + K + h_{TOP}(\sigma))S(\Lambda)].$$

□

Proof of Theorem 9. Obviously, for $\phi \in \mathcal{H}$ and $\gamma \in \mathbb{R}$, we have $\|\phi + \gamma\|_c = \|\phi\|_c$. Moreover, given $\phi, \psi \in \mathcal{H}$, there exists $\bar{\gamma} \in \mathbb{R}$ such that $\|\phi - \psi\|_c = \|\phi - \psi + \bar{\gamma}\|_0$.

As \mathcal{G} commutes with constants, replacing ψ by $\psi - \min \psi$ and ϕ by $\phi + \bar{\gamma} - \min \psi$, without loss of generality, we may assume

$$\min \psi = 0 \quad \text{and} \quad \|\phi - \psi\|_c = \|\phi - \psi\|_0.$$

We suppose yet $\phi \neq \psi$, since otherwise there is nothing to argue.

Take then $\mathbf{y} \in \Sigma_{\mathbf{M}}^*$ satisfying

$$\|\mathcal{G}^+(\phi) - \mathcal{G}^+(\psi)\|_0 = |\mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y})|.$$

By interchanging the roles of ϕ and ψ if necessary, we suppose that

$$|\mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y})| = \mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y}).$$

Since $\min \psi = 0$, taking any point $\mathbf{x} \in \Sigma_{\mathbf{M}}$, we get

$$\|\phi - \psi\|_c \leq \|\phi - \phi(\mathbf{x}) - \psi\|_0 \leq \|\phi - \phi(\mathbf{x})\|_0 + \|\psi\|_0 \leq \text{Lip}(\phi) + \text{Lip}(\psi) \leq 2K.$$

Note that $\|\phi - \psi\|_c = \|\phi - \psi\|_0$ implies $\min(\phi - \psi) = -\max(\phi - \psi)$. In particular, $\min(\phi - \psi) = -\|\phi - \psi\|_c$. So there exists a point $\bar{\mathbf{x}} \in \Sigma_{\mathbf{M}}$ such that

$$(\phi - \psi)(\bar{\mathbf{x}}) = -\|\phi - \psi\|_c < 0.$$

Hence, when $\mathbf{x} \in \Sigma_{\mathbf{M}}$ verifies $d(\mathbf{x}, \bar{\mathbf{x}}) \leq \frac{\|\phi - \psi\|_c}{4K}$, we obtain

$$\begin{aligned} \phi(\mathbf{x}) - \psi(\mathbf{x}) &\leq |\phi(\mathbf{x}) - \phi(\bar{\mathbf{x}})| + |\psi(\bar{\mathbf{x}}) - \psi(\mathbf{x})| + (\phi - \psi)(\bar{\mathbf{x}}) \\ &\leq 2K \frac{\|\phi - \psi\|_c}{4K} - \|\phi - \psi\|_c \\ &= -\frac{\|\phi - \psi\|_c}{2} < 0. \end{aligned} \tag{4}$$

Let then $\mu_{\mathbf{y}} \in \mathcal{M}_{\sigma}$ be such that

$$\mathcal{G}^+(\phi)(\mathbf{y}) = \int_{\Sigma_{\mathbf{M}}} (A(\mathbf{y}, \mathbf{x}) + \phi(\mathbf{x})) \, d\mu_{\mathbf{y}}(\mathbf{x}) + h_{\mu_{\mathbf{y}}}(\sigma).$$

As in the previous proposition, it follows

$$\mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y}) \leq \int_{\Sigma_{\mathbf{M}}} (\phi(\mathbf{x}) - \psi(\mathbf{x})) \, d\mu_{\mathbf{y}}(\mathbf{x}).$$

So if $B_{\frac{\|\phi - \psi\|_c}{4K}}(\bar{\mathbf{x}})$ denotes the closed ball of radius $\frac{\|\phi - \psi\|_c}{4K} \in (0, 1)$ and center $\bar{\mathbf{x}} \in \Sigma_{\mathbf{M}}$, from (4) and lemma 10, we verify

$$\begin{aligned} \mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y}) &\leq \int_{\Sigma_{\mathbf{M}} - B_{\frac{\|\phi - \psi\|_c}{4K}}(\bar{\mathbf{x}})} (\phi(\mathbf{x}) - \psi(\mathbf{x})) \, d\mu_{\mathbf{y}}(\mathbf{x}) \\ &\leq \|\phi - \psi\|_0 \left(1 - \mu_{\mathbf{y}}\left(B_{\frac{\|\phi - \psi\|_c}{4K}}(\bar{\mathbf{x}})\right)\right) \\ &\leq \|\phi - \psi\|_0 (1 - C\|\phi - \psi\|_c^{\alpha}), \end{aligned}$$

where $C := \Gamma/(4K)^{\alpha} > 0$.

As $\|\mathcal{G}^+(\phi) - \mathcal{G}^+(\psi)\|_c \leq \|\mathcal{G}^+(\phi) - \mathcal{G}^+(\psi)\|_0 = \mathcal{G}^+(\phi)(\mathbf{y}) - \mathcal{G}^+(\psi)(\mathbf{y})$, the proof is complete. \square

Theorem 1 is then a direct consequence of Theorem 9, the fact that

$$\text{Lip}(\mathcal{G}^+(\phi)) \leq \|A\|_0 + \text{Lip}(A) \quad \forall \phi \in \mathcal{H},$$

and the following fixed point theorem due to D. A. Gomes and E. Valdinoci (for a proof, see Appendix A of [8]).

A Banach–Caccioppoli-type Theorem. *Let \mathbf{F} be a closed subset of a Banach space, endowed with a norm $\|\cdot\|$. Suppose that $G : \mathbf{F} \rightarrow \mathbf{F}$ is so that*

$$\|G(\phi) - G(\psi)\| \leq (1 - C\|\phi - \psi\|^\alpha) \|\phi - \psi\|,$$

for all $\phi, \psi \in \mathbf{F}$ and some given constants $C, \alpha > 0$. Then there exists a unique $\phi^+ \in \mathbf{F}$ such that $G(\phi^+) = \phi^+$. Moreover, given any $\phi_0 \in \mathbf{F}$, we have

$$\phi^+ = \lim_{n \rightarrow +\infty} G^n(\phi_0).$$

References

- [1] N. Bacaer, Convergence of numerical methods and parameter dependence of min-plus eigenvalue problems, Frenkel-Kontorova models and homogenization of Hamilton-Jacobi equations, *ESAIM: Mathematical Modelling and Numerical Analysis* **35** (2001), 1185-1195.
- [2] W. Chou and R. J. Duffin, An additive eigenvalue problem of physics related to linear programming, *Advances in Applied Mathematics* **8** (1987), 486-498.
- [3] W. Chou and R. Griffiths, Ground states of one-dimensional systems using effective potentials, *Physical Review B* **34** (1986), 6219-6234.
- [4] M. C. Concordel, Periodic homogenization of Hamilton-Jacobi equations: additive eigenvalues and variational formula, *Indiana University Mathematical Journal* **45** (1996), 1095-1118.
- [5] G. Contreras, A. O. Lopes and Ph. Thieullen, Lyapunov minimizing measures for expanding maps of the circle, *Ergodic Theory and Dynamical Systems* **21** (2001), 1379-1409.
- [6] J. P. Conze and Y. Guivarc'h, *Croissance des sommes ergodiques et principe variationnel*, manuscript, circa 1993.
- [7] D. A. Gomes, A. O. Lopes and J. Mohr, The Mather measure and a large deviation principle for the entropy penalized method, to appear in *Communications in Contemporary Mathematics*.
- [8] D. A. Gomes and E. Valdinoci, Entropy penalization methods for Hamilton-Jacobi equations, *Advances in Mathematics* **215** (2007), 94-152.
- [9] O. Jenkinson, Ergodic optimization, *Discrete and Continuous Dynamical Systems, Series A* **15** (2006), 197-224.
- [10] A. O. Lopes, J. Mohr, R. Souza and Ph. Thieullen, Negative entropy, zero temperature and stationary Markov chains on the interval, *Bulletin of the Brazilian Mathematical Society* **40** (2009), 1-52.
- [11] A. O. Lopes, E. R. Oliveira and Ph. Thieullen, The dual potential, the involution kernel and transport in ergodic optimization, *preprint*, 2008.
- [12] I. Molchanov, *Theory of random sets*, Springer-Verlag, London, 2005.
- [13] W. Parry and M. Pollicott, Zeta functions and the periodic orbit structure of hyperbolic dynamics, *Astérisque* **187-188** (1990).
- [14] S. T. Rachev and L. Rüschendorf, *Mass transportation problems – Volume I: Theory, Volume II: Applications*, Springer-Verlag, New York, 1998.
- [15] M. Salmhofer, *Renormalization: an introduction*, Springer-Verlag, Berlin, 1999.
- [16] C. Villani, *Topics in optimal transportation*, AMS, Providence, 2003.
- [17] C. Villani, *Optimal transport: old and new*, Springer-Verlag, Berlin, 2009.