

Bayesian estimation of a skew-t stochastic volatility model

C. A. Abanto-Valle^{†1}, V. H. Lachos[‡] and Dipak K. Dey[§]

[†]Department of Statistics, Federal University of Rio de Janeiro, Brazil

[‡]Department of Statistics, Campinas State University, Brazil

[§]Department of Statistics, University of Connecticut, USA

Abstract

In this paper we present a stochastic volatility (SV) model assuming that the return shock has a skew-Student-t distribution. This allows a parsimonious, flexible treatment of asymmetry and heavy tails in the conditional distribution of returns. An efficient Markov chain Monte Carlo estimation method is described that exploits a skew-normal mixture representation of the error distribution with a gamma distribution as the mixing distribution. We apply the methodology to the NASDAQ daily index returns.

keywords: Markov chain Monte Carlo, non-Gaussian and nonlinear state space models, skew-Student-t, stochastic volatility.

1 Introduction

A large literature in financial econometrics has documented stylized facts which are frequently found in stock and foreign exchange returns: skewness, heavy-tailedness and volatility clustering. These properties are crucial not only for describing the return distributions but also for asset allocation, option pricing, forecasting and risk management.

¹Correspondence to: Carlos A. Abanto-Valle, Department of Statistics, Federal University of Rio de Janeiro, Caixa Postal 68530, CEP: 21945-970, RJ, Brazil. E-mail: cabantovalle@im.ufrj.br

Stochastic volatility (SV) models were introduced in the financial literature for describing time-varying volatilities (Taylor, 1982; 1986). Various extensions of the simple SV model with a normal errors have been discussed in the literature. For instance, many empirical studies have shown strong evidence of heavy-tailed conditional mean errors in financial time series (see for example Mandelbrot, 1963; Fama, 1965; Chib *et al.*, 2002; Jacquier *et al.*, 2004). In this context, recently Abanto-Valle *et al.* (2010) extended the basic SV model by assuming the flexible class of scale mixtures of normal distributions (Lange and Sinsheimer, 1993). The empirical evidence on the presence of asymmetry in the distribution of financial returns is not as clear-cut even though asymmetry plays a non-trivial role in shaping economic decisions. Corrado and Su (1997) suggests that fat tails and asymmetry jointly determine the so-called “volatility smile” in option pricing using the Black-Scholes approach and that explicit account of them improve accuracy in option pricing. Peiro (1999) provides further evidence of asymmetry in returns, both from stock market indices and from individual assets. Further, Mittnik and Paoletta (2000) argue that skewness and heavy tails should be taken into account explicitly in Value-at-Risk forecasts. Cappuccio *et al.* (2006) found empirical evidence on asymmetry in financial returns using a simple stochastic volatility modeling both skewness and heavy tails assuming that the conditional distribution of returns is a skew-generalized error distribution.

In this paper, in order to model simultaneously skewness and heavy-tailedness, we extend the SV model by assuming skew-Student-t (ST) introduced by Azzalini and Capitanio (2003) and hence the SV-ST is defined. Inference in the SV-ST model is performed under a Bayesian paradigm via MCMC methods, which permits to obtain the posterior distribution of parameters by simulation starting from reasonable prior assumptions on the parameters.

The remainder of this paper is organized as follows. Section 2 shows a brief review about skew-normal (Azzalini, 1986) and skew-t distributions and their properties. Section 3 describes the SV-ST model through Bayesian estimation procedure using MCMC methods. Section 4 is devoted to application

and model comparison among the SV-ST model against the SV-N, SV-T and SV-SN models using the NASDAQ data set. Finally, some concluding remarks as well as future developments are deferred to Section 5.

2 The univariate skew-normal and skew-t distributions

We start by giving an important notation that will be used throughout the paper and present a review of the univariate skew normal (SN) and skew-t (ST) distributions and a study of some related properties of those distributions.

A univariate random variable X is said to follow a skew-normal distribution, $X \sim \mathcal{SN}(\zeta, \omega^2, \lambda)$, with location, scale and asymmetry parameters given by ζ , ω^2 and λ , respectively, if the density of this distribution has the form

$$p(x | \zeta, \omega^2, \lambda) = \frac{2}{\omega} \phi\left(\frac{x - \zeta}{\omega}\right) \Phi\left(\frac{\lambda}{\omega}(x - \zeta)\right), \quad (1)$$

where $\phi(\cdot)$ and $\Phi(\cdot)$ are, respectively, the probability density function (pdf) and the cumulative distribution function (cdf) of the standard normal distribution. When $\lambda = 0$, the density in equation (1) becomes $\mathcal{N}(\zeta, \sigma^2)$ (see [Azzalini, 2005](#), for a comprehensive review). In the next sections, we use the following stochastic representation of the SN distribution ([Azzalini, 1986](#); [Henze, 1986](#)). Let $W \sim \mathcal{N}_{[0, \infty)}(0, 1)$ and $\varepsilon \sim \mathcal{N}(0, 1)$, independently, and let $\delta \in (-1, 1)$, where $\mathcal{N}_{[0, \infty)}(\cdot, \cdot)$ and $\mathcal{N}(\cdot, \cdot)$ indicate the truncated normal and normal distribution, respectively. The random variable X , defined by

$$X = \zeta + \omega\delta W + \omega\sqrt{1 - \delta^2}\varepsilon, \quad (2)$$

follows a univariate skew-normal distributions, that is, $X \sim \mathcal{SN}(\zeta, \omega^2, \lambda)$, where $\lambda = \delta/\sqrt{1 - \delta^2}$.

The kurtosis coefficient of a skew-normal distribution is restricted to the interval $[3, 3.8692]$. To achieve a higher degree of excess kurtosis, the skew-t distribution has been introduced by [Branco and](#)

Dey (2001) and Azzalini and Capitanio (2003). A univariate random variable X follows the scalar skew-t distribution, $X \sim \mathcal{ST}(\zeta, \omega^2, \lambda, \nu)$, if it has the following stochastic representation

$$X = \zeta + U^{-1/2}\omega\delta W + U^{-\frac{1}{2}}\omega(1 - \delta^2)^{\frac{1}{2}}\varepsilon, \quad (3)$$

where $W \sim \mathcal{N}_{[0, \infty)}(0, 1)$, $\varepsilon \sim \mathcal{N}(0, 1)$ and $U \sim \mathcal{G}(\frac{\nu}{2}, \frac{\nu}{2})$ are independently distributed. The Gamma distribution $\mathcal{G}(a, b)$ is defined with density $p(u | a, b) = b^a u^{a-1} e^{-bu} / \Gamma(a)$. The pdf of X is then given by

$$f(X | \zeta, \omega^2, \lambda, \nu) = \frac{2}{\omega} t_\nu \left(\frac{x - \zeta}{\omega} \right) T_{\nu+1} \left(\lambda \omega^{-1} (x - \zeta) \sqrt{\frac{\nu + 1}{\nu + \omega^{-2} (x - \zeta)^2}} \right), \quad (4)$$

where $t_\nu(\cdot)$ and $T_\nu(\cdot)$ denote the pdf and cdf of a standard Student-t distribution with ν degrees of freedom. From (3), we have that

$$E(X) = \zeta + \sqrt{\frac{2}{\pi}} k_1 \omega \delta, \quad (5)$$

$$V(X) = \omega^2 k_2 - \frac{2}{\pi} k_1^2 \omega^2 \delta^2, \quad (6)$$

where $\delta = \lambda / \sqrt{1 + \lambda^2}$ and $k_m = E(U^{-m/2})$. $E(\cdot)$ and $V(\cdot)$ denote the expected value and variance, respectively. The skew-t nests the traditional symmetric Student's t distribution as a special case when $\lambda = 0$, and the conditional normal distribution as $\nu \rightarrow \infty$, and can capture left-tailed or negative skewness when $\lambda < 0$, and positive skewness when $\lambda > 0$.

To interpret the parameters (λ, ν) in relation to the skewness and heavy-tailedness, skew-t densities are plotted using several combinations of the parameter values in Figure 1 with ζ and ω fixed at 0 and 2, respectively. In Figure 1, left, the densities are drawn using $\lambda = 0, -2, -4, -8$ with ν fixed at 5. As mentioned, $\lambda = 0$ corresponds to a symmetric Student's t-density. A lower value of λ implies a more negative skewness or left-skewness as well as heavier tails. Figure 1, right, shows the densities for ν at 2, 4, 10 and 15 with λ fixed equal to -2. As ν becomes larger, the density becomes less skewed and has lighter tails. Hence the skewness and heavy-tailedness are determined jointly by the combination of the parameter values of λ and ν .




Figure 1: The skew-t distribution. Top: $\zeta = 0, \omega = 2, \nu = 5$ (fixed), $\lambda = 0, -2, -4, -8$. Bottom: $\zeta = 0, \omega = 2, \lambda = -2$ (fixed), $\nu = 2, 4, 10$ and 15 .

3 The skew-t stochastic volatility model

3.1 The model

In order to account for both the excess kurtosis and skewness in stock returns, we introduce the stochastic volatility model with skew-t errors (SV-ST), which is defined as

$$y_t = e^{\frac{h_t}{2}} \epsilon_t, \quad (7a)$$

$$h_{t+1} = \mu + \varphi(h_t - \mu) + \sigma_\eta \eta_t, \quad (7b)$$

where y_t and h_t are, respectively, the compounded return and the log-volatility at time t . We assume that $|\varphi| < 1$, i.e., the log-volatility process is stationary and that the initial value $h_1 \sim \mathcal{N}(\mu, \frac{\sigma_\eta^2}{1-\varphi^2})$, $\epsilon_t \sim \mathcal{ST}(\zeta, \omega^2, \lambda, \nu)$ and $\eta_t \sim \mathcal{N}(0, 1)$ are uncorrelated. The SV-ST defined by equations (7a) and (7b) can be written hierarchically using the stochastic representation of the skew-t distribution in (3), as

$$y_t = (\zeta + \omega\delta W_t U_t^{-\frac{1}{2}})e^{\frac{h_t}{2}} + e^{\frac{h_t}{2}} U_t^{-\frac{1}{2}} \omega(1 - \delta^2)^{\frac{1}{2}} \epsilon_t, \quad (8a)$$

$$h_{t+1} = \mu + \varphi(h_t - \mu) + \sigma_\eta \eta_t, \quad (8b)$$

$$W_t \sim \mathcal{N}_{[0, \infty)}(0, 1), \quad (8c)$$

$$U_t | \nu \sim \mathcal{G}(\frac{\nu}{2}, \frac{\nu}{2}), \quad (8d)$$

where ϵ_t and η_t are mutually independent and normally distributed with zero mean and unit variance, $\delta = \frac{\lambda}{\sqrt{1+\lambda^2}}$. We set ζ and ω in such a way that $E(y_t | h_t) = 0$ and $V(y_t | h_t) = e^{h_t}$. In this setup, equations (8a) and (8b), with $\lambda = 0$ (equivalently $\delta = 0$) and $U_t = 1 \forall t$ define the SV model with normal distribution (SV-N). Equations (8a),(8b) and (8d) with $\lambda = 0$ define the the SV model with Student-t distribution (SV-T) defined by [Abanto-Valle *et al.* \(2010\)](#). Finally, equations (8a),(8b) and (8c) with $U_t = 1, \forall t$ results the SV model with skew normal distribution (SV-SN).

3.2 Parameter estimation via MCMC

Let $\boldsymbol{\theta} = (\varphi, \sigma_\eta^2, \nu, \lambda)'$ be the full parameter vector of the entire class of SV-ST model, $\mathbf{h}_{1:T} = (h_1, \dots, h_T)'$ be the vector of the log volatilities, $\mathbf{U}_{1:T} = (U_1, \dots, U_T)'$ be the mixing variables, $\mathbf{W}_{1:T} = (W_1, \dots, W_T)'$ and $\mathbf{y}_{1:T} = (y_1, \dots, y_T)'$ be the information available up to time T , while ν is the degrees of freedom parameter vector associated with the mixture distribution and λ the skewness parameter. The Bayesian approach to estimate the parameters in the SV-ST model uses the data augmentation principle, which considers $\mathbf{h}_{1:T}$, $\mathbf{W}_{1:T}$ and $\mathbf{U}_{1:T}$ as latent variables. The joint posterior density of parameters and latent

unobservable variables can be written as

$$\begin{aligned}
p(\boldsymbol{\theta}, \mathbf{W}_{1:T}, \mathbf{U}_{1:T}, \mathbf{h}_{0:T} \mid \mathbf{y}_{1:T}) &\propto p(\mathbf{y}_{1:T} \mid \boldsymbol{\theta}, \mathbf{W}_{1:T}, \mathbf{U}_{1:T}, \mathbf{h}_{1:T}) \\
&\times p(\mathbf{h}_{0:T} \mid \boldsymbol{\theta})p(\mathbf{W}_{1:T})p(\mathbf{U}_{1:T} \mid \boldsymbol{\theta})p(\boldsymbol{\theta}),
\end{aligned} \tag{9}$$

where $p(\boldsymbol{\theta})$ is the prior distribution. Since the posterior density $p(\mathbf{h}_{1:T}, \mathbf{W}_{1:T}, \mathbf{U}_{1:T}, \boldsymbol{\theta} \mid \mathbf{y}_{1:T})$ does not have closed form, we first sample the parameters $\boldsymbol{\theta}$, followed by the latent variables $\mathbf{W}_{1:T}$, $\mathbf{U}_{1:T}$ and $\mathbf{h}_{1:T}$ using Gibbs sampling. The sampling scheme is described by Algorithm 1. Sampling the log-volatilities $\mathbf{h}_{1:T}$ in step 5 of Algorithm 1 is the most difficult task due to the nonlinear setup in the observational equation (8a). In order to avoid the higher correlations due to the Markovian structure of the h_t 's, in the next subsection we develop a multi-move block sampler to sample $\mathbf{h}_{0:T}$ by blocks (Shephard and Pitt 1997; Watanabe and Omori 2004; Abanto-Valle *et al.* 2010). Details on the full conditionals of $\boldsymbol{\theta}$ and the latent variables $\mathbf{U}_{1:T}$ and $\mathbf{W}_{1:T}$ are given in Appendix.

Algorithm 1

1. Set $i = 0$ and get starting values for the parameters $\boldsymbol{\theta}^{(i)}$ and the latent quantities $\mathbf{W}_{1:T}^{(i)}$, $\mathbf{U}_{1:T}^{(i)}$ and $\mathbf{h}_{1:T}^{(i)}$.
2. Generate $\boldsymbol{\theta}^{(i+1)}$ in turn from its full conditional distribution, given $\mathbf{y}_{1:T}$, $\mathbf{h}_{1:T}^{(i)}$, $\mathbf{W}_{1:T}^{(i)}$ and $\mathbf{U}_{1:T}^{(i)}$.
3. Draw $\mathbf{W}_{1:T}^{(i+1)} \sim p(\mathbf{W}_{1:T} \mid \boldsymbol{\theta}^{(i)}, \mathbf{U}_{1:T}^{(i)}, \mathbf{h}_{1:T}^{(i)}, \mathbf{y}_{1:T})$.
4. Draw $\mathbf{U}_{1:T}^{(i+1)} \sim p(\mathbf{U}_{1:T} \mid \boldsymbol{\theta}^{(i+1)}, \mathbf{W}_{1:T}^{(i+1)}, \mathbf{h}_{1:T}^{(i)}, \mathbf{y}_{1:T})$.
5. Generate $\mathbf{h}_{1:T}$ by blocks as:
 - i) For $l = 1, \dots, K$, the knot positions are generated as k_l , the floor of $[T \times \{(l + u_l)/(K + 2)\}]$, where the u_l 's are independent realizations of the uniform random variable on the interval (0,1).

- ii) For $l = 1, \dots, K$, generate $h_{k_{l-1}+1:k_l-1}$ jointly conditional on $\mathbf{y}_{k_{l-1}:k_l-1}$, $\boldsymbol{\theta}^{(i+1)}$, $\mathbf{W}_{k_{l-1}+1:k_l-1}^{(i+1)}$, $\mathbf{U}_{k_{l-1}+1:k_l-1}^{(i+1)}$, $h_{k_{l-1}}^{(i)}$ and $h_{k_l}^{(i)}$.
- iii) For $l = 1, \dots, K$, draw $h_{k_l}^{(i+1)}$ conditional on $\mathbf{y}_{1:T}$, $\boldsymbol{\theta}^{(i)}$, $W_{k_l}^{(i+1)}$, $U_{k_l}^{(i+1)}$, $h_{k_{l-1}}^{(i+1)}$ and $h_{k_{l+1}}^{(i+1)}$.

6. Set $i = i + 1$ and return to 2 until convergence is achieved.

In the SV-ST model considered so far, an important modelling assumption is the regularization penalty $p(\nu)$ on the tail thickness. A default Jeffreys' prior was developed by [Fonseca et al. \(2008\)](#), with a number of desirable properties particularly when learning a fat-tail from a finite dataset. The default Jeffreys's prior for ν takes the form

$$p(\nu) \propto \left(\frac{\nu}{\nu + 3} \right)^{\frac{1}{2}} \left\{ \psi' \left(\frac{\nu}{2} \right) - \psi' \left(\frac{\nu + 1}{2} \right) - \frac{2(\nu + 3)}{\nu(\nu + 1)^2} \right\}^{\frac{1}{2}}, \quad (10)$$

where $\psi'(a) = \frac{d\{\psi(a)\}}{da}$ and $\psi(a) = \frac{d\{\log \Gamma(a)\}}{da}$ are the trigamma and digamma functions, respectively. The interesting feature of this prior is its behavior as ν goes to infinity and it has polynomial tails of the form $p(\nu) \propto \nu^{-4}$. In this case, the tail of the prior decays rather fast for large values of ν and assessing the degree of tail thickness can require prohibitively large samples. To the skewness parameter, we assume that $\lambda \sim t_{0.5}(0.0, \frac{\pi^2}{4})$, a Jeffreys' prior suggested by [Bayes and Branco \(2007\)](#).

3.2.1 Block sampler

In order to simulate $\mathbf{h}_{1:T} = (h_1, \dots, h_T)'$ in the SV-ST model, we consider a two-step process: first, we simulate h_1 conditional on $\mathbf{h}_{2:T}$, next $\mathbf{h}_{2:T}$ conditional on h_1 . To sample the vector $\mathbf{h}_{2:T}$, we develop a multi-move block algorithm. In our block sampler, we divide it into $K + 1$ blocks, $\mathbf{h}_{k_{l-1}+1:k_l-1} = (h_{k_{l-1}+1}, \dots, h_{k_l-1})'$ for $l = 1, \dots, K + 1$, with $k_0 = 1$ and $k_{K+1} = T$, where $k_l - 1 - k_{l-1} \geq 2$ is the size of the l -th block. We sample the block of disturbances $\boldsymbol{\eta}_{k_{l-1}:k_l-2} = (\eta_{k_{l-1}}, \dots, \eta_{k_l-2})'$ given the end conditions $h_{k_{l-1}}$ and h_{k_l} instead of $\mathbf{h}_{k_{l-1}+1:k_l-1}$. In order to facilitate the exposition, we omit the dependence on $\boldsymbol{\theta}$, $\mathbf{W}_{t+1:t+k}$ and $\mathbf{U}_{t+1:t+k}$, and suppose that $k_{l-1} = t$ and $k_l = t + k + 1$ for the l -th block,

such that $t + k < T$. Then $\boldsymbol{\eta}_{t:t+k-1} = (\eta_t, \dots, \eta_{t+k-1})'$ are sampled at once from their full conditional distribution $f(\boldsymbol{\eta}_{t:t+k-1} | h_t, h_{t+k+1}, \mathbf{y}_{t:t+k})$, which without the constant terms is expressed in log scale as

$$\begin{aligned} \log f(\boldsymbol{\eta}_{t:t+k-1} | h_t, h_{t+k+1}) &= \text{const} - \frac{1}{2} \sum_{r=t}^{t+k-1} \eta_r^2 + \sum_{r=t+1}^{t+k} l(h_r) \\ &\quad - \frac{1}{2\sigma_\eta^2} [h_{t+k+1} - \mu - \varphi(h_{t+k} - \mu)]^2 \mathbb{I}(t+k < T), \end{aligned}$$

where $\mathbb{I}(\cdot)$ is an indicator function. We denote the first and second derivatives of $l(h_r)$ with respect to h_r by l' and l'' , where $l(h_r) = \log p(y_r | \nu, \lambda, W_r, U_r, h_r)$ is obtained from equation (8a). As (11) does not have closed form, we use the Metropolis-Hastings acceptance-rejection algorithm (Tierney, 1994; Chib and Greenberg, 1995) to sample from. We propose to use the following artificial Gaussian state space model as a proposed density to simulate the block $\boldsymbol{\eta}_{t+1:t+k}$

$$\hat{y}_r = h_r + \xi_r, \quad \xi_r \sim \mathcal{N}(0, d_r), \quad r = t+1, \dots, t+k, \quad (11)$$

$$h_{r+1} = \mu + \varphi(h_r - \mu) + \sigma_\eta \eta_r, \quad \eta_r \sim \mathcal{N}(0, 1), \quad r = t, t+1, \dots, t+k-1, \quad (12)$$

where the auxiliary variables d_r and \hat{y}_r for $r = t+1, \dots, t+k-1$ and $t+k = T$ are defined as follows:

$$\begin{aligned} d_r &= -\frac{1}{l''_F(\hat{h}_r)}, \\ \hat{y}_r &= \hat{h}_r + d_r l'(\hat{h}_r). \end{aligned} \quad (13)$$

For $r = t+k < T$, it follows that

$$\begin{aligned} d_r &= \frac{\sigma_\eta^2}{\varphi^2 - \sigma_\eta^2 l''_F(\hat{h}_{t+k})}, \\ \hat{y}_r &= d_r \left[l'(\hat{h}_r) - l''_F(\hat{h}_r) \hat{h}_r + \frac{\varphi}{\sigma_\eta^2} [h_{r+1} - \mu(1 - \varphi)] \right]. \end{aligned} \quad (14)$$

We obtain the measurement equation (11) by a second-order expansion of l_r around some preliminary estimate of η_r , denoted by $\hat{\eta}_r$, where \hat{h}_r is the estimate of h_r equivalent to $\hat{\eta}_r$, and

$$l''_F(h_r) = E[l''(h_r)] = -\frac{1}{2} - \frac{(\zeta + \omega \delta W_t U_t^{-\frac{1}{2}})^2}{4\omega^2(1 - \delta^2)} U_r, \quad (15)$$

which is everywhere strictly negative. The expectation in (15) is taken with respect to y_r conditional on $h_r, W_r, U_r, \boldsymbol{\theta}$. Since (11)-(12) define a Gaussian state space model, we can apply de Jong and Shephard's simulation smoother (de Jong and Shephard, 1995) to perform the sampling. We denote this density by g . Since f is not bounded by g , we use the Metropolis-Hastings acceptance-rejection algorithm to sample from f , as recommended by Chib and Greenberg (1995). In the SV-SN case, we use the same procedure with $U_t = 1$ for $t = 1, \dots, T$.

The procedure to select the expansion block $\hat{\mathbf{h}}_{t+1:t+k}$ is described in the Algorithm 2.

Algorithm 2

1. Initialize $\hat{\mathbf{h}}_{t+1:t+k}$.
2. Evaluate recursively $l'(\hat{h}_r)$ and $l''_F(\hat{h}_r)$ for $r = t + 1, \dots, t + k$.
3. Conditional on the current values of the vector of parameters $\boldsymbol{\theta}, \mathbf{U}_{t+1:t+k}, \mathbf{W}_{t+1:t+k}, h_t$ and h_{t+k+1} , define the auxiliary variables \hat{y}_r and d_r using equations (13) or (14) for $r = t + 1, \dots, t + k$.
4. Consider the linear Gaussian state-space model in (11) and (12). Apply the Kalman filter and a disturbance smoother (Koopman, 1993) and obtain the posterior mean of $\boldsymbol{\eta}_{t:t+k}$ ($\mathbf{h}_{t:t+k}$) and set $\hat{\boldsymbol{\eta}}_{t:t+k}$ ($\hat{\mathbf{h}}_{t:t+k}$) to this value.
5. Return to step 2 and repeat the procedure until achieving convergence.

Finally, we describe the updating procedure for h_1 and the knot conditions h_{k_l} , for $l = 1, \dots, K$. We simulate $h_1 | h_2, \boldsymbol{\theta} \sim \mathcal{N}(\mu + \varphi(h_1 - \mu), \sigma_\eta^2)$. As the density $p(h_{k_l} | h_{k_l-1}, h_{k_l+1})$ does not have a closed form, we use the Metropolis-Hastings algorithm with proposal $\mathcal{N}(\frac{\mu(1-\varphi)^2 + \varphi(h_{k_l-1} + h_{k_l+1})}{1+\varphi^2}, \frac{\sigma_\eta^2}{1+\varphi^2})$. Let $h_{k_l}^p$ and $h_{k_l}^{(i-1)}$ denote the proposal value and the previous iteration value. Thus, the acceptance probability is given by $\alpha_{MH} = \min\{1, \frac{Q(h_{k_l}^p)}{Q(h_{k_l}^{(i-1)})}\}$, where $Q(h_{k_l})$ is the conditional density of $y_{k_l} | \boldsymbol{\theta}, W_{k_l}, U_{k_l}, h_{k_l}$.




Figure 2: Compounded NASDAQ returns from January 7, 1988 to July 3, 2003. The left panel shows the plot of the raw series and the right panel the histogram of returns.

4 Empirical Application

This section analyzes the daily closing prices of the NASDAQ Composite index. The NASDAQ Composite is a stock market index of the common stocks and similar securities listed on the NASDAQ stock market, meaning that it has over 3000 components. It is highly followed in the U.S. as an indicator of the performance of stocks of technology and growth companies. Since both U.S. and non-U.S. companies are listed on the NASDAQ stock market, the index is not exclusively a U.S. index. The data set was obtained from the Yahoo finance web site, available to download at <http://finance.yahoo.com>. The period of analysis is July 5, 1988 - July 3, 2003, which yields 3784 observations. Throughout, we work with the compounded return expressed as a percentage, $y_t = 100 \times (\log P_t - \log P_{t-1})$, where P_t is the closing price on day t . The compounded NASDAQ index returns are plotted in Figure 2 as a time series plot and also as a histogram. We clearly identify the period of elevated volatility around of the turn of the Millennium associated with the collapse of the Tech bubble. We are particularly interested in understanding the importance of excess of kurtosis and skewness in the NASDAQ index return and

we avoid confounding our results by this highly influential outlier. There are some interesting pattern we observe in this history of NASDAQ returns. The average return is 0.04 percent daily with a daily standard deviation of 1.54. Returns exhibit negative skewness of -0.02 and kurtosis of 9.13. Note also that the returns have a large range (minimum, -10.16 and maximum, 13.25). We use the the Jarque-Bera (JB) statistics to test the normality assumption of the returns. In light of the JB statistics (5923.6), the null hypothesis of normality is rejected (p-value of 0.00) due to negative skewness and excess of kurtosis.

Table 1: Estimation results for the NASDAQ returns. First row: Posterior mean. Second row: Posterior 95% credible interval in parentheses. Third row: CD statistics.

Parameter	SV-N	SV-T	SV-SN	SV-ST
	0.0623	0.0457	0.0417	0.0314
μ	(-0.9523,0.9764)	(-1.3032,1.2315)	(-1.1524,1.0409)	(-1.5076,1.3985)
	0.70	0.94	-0.91	-0.66
	0.9944	0.9963	0.9954	0.9967
φ	(0.9897,0.9984)	(0.9926,0.9994)	(0.9745,0.9947)	(0.9932,0.9995)
	1.12	-0.43	0.11	-1.58
	0.0172	0.0118	0.0143	0.0107
σ_{η}^2	(0.0115,0.0246)	(0.0075,0.0171)	(0.0097,0.0204)	(0.0070,0.0155)
	-0.13	0.39	0.81	0.87
	-	-	-1.3908	-1.1528
λ	-	-	(-1.6280,-1.1470)	(-1.4020,-0.8820)
	-	-	0.23	-1.65
	-	19.3369	-	19.6797
ν	-	(11.3700,35.3600)	-	(11.4000,36.5000)
	-	-1.78	-	1.72
	-	0.0564	-	0.0556
$\frac{1}{\nu}$	-	(0.0283,0.0879)	-	(0.0274,0.0877)
	-	1.51	-	-1.36

Now, we analyze the NASDAQ index returns with the aim of providing robust inference. In our analysis, we fit and compare the SV-N, SV-T, SV-SN and SV-ST models. In all cases, we simulated the h_t 's in a multi-move fashion with stochastic knots based on the method described in Section 3.2. We fix the number of blocks K to be 95 in such a way that each block contained 40 h_t 's on average. We set the prior distribution of the common parameters as: $\varphi \sim \mathcal{N}_{(-1,1)}(0.95, 100)$, $\sigma^2 \sim \mathcal{IG}(2.5, 0.025)$, $\mu \sim \mathcal{N}(0, 100)$. For the parameter φ the priors' mean and variance are 0.0032 and 0.3328, respectively. This prior setup is equivalent to the uniform distribution on interval $(-1, 1)$, which gives zero mean and variance of 0.3333. We assume that $\lambda \sim t_{0.5}(0.0, \frac{\pi^2}{4})$, a Jeffreys' prior suggested by Bayes and Branco (2007). Finally, for ν , we assume the prior given by equation (10). All the calculations were performed running stand-alone code developed by us using an open source C++ library for statistical computation, the Scythe statistical library (Pemstein *et al.*, 2007), which is available for free download at <http://scythe.wustl.edu>.

For all models, we conducted the MCMC simulation for 50000 iterations. In all cases, the first 10000 draws were discarded as a burn-in period. In order to reduce the autocorrelation between successive values of the simulated chain, only every 20th values of the chain were stored. With the resulting 2000 values, we calculated the posterior means, the 95% credible intervals and the convergence diagnostic (CD) statistics (Geweke, 1992). If the sequence of the recorded MCMC output is stationary, it converges in distribution to the standard normal. According to the CD the null hypothesis that the sequence of 2000 draws is stationary was accepted at the 5% level, $CD \in (-1.96, 1.96)$, for all the parameters in all the models considered here. Table 1 summarizes the results.

From Table 1, consistent with the existing evidence of great persistence in the log-volatility process, we found that the posterior means of φ and 95% posterior credible intervals very close to the unity. Being the posterior mean of φ of the SV-ST model slightly higher than those of the other three models. The posterior mean of σ_η^2 is smaller in the SV-ST than those of the SV-N, SV-T and the SV-SN models, indicating that the log-volatility process of the SV-ST is less variable than those of the other ones.

In the SV-T and SV-ST models, the magnitude of the tail-fatness is measured by the degrees of freedom, ν , parameter. We found that the posterior mean of ν are the SV-T are 19.34 and 19.68, respectively, which indicates tail-fatness. In Table 1, we report the posterior mean of $1/\nu$, for both models, which, in both cases, are over 3.5 standard deviation from zero. Since the SV-N and SV-SN models are nested in the limit when $1/\nu$ approaches to zero this provides strong evidence of heavy-tailness of conditional distributions of the returns.

Regarding the skewness parameter, λ , in the SV-SN and SV-ST models, we found that the posterior means are -1.3908 and -1.1528, respectively. In both models, the 95% credible interval does not contain zero, that is the negativity of λ is credible. This supports the strong evidence of skewnesses in the NASDAQ data set.

The magnitudes of the mixing parameter U_t are associated with extremeness of the corresponding observations. In the Bayesian paradigm, the posterior mean of the mixing parameter can be used to identify a possible outlier (see, for instance, Rosa *et al.*, 2003). The SV-T and SV-ST models can accommodate an outlier by inflating the variance component for that observation in the conditional distribution with smaller U_t value. This fact is shown in Figure 3 where we depicted the posterior mean of the mixing variable U_t for the SV-T (top panel) and SV-ST (bottom panel) models, respectively.

To assess the goodness of the estimated models, we calculate the Bayesian predictive information criteria, BPIC (Ando, 2006; 2007). The BPIC criterion is defined as

$$BPIC = -2E_{\boldsymbol{\theta}|\mathbf{y}_{1:T}}[\log\{p(\mathbf{y}_{1:T} | \boldsymbol{\theta})\}] + 2T\hat{b}, \quad (16)$$

where \hat{b} is given by

$$\hat{b} \approx \frac{1}{T} \left\{ E_{\boldsymbol{\theta}|\mathbf{y}_{1:T}}[\log\{p(\mathbf{y}_{1:T} | \boldsymbol{\theta})p(\boldsymbol{\theta})\}] - \log[p(\mathbf{y}_{1:T} | \hat{\boldsymbol{\theta}})p(\hat{\boldsymbol{\theta}})] + \text{tr}\{J_T^{-1}(\hat{\boldsymbol{\theta}})I_T(\hat{\boldsymbol{\theta}})\} + 0.5q \right\}. \quad (17)$$

Here q is the dimension of $\boldsymbol{\theta}$, $E_{\boldsymbol{\theta}|\mathbf{y}_{1:T}}[\cdot]$ denotes the expectation with respect to the posterior distribution,





Figure 3: NASDAQ data set: posterior smoothed mean of mixture variable U_t for the SV-T (top panel) and SV-ST (bottom panel) models.

$\hat{\boldsymbol{\theta}}$ is the posterior mode, and

$$I_T(\hat{\boldsymbol{\theta}}) = \frac{1}{T} \sum_{t=1}^T \left(\frac{\partial \eta_T(y_t, \boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \frac{\partial \eta_T(y_t, \boldsymbol{\theta})}{\partial \boldsymbol{\theta}'} \right) \Big|_{\boldsymbol{\theta}=\hat{\boldsymbol{\theta}}},$$

$$J_T(\hat{\boldsymbol{\theta}}) = \frac{1}{T} \sum_{t=1}^T \left(\frac{\partial^2 \eta_T(y_t, \boldsymbol{\theta})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}'} \right) \Big|_{\boldsymbol{\theta}=\hat{\boldsymbol{\theta}}},$$

with $\eta_T(y_t, \boldsymbol{\theta}) = \log p(y_t | \mathbf{y}_{1:t-1}, \boldsymbol{\theta}) + \log p(\boldsymbol{\theta})/T$.

In the SV-N, SV-T, SV-SN, and SV-ST models, the log-likelihood function, $\log p(\mathbf{y}_{1:T} | \boldsymbol{\theta})$, is estimated using the auxiliary particle filter (see, e.g., [Pitt and Shephard, 1999](#)) with 10000 particles. From [Table 2](#), the BPIC criterion indicates the SV-ST model is the best model among all the models considered here, suggesting that the NASDAQ index return data demonstrate sufficient departure from underlying normality assumptions and asymmetry.

In [Figure 4](#), we plot the smoothed mean of $e^{\frac{h_t}{2}}$ obtained from the MCMC output for the SV-N (solid line) and the SV-ST (dotted line). From a practical point of view, we are mainly interested in whether we find a significant difference between the two series. Therefore, in the bottom panel of [Figure 9](#), we plot the smoothed mean of the difference of $e^{\frac{h_t}{2}}$ obtained from the SV-N and SV-T models. Some extreme returns make the differences clear. This can have a substantial impact, for instance, in the valuation of derivative instruments and several strategic or tactical asset allocation topics.

5 Conclusions

In this article, we presented a Bayesian implementation of the stochastic volatility model with skew-Student-t (SV-ST) errors as an alternative to the normal assumption of the conditional distribution of





Figure 4: NASDAQ data set. Top: Posterior smoothed mean of $e^{\frac{h_t}{2}}$. SV-N (solid line) and SV-ST (dotted line). Bottom: Posterior smoothed mean of the difference of $e^{\frac{h_t}{2}}$ in both models.

Table 2: Nasdaq return data set. BPIC: Bayesian predictive information criterion.

Model	BPIC	Ranking
SV-N	15628.4	3
SV-T	11662.6	2
SV-SN	16012.4	4
SV-ST	11575.3	1

the returns. The SV-ST model allows a parsimonious yet flexible treatment of both skewness and tail thickness. Under a Bayesian perspective, we developed a fast and efficient MCMC sampling procedure to estimate all the parameters and latent quantities in our proposed SV-ST model. We use objective priors for the degrees of freedom and the skewness parameters, ν and λ , based on [Fonseca *et al.* \(2008\)](#) and [Bayes and Branco \(2007\)](#), respectively. As a by product of the MCMC algorithm, we were able to produce an estimate of the latent information process which can be used in financial modelling. The use of mixing variable, $\mathbf{U}_{1:T}$ not only simplifies the full conditional distributions required for the Gibbs sampling algorithm, but also provides a mean for outlier diagnostics. We illustrated our methods through an empirical application of the NASDAQ return series, which showed that the SV-ST model provides better fit than the SV-N, SV-T and SV-SN models in terms of parameter estimates, interpretation and robustness aspects. On the other hand, since the posterior mean and 95% posterior credibility interval of the parameter λ contains only negative values, we can conclude that there is a strong evidence of skewness in the NASDAQ data set.

Acknowledgements

The research of Carlos A. Abanto-Valle was supported by CNPq. V. H. Lachos acknowledges financial support from FAPESP and CNPq.

Appendix A: The full conditionals

In this appendix, we describe the full conditional distributions for the parameters and the mixing latent variables $\mathbf{U}_{1:T}$ and $\mathbf{W}_{1:T}$ for the SV-ST model model.

Full conditional distribution of μ , φ and σ_η^2

The prior distributions of the common parameters are set as: $\mu \sim N(\bar{\mu}, \sigma_\mu^2)$, $\varphi \sim \mathcal{N}_{(-1,1)}(\bar{\varphi}, \sigma_\varphi^2)$, $\sigma_\eta^2 \sim \mathcal{IG}(\frac{T_0}{2}, \frac{M_0}{2})$. We have the following full conditional for μ :

$$\mu \mid \mathbf{h}_{1:T}, \varphi, \sigma_\eta^2 \sim \mathcal{N}\left(\frac{b_\mu}{a_\mu}, \frac{1}{a_\mu}\right), \quad (\text{A.1})$$

where $a_\mu = \frac{1}{\sigma_\alpha^2} + \frac{(T-1)(1-\varphi)^2}{\sigma_\eta^2} + \frac{(1-\varphi)^2}{\sigma_\eta^2}$ and $b_\mu = \frac{\bar{\mu}}{\sigma_\mu^2} + \frac{(1-\varphi^2)}{\sigma_\eta^2} h_1 + \frac{\sum_{t=1}^{T-1} (h_{t+1} - \varphi h_t)(1-\varphi)}{\sigma_\eta^2}$. In a similar way, the conditional posterior of φ is given by

$$p(\varphi \mid \mathbf{h}_{1:T}, \mu, \sigma_\eta^2) \propto Q(\varphi) \exp\left\{-\frac{a_\varphi}{2} \left(\psi - \frac{b_\varphi}{a_\varphi}\right)^2\right\} \mathbb{I}_{|\varphi| < 1}, \quad (\text{A.2})$$

where $Q_\varphi = \sqrt{1-\varphi^2} \exp\left\{-\frac{1}{2\sigma_\eta^2} [(1-\varphi^2)(h_1 - \mu)^2]\right\}$, $a_\varphi = \frac{\sum_{t=1}^{T-1} (h_t - \mu)^2}{\sigma_\eta^2} + \frac{1}{\sigma_\psi^2}$, $b_\varphi = \frac{\sum_{t=1}^{T-1} (h_t - \mu)(h_{t+1} - \mu)}{\sigma_\eta^2} + \frac{\bar{\varphi}}{\sigma_\varphi^2}$ and $\mathbb{I}_{|\varphi| < 1}$ is an indicator variable. As $p(\varphi \mid \mathbf{h}_{0:T}, \alpha, \sigma_\eta^2)$ in (A.2) does not have closed form, we sample from it by using the Metropolis-Hastings algorithm with truncated $\mathcal{N}_{(-1,1)}(\frac{b_\psi}{a_\psi}, \frac{1}{a_\psi})$ as the proposal density.

Finally, the full conditional of σ_η^2 is $\mathcal{IG}(\frac{T_1}{2}, \frac{M_1}{2})$, where $T_1 = T_0 + T$ and $M_1 = M_0 + [(1 - \psi^2)(h_1 - \mu)^2] + \sum_{t=1}^{T-1} [h_{t+1} - \mu - \psi(h_t - \mu)]^2$.

Full conditional of ν , λ , U_t and W_t

We, set ζ and ω in such a way that $E(y_t \mid h_t) = 0$ and $V(y_t \mid h_t) = e^{h_t}$. So, we have $\zeta = -\sqrt{\frac{2}{\pi}} k_1 \delta \omega$ and $\omega^2 = \left[k_2 - \frac{2}{\pi} k_1^2 \delta^2 \right]^{-1}$, where $k_1 = \sqrt{\frac{\nu}{2}} \frac{\Gamma(\frac{\nu-1}{2})}{\Gamma(\frac{\nu}{2})}$, $k_2 = \frac{\nu}{\nu-2}$ and $\delta = \frac{\lambda}{\sqrt{1+\lambda^2}}$. Then the full the conditionals of

ν and λ follows:

$$p(\nu | \cdot) \propto \left(\frac{\nu}{\nu+3} \right)^{\frac{1}{2}} \left\{ \psi' \left(\frac{\nu}{2} \right) - \psi' \left(\frac{\nu+1}{2} \right) - \frac{2(\nu+3)}{\nu(\nu+1)^2} \right\}^{\frac{1}{2}} \\ \times \left(\frac{1}{\omega} \right)^T e^{-\frac{1}{2\omega^2(1-\delta^2)} \sum_{t=1}^T U_t e^{-h_t} (y_t - \zeta - \omega \delta W_t U_t^{-\frac{1}{2}} e^{\frac{h_t}{2}})^2}, \quad (\text{A.3})$$

$$p(\lambda | \cdot) \propto \left(1 + \frac{2\lambda}{\pi^2} \right)^{-\frac{3}{4}} \left(\frac{1}{1-\delta^2} \right)^{\frac{T}{2}} e^{-\frac{1}{2\omega^2(1-\delta^2)} \sum_{t=1}^T U_t e^{-h_t} (y_t - \zeta - \omega \delta W_t U_t^{-\frac{1}{2}} e^{\frac{h_t}{2}})^2}. \quad (\text{A.4})$$

Since the above full conditional distributions are not in any known closed form, we must simulate ν and λ using the Metropolis-Hastings algorithm. The proposal density used are $\mathcal{N}(\nu > 2)(\mu_\nu, \tau_\nu^2)$ and $\mathcal{N}(\mu_\lambda, \tau_\lambda^2)$, with $\mu_\nu = x - \frac{q'(x)}{q''(x)}$ and $\tau_\nu^2 = \max\{0.001, (-q''(x))^{-1}\}$ for $\nu = \nu$ or λ , where x is the value of the previous iteration, $q(\cdot)$ is the logarithm of the conditional posterior density, and $q'(\cdot)$ and $q''(\cdot)$ are the first and second derivatives respectively.

As $U_t \sim \mathcal{G}(\frac{\nu}{2}, \frac{\nu}{2})$, the conditional posterior of U_t is given by

$$p(U_t | h_t, W_t, \nu, \lambda) \propto Q(U_t) U_t^{\frac{\nu+1}{2}-1} e^{-\frac{U_t}{2} [\nu + \frac{e^{-h_t} (y_t - \zeta e^{\frac{h_t}{2}})^2}{\omega^2(1-\delta^2)}]}, \quad (\text{A.5})$$

where $Q(U_t) = e^{\frac{U_t^{\frac{1}{2}} \delta W_t e^{-\frac{h_t}{2}} (y_t - \zeta e^{\frac{h_t}{2}})}{\omega(1-\delta^2)}}$. As $p(U_t | h_t, W_t, \nu, \lambda)$ in (A.5) does not have closed form, we sample from it by using the Metropolis-Hastings algorithm with $\mathcal{G}(\frac{\nu+1}{2}, \frac{1}{2} [\nu + \frac{e^{-h_t} (y_t - \zeta e^{\frac{h_t}{2}})^2}{\omega^2(1-\delta^2)}])$ as the proposal density. Finally, from equations (8a) and (8c), we have the full conditional of W_t is the $\mathcal{N}_{[0, \infty)}(\frac{\delta U_t^{-\frac{1}{2}} e^{-\frac{h_t}{2}} [y_t - \zeta e^{\frac{h_t}{2}}]}{\omega}, \frac{1}{1-\delta^2})$.

References

- Abanto-Valle, C. A. Bandyopadhyay, D. Lachos, V. H. and Enriquez, I. (2010) Robust Bayesian Analysis of Heavy-tailed Stochastic Volatility Models using Scale Mixtures of Normal Distributions. *Computational Statistics & Data Analysis*, 54, 2883–2898.
- Ando, T. (2006) Bayesian Inference for nonlinear and non-gaussian stochastic volatility model with leverage effect. *Journal of Japan Statistical Society*, 36, 173–197.

- Ando, T. (2007) Bayesian predictive information criterion for the evaluation of hierarchical Bayesian and empirical Bayes models. *Biometrika*, 94, 443–458.
- Azzalini, A. (1986) Further results on a class of distributions which include the normal ones. *Statistica*, 46, 199–208.
- Azzalini, A. (2005) The Skew-normal distribution and related multivariate families. *Scandinavian Journal of Statistics*, 32, 159–188.
- Azzalini, A. and Capitanio, A. (2003) Distributions generated by perturbation of symmetry with emphasis on a multivariate skew-t distribution. *Journal of Royal Statistical Society: Series B*, 65, 367–389.
- Bayes, C. L. and Branco, M. D. (2007) Bayesian inference for the skewness parameter of the scalar skew-normal distribution. *Brazilian Journal of Probability and Statistics*, 21, 141–163.
- Branco, M. D. and Dey, D. K. (2001) A general class of multivariate skew-elliptical distributions. *Journal of Multivariate Analysis*, 79, 99–113.
- Cappuccio, N. Lubian, D. and Raggi, D. (2006) Investigating asymmetry in U.S. stock market indexes: evidence from a stochastic volatility model. *Applied Financial Economics*, 16, 479–490.
- Chib, S. and Greenberg, E. (1995) Understanding the Metropolis-Hastings algorithm. *The American Statistician*, 49, 327–335.
- Chib, S. Nardari, F. and Shepard, N. (2002) Markov Chain Monte Carlo methods for stochastic volatility models. *Journal of Econometrics*, 108, 281–316.
- Corrado, C. J. and Su, T. (1997) Implied volatility skews and stock index skewness and kurtosis implied by S&P500 index option prices. Working paper University of Missouri - Columbia.

- de Jong, P. and Shephard, N. (1995) The Simulation Smoother for Time Series Models. *Biometrika*, 82, 339–350.
- Fama, E. (1965) Portfolio analysis in a stable paretian market. *Managament Science*, 11, 404–419.
- Fonseca, T. C. O. Ferreira, M. A. R. and Migon, H. S. (2008) Objective Bayesian analysis for the Student-t regression model. *Biometrika*, 95, 325–333.
- Geweke, J. (1992) Evaluating the accuracy of sampling-based approaches to the calculation of posterior moments. In: *Bayesian Statistics. Vol. 4.*, eds. J. M. Bernardo J. O. Berger A. P. Dawid and A. F. M. Smith Oxford, U.K.: Oxford University Press pp. 169–193.
- Henze, N. (1986) A probabilistic representation of the skew-normal distribution. *Scandinavian Journal of Statistics*, 13, 271–275.
- Jacquier, E. Polson, N. and Rossi, P. (2004) Bayesian analysis of stochastic volatility models with Fat-tails and correlated errors. *Journal of Econometrics*, 122, 185–212.
- Koopman, S. (1993) Disturbance smoothers for State Space models. *Biometrika*, 80, 117–126.
- Lange, K. L. and Sinsheimer, J. S. (1993) Normal/independent distributions and their applications in robust regression. *J. Comput. Graph. Stat*, 2, 175–198.
- Mandelbrot, B. (1963) The variation of certain speculative prices. *Journal of Business*, 36, 314–419.
- Mittnik, S. and Paoella, M. S. (2000) Conditional density and value-at-risk prediction of asian currency exchange rates. *Journal of Forecasting*, 19, 313–333.
- Peiro, A. (1999) Skewness in financial returns. *Journal of Banking and Finance*, 23, 847–862.
- Pemstein, D. Quinn, K. V. and Martin, A. D. (2007) The Scythe statistical library: An open source C++ library for statistical computation. *Journal of Statistical Software*, V, 1–29.

- Pitt, M. and Shephard, N. (1999) Filtering via simulation: Auxiliary particle filter. *Journal of the American Statistical Association*, 94, 590–599.
- Rosa, G. J. M. Padovani, C. R. and Gianola, D. (2003) Robust linear mixed models with Normal/Independent distributions and Bayesian MCMC implementation. *Biometrical Journal*, 45, 573–590.
- Shephard, N. and Pitt, M. (1997) Likelihood analysis of non-Gaussian measurements time series. *Biometrika*, 84, 653–667.
- Taylor, S. (1982) Financial returns modelled by the product of two stochastic processes—a study of the daily sugar prices 1961–75. In: *Time Series Analysis: Theory and Practice, Vol 1*, (eds O. Anderson) Amsterdam: North-Holland pp. 203–226.
- Taylor, S. (1986) *Modeling Financial Time Series* Chichester: Wiley.
- Tierney, L. (1994) Markov chains for exploring posterior distributions (with discussion). *Annals of Statistics*, 21, 1701–1762.
- Watanabe, T. and Omori, Y. (2004) A multi-move sampler for estimate non-Gaussian time series model: Comments on Shepard and Pitt (1997). *Biometrika*, 91, 246–248.