

# AVERAGING METHODS FOR STUDYING THE PERIODIC ORBITS OF DISCONTINUOUS DIFFERENTIAL SYSTEMS

JAUME LLIBRE<sup>1</sup>, DOUGLAS D. NOVAES<sup>2</sup> AND MARCO ANTONIO TEIXEIRA<sup>2</sup>

ABSTRACT. The main objective of this work is to extend the averaging method for studying the periodic orbits of a class of differential equations with discontinuous second member. Thus, overall results are presented to ensure the existence of limit cycles of such systems. Certainly these results represent new insights in averaging, in particular its relation with non smooth dynamical systems theory. An application is presented in careful detail.

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

The discontinuous differential systems, i.e. differential equations with discontinuous right-hand sides, is a subject that has been developed very fast these last years. It has become certainly one of the common frontiers between Mathematics, Physics and Engineering. Thus certain phenomena in control systems, impact in mechanical systems and nonlinear oscillations are the main sources of motivation of their study, see for more details Teixeira [21].

The knowledge of the existence or not of periodic solutions is very important for understanding the dynamics of the differential systems. One of good tools for study the periodic solutions is the averaging theory, see for instance the books of Sanders and Verhulst [19] and Verhulst [23]. We point out that the method of averaging is a classical and matured tool that provides a useful means to study the behaviour of nonlinear smooth dynamical systems. The method of averaging has a long history that starts with the classical works of Lagrange and Laplace who provided an intuitive justification of the process. The first formalization of this procedure was given by Fatou in 1928 [8]. Very important practical and theoretical contributions in the averaging theory were made by Krylov and Bogoliubov [2] in the 1930s and Bogoliubov [1] in 1945. The principle of averaging has been extended in many directions for both finite- and infinite-dimensional differentiable systems. The classical results for studying the periodic orbits of differential systems need at least that those systems be of class  $C^2$ . Recently Buica and Llibre [4] extended the averaging theory for studying periodic orbits to continuous differential systems using mainly the Brouwer degree.

The main objective of this paper is to extend the averaging theory for studying periodic orbits to discontinuous differential systems using again the Brouwer degree.

Let  $D$  be an open subset of  $\mathbb{R}^n$ . We shall denote the points of  $\mathbb{R} \times D$  as  $(t, x)$ , and we shall call the variable  $t$  as the time. Let  $h : \mathbb{R} \times D \rightarrow \mathbb{R}$  be a  $C^1$  function having the  $0 \in \mathbb{R}$  as a regular value, and let  $\mathcal{M} = h^{-1}(0)$ .

---

2010 *Mathematics Subject Classification.* 37G15, 37C80, 37C30.

*Key words and phrases.* periodic solution, averaging theory, discontinuous differential system.

Let  $X, Y : \mathbb{R} \times D \rightarrow \mathbb{R}^n$  be two continuous vector fields. Assume that the functions  $h, X$  and  $Y$  are  $T$ -periodic in the variable  $t$ . Now we define a *discontinuous piecewise differential system*

$$(1) \quad x'(t) = Z(t, x) = \begin{cases} X(t, x) & \text{if } h(t, x) > 0, \\ Y(t, x) & \text{if } h(t, x) < 0. \end{cases}$$

Denote  $Z = (X, Y)$ .

Here we deal with a different formulation for the discontinuous differential system (1). Let  $\text{sign}(u)$  be the sign function defined in  $\mathbb{R} \setminus \{0\}$  as

$$\text{sign}(u) = \begin{cases} 1 & \text{if } u > 0, \\ -1 & \text{if } u < 0. \end{cases}$$

Then the discontinuous differential system (1) can be written using the function  $\text{sign}(u)$  as

$$(2) \quad x'(t) = Z(t, x) = F_1(t, x) + \text{sign}(h(t, x))F_2(t, x),$$

where

$$F_1(t, x) = \frac{1}{2}(X(t, x) + Y(t, x)) \quad \text{and} \quad F_2(t, x) = \frac{1}{2}(X(t, x) - Y(t, x)).$$

To work with the discontinuous differential system (2) we should introduce the regularization process, where the discontinuous vector field  $Z(t, x)$  is approximated by an one-parameter family of continuous vector fields  $Z_\delta(t, x)$  such that  $\lim_{\delta \rightarrow 0} Z_\delta = Z(t, x)$ .

In [20] Sotomayor and Teixeira introduced a regularization for the discontinuous vector fields in  $\mathbb{R}^2$  having a line of discontinuity and, using this technique, they proved generically that its regularization provides the same extension of the orbits through the line of discontinuity that the one given by the Filippov's rules, see [8]. Later on Llibre and Teixeira [18] studied the regularization of generic discontinuous vector fields in  $\mathbb{R}^3$  having a surface of discontinuity, and proved that  $\lim_{\delta \rightarrow 0} Z_\delta$  essentially agrees with Filippov's convention in dimension three. Finally, in [21] Teixeira generalized the regularization procedure to finite dimensional discontinuous vector fields.

In [16] Llibre, da Silva and Teixeira studied singular perturbations problems in dimension three which are approximations of discontinuous vector fields proving that the regularization process developed in [18] produces a singular problem for which the discontinuous set is a center manifold, moreover, they proved that the definition of sliding vector field coincides with the reduced problem of the corresponding singular problem for a class of vector fields.

In all these regularizations (except the one using singular perturbation theory) a transition function is used to average the vector fields  $X$  and  $Y$  on the set of discontinuity in order to get a family of continuous vector fields that approximates the discontinuous one.

A continuous function  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  is a *transition function* if  $\phi(u) = -1$  for  $u \leq -1$ ,  $\phi(u) = 1$  for  $u \geq 1$  and  $\phi'(u) > 0$  if  $u \in (-1, 1)$ . The  $\phi$ -regularization of  $Z = (X, Y)$  is the one-parameter family of continuous functions  $Z_\delta$  with  $\delta \in (0, 1]$

given by

$$Z_\delta(t, x) = \frac{1}{2}(X(t, x) + Y(t, x)) + \frac{1}{2}\phi_\delta(h(t, x))(X(t, x) - Y(t, x)),$$

with

$$(3) \quad \phi_\delta(u) = \phi\left(\frac{u}{\delta}\right).$$

Note that for all  $(t, x) \in (\mathbb{R} \times D) \setminus \mathcal{M}$  we have that  $\lim_{\delta \rightarrow 0} Z_\delta(t, x) = Z(t, x)$ .

The formulation (2) of the discontinuous differential system (1) admits a natural regularization. Define the transition function  $\phi$  as

$$(4) \quad \phi(u) = \begin{cases} 1 & \text{if } u \geq 1, \\ u & \text{if } -1 < u < 1, \\ -1 & \text{if } u \leq -1. \end{cases}$$

Let  $\phi_\delta : \mathbb{R} \rightarrow \mathbb{R}$  be the continuous function defined in (3). It is clear that

$$(5) \quad \lim_{\delta \rightarrow 0} \phi_\delta(u) = \text{sign}(u),$$

and

$$Z_\delta(t, z) = F_1(t, x) + \phi_\delta(h(t, x))F_2(t, x),$$

is the  $\phi$ -regularization of the discontinuous differential system (1).

If  $\Sigma = \{0\} \times D$ , then doing a time translation (if necessary) we always can assume that  $\Sigma \not\subseteq \mathcal{M}$ . As usual  $\nabla h$  denotes the gradient of the function  $h$ , and  $\nabla_x h$  denotes the gradient of the function  $h$  restricted to the variable  $x$ .

Our main result on the periodic orbits of the discontinuous differential system 2 using averaging is given in the next theorem. Its proof uses the theory of Brouwer degree for finite dimensional spaces (see the appendix A for a definition of the Brouwer degree  $d_B(f, V, 0)$ ), and is based on the averaging theory for non-smooth differential system stated by Buica and Llibre [4] (see Appendix B).

**Theorem 1.** *We consider the following discontinuous differential system*

$$(6) \quad x'(t) = \varepsilon F(t, x) + \varepsilon^2 R(t, x, \varepsilon),$$

with

$$\begin{aligned} F(t, x) &= F_1(t, x) + \text{sign}(h(t, x))F_2(t, x), \\ R(t, x, \varepsilon) &= R_1(t, x, \varepsilon) + \text{sign}(h(t, x))R_2(t, x, \varepsilon), \end{aligned}$$

where  $F_1, F_2 : \mathbb{R} \times D \rightarrow \mathbb{R}^n$ ,  $R_1, R_2 : \mathbb{R} \times D \times (-\varepsilon_0, \varepsilon_0) \rightarrow \mathbb{R}^n$  and  $h : \mathbb{R} \times D \rightarrow \mathbb{R}$  are continuous functions,  $T$ -periodic in the variable  $t$  and  $D$  is an open subset of  $\mathbb{R}^n$ . We also suppose that  $h$  is a  $C^1$  function having 0 as a regular value. Denote by  $\mathcal{M} = h^{-1}(0)$ , by  $\Sigma = \{0\} \times D \not\subseteq \mathcal{M}$ , by  $\Sigma_0 = \Sigma \setminus \mathcal{M} \neq \emptyset$ , and its elements by  $z \equiv (0, z) \notin \mathcal{M}$ .

Define the averaged function  $f : D \rightarrow \mathbb{R}^n$  as

$$(7) \quad f(x) = \int_0^T F(t, x) dt.$$

We assume the following three conditions.

- (i)  $F_1, F_2, R_1, R_2$  and  $h$  are locally  $L$ -Lipschitz with respect to  $x$ ;
- (ii) for  $a \in \Sigma_0$  with  $f(a) = 0$ , there exist a neighbourhood  $V$  of  $a$  such that  $f(z) \neq 0$  for all  $z \in \overline{V} \setminus \{a\}$  and  $d_B(f, V, 0) \neq 0$ .

- (iii) If  $\partial h/\partial t \neq 0$ , then for all  $(t, z) \in \mathcal{M}$  we have that  $(\partial h/\partial t)(t, z) \neq 0$ ; and if  $\partial h/\partial t \equiv 0$ , then  $\langle \nabla_x h, F_1 \rangle^2 - \langle \nabla_x h, F_2 \rangle^2 > 0$  for all  $(t, z) \in [0, T] \times \mathcal{M}$ .

Then, for  $|\varepsilon| > 0$  sufficiently small, there exists a  $T$ -periodic solution  $x(\cdot, \varepsilon)$  of system (6) such that  $x(0, \varepsilon) \rightarrow a$  as  $\varepsilon \rightarrow 0$ .

Theorem 1 is proved in section 2.

We remark that condition (iii) of Theorem 1 changes according with the fact that the hypersurface of discontinuity  $\mathcal{M}$  depends or not on the time .

The averaging theory has already been used in discontinuous dynamical systems. Thus, for instance, Llibre, Novaes and Teixeira [13] have used the averaging theory to provide sufficient conditions for the existence of periodic solutions of the planar double pendulum with discontinuous perturbation. Llibre and Teixeira [17] have used the averaging theory to provide lower bounds for the maximum number of limit cycles for the  $m$ -piecewise discontinuous polynomial differential equations. In [5] Cardin, Carvalho and Llibre have used the averaging theory to study the bifurcation of limit cycles from the periodic orbits of two and four dimensional linear center in  $\mathbb{R}^n$  perturbed inside in a class of discontinuous piecewise linear differential systems. Finally, in [15] Llibre and Rong have used the averaging theory to study the number of limit cycles of the discontinuous piecewise linear differential systems in  $\mathbb{R}^{2n}$  with two zones separated by a hyperplane.

In Theorem 1 we have extended to general discontinuous differential systems the ideas used in the previous mentioned papers for particular discontinuous differential systems.

Now we shall do an application of Theorem 1 to the discontinuous piecewise linear differential systems separated by a straight line. These differential systems have been studied recently by Han and Zhang [10], and Huan and Yuang [11], among other papers. In [10] some results about the existence of two limit cycles appeared, so that the authors conjectured that the maximum number of limit cycles for this class of piecewise linear differential systems is exactly two. This conjecture is analogous to Conjecture 1 in the discussion of Tonnelier in [22]. However, by considering a specific family of discontinuous PWL differential systems with two linear zones sharing the equilibrium position, in [11] strong numerical evidence about the existence of three limit cycles was obtained, and a proof was provided by Llibre and Ponce [14]. This example represents up to now the first discontinuous piecewise linear differential system with two zones and 3 limit cycles surrounding a unique equilibrium. Now we shall provide a new proof of the existence of these three limit cycles using the averaging theory through our Theorem 1.

In polar coordinates  $(r, \theta)$  given by  $x = r \cos \theta$  and  $y = r \sin \theta$ , the planar discontinuous piecewise linear differential system with two zones separated by a straight line corresponding to the system studied in the paper [14] is

$$(8) \quad \frac{dr}{d\theta} = F(\theta, r) = \begin{cases} \varepsilon \frac{19}{50} r & \text{if } r \cos \theta \geq 1, \\ \varepsilon \frac{2300 \cos(2\theta) - 4623 \sin(2\theta) - 300}{1500} r & \text{if } r \cos \theta < 1, \end{cases}$$

where we have multiplied the right hand side of the system of [14] by the small parameter  $\varepsilon$ . Our main contributions in this application is to provide the explicit

analytic equations defining the limit cycles of the discontinuous piecewise linear differential system with two zones (8).

**Theorem 2.** *Any limit cycle of the discontinuous piecewise linear differential system with two zones (8) which intersects the straight line  $x = 1$  in two points  $(r_0, \theta_0)$  and  $(r_1, \theta_1)$  with  $-\pi/2 < \theta_0 < 0 < \theta_1 < \pi/2$ ,  $r_k \cos \theta_k = 1$  for  $k = 0, 1$ , and  $r_0 > 1$  and  $\theta_1$  must satisfy the following two equations*

$$\begin{aligned}
 & \exp\left(\frac{19(\theta_1 - \theta_0)}{50}\right) r_0 \cos \theta_1 - 1 = 0, \\
 & \frac{19(\theta_1 - \theta_0)}{50} + \frac{1}{5} \arctan\left(\frac{1}{15} \sec \theta_0 (23 \cos \theta_0 - 100 \sin \theta_0)\right) \\
 (9) \quad & - \frac{1}{5} \arctan\left(\frac{1}{15} \sec \theta_1 (23 \cos \theta_1 - 100 \sin \theta_1)\right) \\
 & - \frac{1}{2} \log(|4623 \cos(2\theta_0) + 2300 \sin(2\theta_0) - 5377|) \\
 & + \frac{1}{2} \log(|4623 \cos(2\theta_1) + 2300 \sin(2\theta_1) - 5377|) - \frac{2\pi}{5} = 0,
 \end{aligned}$$

where  $\theta_0 = \arccos(1/r_0) - \pi$ , and the determination of the arctan is in the interval  $(-\pi/2, \pi/2)$  and of the arccos in the interval  $(0, \pi)$ .

On the other hand, any limit cycle of system (8) which intersects the straight line  $x = 1$  in two points  $(r_0, \theta_0)$  and  $(r_1, \theta_1)$  with  $0 < \theta_0 < \theta_1 < \pi/2$ ,  $r_k \cos \theta_k = 1$  for  $k = 0, 1$ , and  $r_0 > 1$  and  $\theta_1$  must satisfy the equations (9), but now in both equations  $\theta_0 = \arccos(1/r_0)$ .

We recall that a *limit cycle* of system (8) is an isolated periodic orbit of that system in the set of all periodic orbits of the system. It is well known that the study of the limit cycles of the differential systems in dimension two is one of the main problems of the qualitative theory of differential systems in dimension two, see for instance the surveys of Ilyashenko [9] and Jibin Li [12].

In fact, as we shall see in the proof of Theorem 2 equations (9) have three solutions, providing the three limit cycles of Figure 1.

## 2. PROOF OF THEOREM 1

For proving Theorem 1 we need some preliminary lemmas. In what follows we always assume that all hypotheses of Theorem 1 hold.

**Lemma 3.** *If  $\partial h / \partial t \not\equiv 0$ , then for  $z \in D$  and  $t \in [0, T]$  the map  $h_z : t \mapsto h(t, z)$  has finitely many zeros in  $[0, T]$ .*

*Proof.* From the definition of  $h$  we know that the map  $h_z : t \mapsto h(t, z)$  is  $C^1$ . Suppose that there exist an increasing sequence  $(t_i)_{i \in \mathbb{N}} \subset [0, T]$  such that  $h_z(t_i) = 0$  for all  $i \in \mathbb{N}$ . Then there exist a convergent subsequence  $(t_{i_j})_{j \in \mathbb{N}}$  such that  $t_{i_j} \rightarrow \bar{t} \in [0, T]$ . By continuity of  $h_z$ , it follows that  $h_z(\bar{t}) = 0$ , which implies that  $(\bar{t}, z) \in \mathcal{M}$ . We know, by the Mean Value Theorem, that for each  $j \in \mathbb{N}$  there exist  $s_j \in (t_{i_j}, t_{i_{j+1}})$  such that  $h'_z(s_j) = 0$ . Since  $s_j \rightarrow \bar{t}$ , then  $h'_z(\bar{t}) = 0$ , which is a contradiction with hypothesis (iii). Hence, the map  $h_z : t \mapsto h(t, z)$ , for  $t \in [0, T]$ , vanishes only in a finite subset  $\mathcal{T} \subset [0, T]$ .  $\square$




FIGURE 1. The three limit cycles surrounding the origin.

In the hypothesis (ii) of Theorem 1 we use the Brouwer degree of the averaged function  $f$ . We need to show that the function  $f$  is continuous in order that the Brouwer degree will be well defined, for more details see Appendix A.

**Lemma 4.** *The averaged function (7) is continuous in  $\Sigma_0$ .*

*Proof.* Fix  $z_0 \in \Sigma_0$ . Since the map  $h_{z_0} : t \mapsto h(t, z_0)$  is  $C^1$  and  $(\mathbb{R} \times D) \setminus \mathcal{M}$  is an open subset of  $\mathbb{R} \times D$ . If  $\partial h / \partial t \not\equiv 0$ , by Lemma 3, for  $t \in [0, T]$ , we know that  $h(t, z_0) = 0$  only for a finite subset  $\mathcal{T} = \{t_1, t_2, \dots, t_{k-1}\}$  of  $[0, T]$ . If  $\partial h / \partial t \equiv 0$ , then  $\mathcal{T} = \emptyset$ . Set  $t_0 = 0$  and  $t_k = T$ . Now  $h(t, z_0)h(s, z_0) > 0$  for  $t, s \in (t_i, t_{i+1})$  and  $i = 0, 1, \dots, k-1$ , i.e.  $h(t, z_0)$  does not change its sign in  $(t_i, t_{i+1})$  for  $i = 0, 1, \dots, k-1$ . Now, for  $z \in D$  in some neighborhood of  $z_0$ , we estimate

$$\begin{aligned}
 |f(z) - f(z_0)| &\leq \int_0^T |F_1(t, z_0) - F_1(t, z)| dt \\
 &\quad + \int_0^t |\text{sign}(h(t, z_0))F_2(t, z_0) - \text{sign}(h(t, z))F_2(t, z)| dt \\
 &\leq TL|z_0 - z| + \int_0^t \tilde{F}(t, z_0, z) dt \\
 &\leq TL|z_0 - z| + \sum_{i=0}^{k-1} \int_{t_i}^{t_{i+1}} \tilde{F}(t, z_0, z) dt,
 \end{aligned}$$

where  $\tilde{F}(t, z_0, z) = |\text{sign}(h(t, z_0))F_2(t, z_0) - \text{sign}(h(t, z))F_2(t, z)|$ . Set  $I_i = (t_i, t_{i+1})$  and  $I_i^\gamma = [t_i + \gamma, t_{i+1} - \gamma]$  for  $i = 1, 2, \dots, k-1$ . Note that there exists  $\gamma_0 > 0$  such that  $I_i^\gamma \subset I_i$  for all  $\gamma \leq \gamma_0$ . The set  $I_i^\gamma \times \{z_0\} \subset S$  is compact, then there exist a

ball  $B(z_0, r_\gamma)$  such that  $K_i^\gamma = I_i^\gamma \times \overline{B(z_0, r_\gamma)} \subset S$ . We can assume that  $r_\gamma < \gamma$ . Set  $M_i^\gamma = \sup\{\tilde{F}(t, z_0, z) : (t, z) \in K_i^\gamma\}$ . Note that

$$\begin{aligned} \tilde{F}(t, z_0, z) &\leq |\text{sign}(h(t, z_0))F_2(t, z_0)| + |\text{sign}(h(t, z))F_2(t, z)| \\ &= |F_2(t, z_0)| + |F_2(t, z)|, \end{aligned}$$

if we take  $M_0 = \sup\{|F_2(t, z_0)| + |F_2(t, z)| : (t, z) \in [0, T] \times \overline{B(z_0, r_\gamma)}\}$ , then  $M_i^\gamma \leq M_0$  for all  $i = 1, 2, \dots, k-1$ . Thus

$$\begin{aligned} \int_{t_i}^{t_{i+1}} \tilde{F}(t, z_0, z) &= \int_{I_i^\gamma} \tilde{F}(t, z_0, z) + \int_{t_i}^{t_i+\gamma} \tilde{F}(t, z_0, z) + \int_{t_{i+1}-\gamma}^{t_{i+1}} \tilde{F}(t, z_0, z) \\ &\leq TL|z_0 - z| + 2\gamma M_i^\gamma \leq (TL + 2M_0)\gamma. \end{aligned}$$

Hence

$$|f(z_0) - f(z)| \leq TL\gamma + k(TL + 2M_0)\gamma \leq C_0\gamma.$$

Therefore, given  $\varepsilon > 0$  we can choose  $\gamma = \varepsilon/C_0$ , then for all  $z \in B(z_0, r_\gamma)$  we have  $|f(z_0) - f(z)| < \varepsilon$ . This implies that  $f$  is continuous in  $z_0 \in \Sigma_0$ .  $\square$

Let  $a$  be the point in hypothesis (ii) of Theorem 1. By Lemma 4 there exists a neighborhood  $V$  of  $a$  such that  $f$  is continuous in  $V$ . Hence, by Theorems 9 and 10 (see Appendix A), there exists a unique map that satisfies the properties of the Brouwer degree for the function  $f(z)$  with  $z \in \overline{V}$ , because  $0 \notin f(\partial V)$ . This map is denoted by  $d_B(f, V, 0)$ .

**Lemma 5.** *For  $|\varepsilon|$  sufficiently small the solutions of system (6), in the sense of Filippov, are uniquely defined in any compact set  $[0, T] \times D_1$  contained in  $[0, T] \times D$ .*

To prove Lemma 5 we will need the following proposition, that has been proved in Corollary 1 of section 10 of chapter 1 of [8]. Define

$$\begin{aligned} S^+ &= \{(t, x) \in \mathbb{R} \times D : h(x, t) > 0\}, \\ S^- &= \{(t, x) \in \mathbb{R} \times D : h(x, t) < 0\}. \end{aligned}$$

Note that  $\mathbb{R} \times D = S^- \cup \mathcal{M} \cup S^+$ .

**Proposition 6.** *For every point of the manifold  $\mathcal{M}$  where  $(Xh)(Yh) > 0$ , there is a unique solution passing either from  $S^-$  into  $S^+$ , or from  $S^+$  into  $S^-$ .*

*Proof of Lemma 5.* System (6) can be written as the autonomous system

$$\begin{pmatrix} \tau' \\ x' \end{pmatrix} = \begin{cases} X(\tau, x) & \text{if } h(\tau, x) > 0, \\ Y(\tau, x) & \text{if } h(\tau, x) < 0, \end{cases}$$

in  $\mathbb{R} \times D$ , where

$$\begin{aligned} X(\tau, x) &= \begin{pmatrix} 1 \\ \varepsilon(F_1(\tau, x) + F_2(\tau, x)) + \varepsilon^2(R_1(\tau, x, \varepsilon) + R_2(\tau, x, \varepsilon)) \end{pmatrix}, \\ Y(\tau, x) &= \begin{pmatrix} 1 \\ \varepsilon(F_1(\tau, x) - F_2(\tau, x)) + \varepsilon^2(R_1(\tau, x, \varepsilon) - R_2(\tau, x, \varepsilon)) \end{pmatrix}. \end{aligned}$$

If  $\partial h/\partial t \neq 0$ , then we have that

$$(Xh)(Yh) = \langle \nabla h, X \rangle \langle \nabla h, Y \rangle = \left( \frac{\partial h}{\partial t} \right)^2 (t, x) + \mathcal{O}(\varepsilon).$$

By hypotheses (iii) we know that if  $(t, x) \in \mathcal{M}$ , then  $(\partial h/\partial t)(t, x) \neq 0$ . Hence, for  $|\varepsilon|$  sufficiently small, we conclude that  $(Xh)(Yh)(t, x) > 0$ , for every  $(t, x)$  in any compact set  $[0, T] \times D_1$  with  $D_1 \subset D$ . Applying Proposition 6 the result follows.

If we assume that  $\partial h/\partial t \equiv 0$ . Then

$$(Xh)(Yh) = \langle \nabla h, X \rangle \langle \nabla h, Y \rangle = \varepsilon^2 (\langle \nabla_x h, F_1 \rangle^2 - \langle \nabla_x h, F_2 \rangle^2) + \mathcal{O}(\varepsilon^3).$$

By hypotheses (iii) for  $|\varepsilon|$  sufficiently small we conclude that  $(Xh)(Yh)(t, x) > 0$  for every  $(t, x)$  in any compact set  $[0, T] \times D_1$  with  $D_1 \subset D$ . Again applying Proposition 6 the lemma is proved.  $\square$

Instead of working with the discontinuous differential system (6) we shall work with the continuous differential system

$$(10) \quad x'(t) = \varepsilon F_\delta(t, x) + \varepsilon^2 R_\delta(t, x, \varepsilon),$$

where

$$\begin{aligned} F_\delta(t, x) &= F_1(t, x) + \phi_\delta(h(t, x))F_2(t, x), \\ R_\delta(t, x, \varepsilon) &= R_1(t, x, \varepsilon) + \phi_\delta(h(t, x))R_2(t, x, \varepsilon), \end{aligned}$$

and  $\phi_\delta : \mathbb{R} \rightarrow \mathbb{R}$  is the continuous function defined in (3) and (4), and satisfying (5).

For system (10) the averaged function is defined as

$$f_\delta(z) = \int_0^T F_\delta(t, x) dt.$$

We need to guarantee that hypothesis (i) of Theorem 11 (see appendix A) holds for the functions  $F_\delta$  and  $R_\delta$ . For this purpose we prove the following lemma.

**Lemma 7.** *For  $\delta \in (0, 1]$  the function  $\phi_\delta : \mathbb{R} \rightarrow \mathbb{R}$  defined in (3) with  $\phi$  given by (4) is globally  $1/\delta$ -Lipschitz; i.e. for all  $u_1, u_2 \in \mathbb{R}$  we have that  $|\phi_\delta(u_1) - \phi_\delta(u_2)| \leq (1/\delta)|u_1 - u_2|$ .*

*Proof.* If  $u_1 \leq -\delta < \delta \leq u_2$ , then  $|\phi_\delta(u_1) - \phi_\delta(u_2)| = 2 = (1/\delta)2\delta \leq (1/\delta)|u_1 - u_2|$ .

If  $u_1, u_2 \leq -\delta$  or  $u_1, u_2 \geq \delta$ , then  $|\phi_\delta(u_1) - \phi_\delta(u_2)| = 0 \leq (1/\delta)|u_1 - u_2|$ .

Assume that  $u_1 \in (-\delta, \delta)$ . If  $|u_2| < \delta$ , then  $|\phi_\delta(u_1) - \phi_\delta(u_2)| = (1/\delta)|u_1 - u_2|$ ; and if  $|u_2| \geq \delta$ , then  $|\phi_\delta(u_1) - \phi_\delta(u_2)| \leq \max\{|1/\delta|, |1/u_2|\}|u_1 - u_2| \leq (1/\delta)|u_1 - u_2|$ . This completes the proof of the lemma.  $\square$

**Proposition 8.** *For  $\delta \in (0, 1]$  the functions  $F_\delta$  and  $R_\delta$  are locally Lipschitz with respect to the variable  $x$ .*

*Proof.* Let  $K \subset D$  be a compact subset. Denote  $M = \sup\{|F_2(t, x)| : (t, x) \in [0, T] \times K\}$ , which is well defined by continuity of the function  $(t, x) \mapsto |F_2(t, x)|$  and compactness of the set  $[0, T] \times K$ . For  $x_1$  and  $x_2$  in  $K$  where  $F_1$  and  $h$  are

locally  $L$ -Lipschitz and by Lemma 7, we have

$$\begin{aligned}
|F_\delta(t, x_1) - F_\delta(t, x_2)| &= |F_1(t, x_1) - F_1(t, x_2) \\
&\quad + \phi_\delta(h(t, x_1))F_2(t, x_1) - \phi_\delta(h(t, x_2))F_2(t, x_2)| \\
&\leq |F_1(t, x_1) - F_1(t, x_2)| \\
&\quad + |\phi_\delta(h(t, x_1))F_2(t, x_1) - \phi_\delta(h(t, x_2))F_2(t, x_2)| \\
&\leq L|x_1 - x_2| + |\phi_\delta(h(t, x_1))||F_2(t, x_1) - F_2(t, x_2)| \\
&\quad + |F_2(t, x_2)||\phi_\delta(h(t, x_1)) - \phi_\delta(h(t, x_2))| \\
&\leq 2L|x_1 - x_2| + \frac{M}{\delta}|h(t, x_1) - h(t, x_2)| \\
&\leq \left(2L + \frac{ML}{\delta}\right)|x_1 - x_2| = L_\delta|x_1 - x_2|.
\end{aligned}$$

The proof for  $R_\delta$  is analogous.  $\square$

Now we are ready to prove Theorem 1.

*Proof of Theorem 1.* We will study the Poincaré maps for the discontinuous differential system (6) and for the continuous differential system (10). For each  $z \in D$ , let  $x_\delta(t, z, \varepsilon)$  denote the solution of system (10) such that  $x_\delta(0, z, \varepsilon) = z$ .

Since the differential system (10) is  $T$ -periodic in the variable  $t$ , we can consider system (10) as a differential system defined on the generalized cylinder  $\mathbb{S}^1 \times D$  obtained by identifying  $\Sigma = \{(\tau, x) : \tau = 0\}$  with  $\{(\tau, x) : \tau = T\}$ , see Figure 2. On this cylinder  $\Sigma$  is a section for the flow. Moreover, if  $z \in D$  is the coordinate of a point on  $\Sigma$ , then we consider the Poincaré map  $P_\delta^\varepsilon(z) = x_\delta(T, z, \varepsilon)$  for the points  $z$  such that  $x_\delta(T, z, \varepsilon)$  is defined.

Observe that there exists  $\varepsilon_0 > 0$  such that, whenever  $\varepsilon \in [-\varepsilon_0, \varepsilon_0]$ , the solution  $x_\delta(t, z, \varepsilon)$  is uniquely defined on the interval  $[0, T]$ . Indeed, if  $(t_z^-, t_z^+)$  is the maximal open interval for which the solution passing through  $(0, z)$  is defined. Now we shall apply the local existence and uniqueness theorem for the solutions of these differential, see for example Theorem 1.2.2 of [19]. Note that we can apply that theorem due to the result of Proposition 8. Hence, by the local existence and uniqueness theorem we have that  $t_z^+ > h_z$  and  $h_z = \inf\{T, d \setminus m(\varepsilon)\}$  where  $m(\varepsilon) \geq |\varepsilon F_\delta(t, x) + \varepsilon^2 R_\delta(t, x, \varepsilon)|$  for all  $t \in [0, T]$ , for each  $x$  with  $|x - z| \leq d$  and for every  $z \in D$ . When  $|\varepsilon|$  is sufficiently small,  $m(\varepsilon)$  can be arbitrarily large, in such a way that  $h_z = T$  for all  $z \in D$ . Hence, for  $\varepsilon \in [-\varepsilon_0, \varepsilon_0]$ , the Poincaré map of system (10) is well defined and continuous for every  $z \in D$ .

From the definition of the Poincaré map  $P_\delta^\varepsilon(z)$  its fixed points correspond to periodic orbits of period  $T$  of the differential system (10) defined on the cylinder.

We can define in a similar way the Poincaré map  $P^\varepsilon(z)$  of the discontinuous differential system (6). More precisely, by Lemma 5 and by the continuous dependence of solution of the differential equation with discontinuous righthand side on the initial data (see for instance Corollary 1 of section 8 of chapter 2 of [8]), such Poincaré map is continuous. Again the fixed points of  $P^\varepsilon(z)$  correspond to periodic orbits of the discontinuous differential system (6).

By definition the continuous differential system (10) is  $C^2$  in the variable  $\varepsilon$ . So we do the Taylor expansion of the Poincaré map of system (10) around  $\varepsilon$  up to




FIGURE 2. Generalized cylinder.

order two, and we get

$$(11) \quad P_\delta^\varepsilon(z) = z + \varepsilon f_\delta(z) + \mathcal{O}(\varepsilon^2),$$

where  $f_\delta(z)$  is the averaged function of the continuous differential system (10), for more details see for instance [4].

The referred Poincaré map is the composition of the Poincaré maps of the continuous differential systems, so it is continuous. Therefore, we can repeat the arguments of [4] and we again have that

$$(12) \quad P^\varepsilon(z) = z + \varepsilon f(z) + \mathcal{O}(\varepsilon^2),$$

where  $f(z)$  is the averaged function of the discontinuous differential system (6).

Due to (5) the pointwise limit of the Poincaré map  $P_\delta^\varepsilon(z)$  of system (10), when  $\delta \rightarrow 0$  is the Poincaré map  $P^\varepsilon(z)$  of system (6). Consequently, from (11) and (12) we obtain that the pointwise limit in  $\Sigma$  of the function  $f_\delta$ , when  $\delta \rightarrow 0$  is the function  $f$ .

Let  $a \in V$  be the point satisfying hypotheses (ii) of Theorem 1. Therefore  $f(z) \neq 0$  for all  $z \in \overline{V} \setminus \{a\}$ . Define  $f_0 = f|_{\overline{V}}$ , we know that  $f_0$  is continuous by Lemma 4. Then, we consider the continuous homotopy  $\{f_\delta|_{\overline{V}}, 0 \leq \delta \leq 1\}$ . We claim that there exists a  $\delta_0 \in (0, 1]$  such that  $0 \notin f_\delta(\partial V)$  for all  $\delta \in [0, \delta_0]$ . Now we shall prove the claim.

As usual  $\mathbb{N}$  denotes the set of positive integers. Suppose that there exists a sequence  $(z_m)_{m \in \mathbb{N}}$  in  $\partial V$  such that  $f_{\frac{1}{m}}(z_m) = 0$ . As the sequence  $(z_m)$  is contained in the compact set  $\partial V$ , so there exists a subsequence  $(z_{m_\ell})_{\ell \in \mathbb{N}}$  such  $z_{m_\ell} \rightarrow z_0 \in \partial V$ . Consequently we obtain that  $f(z_0) = 0$ , in contradiction with the hypotheses (ii) of Theorem 1. Hence, the claim is proved.

From the above claim and the property (iii) of Theorem 9 (see Appendix A) we conclude that  $d_B(f_\delta, V, 0) \neq 0$  for  $0 \leq \delta \leq \delta_0$ . Therefore, by the property (i) of Theorem 9 we obtain that  $0 \in f_\delta(V)$ , so there exists  $a_\delta \in V$  such that  $f_\delta(a_\delta) = 0$ .

Since, by continuity, there exists the  $\lim_{\delta \rightarrow 0} a_\delta$  and it is a zero of the function  $f_0 = f|_V$ . This limit is the point  $a$  of the hypotheses (ii) of Theorem 1, because  $a$  is the unique zero of  $f_0$  in  $V$ .

In summary, in order that for every  $\delta \in (0, \delta_0]$  the averaged function  $f_\delta$  satisfy the assumptions (ii) of Theorem 11 (see Appendix B). So it only remains to show that in  $V$  we have that  $f_\delta(z) \neq 0$  for all  $z \in \overline{V} \setminus \{a_\delta\}$ . But this can be achieved in a complete similar way as we proved the above claim. Hence, by Proposition 8 for every  $\delta \in (0, \delta_0]$  the continuous differential system (10) satisfies all the assumptions of Theorem 11. Hence, for  $|\varepsilon|$  sufficiently small there exists a periodic solution  $x_\delta(t, \varepsilon)$  of the continuous differential system (10) such that  $z_{(\delta, \varepsilon)} := x_\delta(0, \varepsilon) \rightarrow a_\delta$  when  $\varepsilon \rightarrow 0$ .

Now, from (11) the point  $z_{(\delta, \varepsilon)}$  is a fixed point of the Poincaré map  $P_\delta^\varepsilon(z)$ , i.e.  $P_\delta^\varepsilon(z_{(\delta, \varepsilon)}) = z_{(\delta, \varepsilon)}$ . Since  $\lim_{\delta \rightarrow 0} P_\delta^\varepsilon(z) = P^\varepsilon(z)$ , it follows that  $z_\varepsilon = \lim_{\delta \rightarrow 0} z_{(\delta, \varepsilon)}$  is a fixed point of the Poincaré map  $P^\varepsilon(z)$ . So, the discontinuous differential system (6) has a periodic solution  $x(t, \varepsilon)$  such that  $z_\varepsilon = x(0, \varepsilon) \rightarrow a$  as  $\varepsilon \rightarrow 0$ . Therefore the theorem is proved.  $\square$

### 3. APPLICATION

In this section we shall prove Theorem 2, by applying Theorem 1 to the discontinuous differential system (8). So, we must compute the integral (7), which for system (8) becomes

$$(13) \quad f(r) = \int_0^{2\pi} F(\theta, r) d\theta,$$

where the function  $F(\theta, r)$  is given in (8).

The solution of the differential system (8) in the half-plane  $x = r \cos \theta \geq 1$  starting at the point  $(r_0, \theta_0)$  with  $r_0 \cos \theta_0 = 1$  and  $\theta_0 \in (-\pi/2, 0)$  is

$$r(\theta) = \exp\left(\frac{19(\theta - \theta_0)}{50}\right) r_0.$$

Therefore, at the point  $(r_1, \theta_1)$  with  $r_1 \cos \theta_1 = 1$  and  $\theta_1 \in (0, \pi/2)$  we have that

$$\exp\left(\frac{19(\theta_1 - \theta_0)}{50}\right) r_0 \cos \theta_1 = 1.$$

This equation coincides with the first equation of (9).

Now computing the integral (13) we obtain exactly the right hand side of the second equation of (9) multiplied by  $r$ . According to Theorem 1 we must find the zeros of this last expression. Since  $r$  cannot be zero the equation for the zeros is reduced exactly to the second equation of (9). In short, by Theorem 1 we have proved that a periodic orbit of system (8) intersects the straight line  $x = 1$  in two points  $(r_0, \theta_0)$  and  $(r_1, \theta_1)$  with  $\theta_0 \in (-\pi/2, 0)$ ,  $\theta_1 \in (0, \pi/2)$ ,  $r_k \cos \theta_k = 1$  for  $k = 0, 1$ , and  $r_0 > 1$  and  $\theta_1$  must satisfy the equations (9).

In [14] it is proved that the discontinuous differential equation (8) has three limit cycles (i) and that their points  $(r_0, \theta_0)$  and  $(r_1, \theta_1)$  are approximately for the inner limit cycle of Figure 1

$$(14) \quad r_0 = 1.013330663139\dots, \quad \theta_0 = 0.162383740477\dots, \quad \theta_1 = 0.5541676264624\dots;$$

for the middle limit cycle of Figure 1

$$(15) \quad r_0 = 1.003945075086\dots, \quad \theta_0 = -0.088680876377\dots, \quad \theta_1 = 0.768002346543\dots;$$

for the external limit cycle of Figure 1

$$(16) \quad r_0 = 1.111870463116\dots, \quad \theta_0 = -0.452434880837\dots, \quad \theta_1 = 1.034197922817\dots$$

It is easy to check that (14), (15) and (16) satisfies the two equations (9). Hence, Theorem 2 is proved.

#### APPENDIX A: BASIC RESULTS ON THE BROUWER DEGREE

In this appendix we present the existence and uniqueness result from the degree theory in finite dimensional spaces. We follow the Browder's paper [3], where are formalized the properties of the classical Brouwer degree.

**Theorem 9.** *Let  $X = \mathbb{R}^n = Y$  for a given positive integer  $n$ . For bounded open subsets  $V$  of  $X$ , consider continuous mappings  $f : \bar{V} \rightarrow Y$ , and points  $y_0$  in  $Y$  such that  $y_0$  does not lie in  $f(\partial V)$  (as usual  $\partial V$  denotes the boundary of  $V$ ). Then to each such triple  $(f, V, y_0)$ , there corresponds an integer  $d(f, V, y_0)$  having the following three properties.*

- (i) *If  $d(f, V, y_0) \neq 0$ , then  $y_0 \in f(V)$ . If  $f_0$  is the identity map of  $X$  onto  $Y$ , then for every bounded open set  $V$  and  $y_0 \in V$ , we have*

$$d(f_0|_V, V, y_0) = \pm 1.$$

- (ii) *(Additivity) If  $f : \bar{V} \rightarrow Y$  is a continuous map with  $V$  a bounded open set in  $X$ , and  $V_1$  and  $V_2$  are a pair of disjoint open subsets of  $V$  such that*

$$y_0 \notin f(\bar{V} \setminus (V_1 \cup V_2)),$$

*then,*

$$d(f_0, V, y_0) = d(f_0, V_1, y_0) + d(f_0, V_2, y_0).$$

- (iii) *(Invariance under homotopy) Let  $V$  be a bounded open set in  $X$ , and consider a continuous homotopy  $\{f_t : 0 \leq t \leq 1\}$  of maps of  $\bar{V}$  in to  $Y$ . Let  $\{y_t : 0 \leq t \leq 1\}$  be a continuous curve in  $Y$  such that  $y_t \notin f_t(\partial V)$  for any  $t \in [0, 1]$ . Then  $d(f_t, V, y_t)$  is constant in  $t$  on  $[0, 1]$ .*

**Theorem 10.** *The degree function  $d(f, V, y_0)$  is uniquely determined by the three conditions of Theorem 9.*

For the proofs of Theorems 9 and 10 see [3].

#### APPENDIX B: BASIC RESULTS ON AVERAGING THEORY

In this appendix we present the basic result from the averaging theory that we shall need for proving the main results of this paper. For a general introduction to averaging theory see for instance the book of Sanders and Verhulst [19].

**Theorem 11.** *We consider the following differential system*

$$(17) \quad x'(t) = \varepsilon F(t, x) + \varepsilon^2 R(t, x, \varepsilon),$$

where  $F : \mathbb{R} \times D \rightarrow \mathbb{R}^n$  and  $R : \mathbb{R} \times U \times (-\varepsilon_f, \varepsilon_f) \rightarrow \mathbb{R}^n$  are continuous functions,  $T$ -periodic in the first variable and  $D$  is an open subset of  $\mathbb{R}^n$ . We define the averaged function  $f : D \rightarrow \mathbb{R}^n$  as

$$(18) \quad f(x) = \int_0^T F(s, x) ds,$$

and assume that

- (i)  $F$  and  $R$  are locally Lipschitz with respect to  $x$ ;
- (ii) for  $a \in D$  with  $f(a) = 0$ , there exist a neighborhood  $V$  of  $a$  such that  $f(z) \neq 0$  for all  $z \in \overline{V} \setminus \{a\}$  and  $d_B(f, V, 0) \neq 0$ .

Then, for  $|\varepsilon| > 0$  sufficiently small, there exist a  $T$ -periodic solution  $x(t, \varepsilon)$  of the system (17) such that  $x(0, \varepsilon) \rightarrow a$  as  $\varepsilon \rightarrow 0$ .

Theorem 11 for studying the periodic orbits of continuous differential systems has weaker hypotheses than the classical result for studying the periodic orbits of smooth differential systems, see for instance Theorem 11.5 of Verhulst [23], where instead of (i) is assumed that

- (j)  $F$ ,  $R$ ,  $D_x F$ ,  $D_x^2 F$  and  $D_x R$  are defined, continuous and bounded by a constant  $M$  (independent of  $\varepsilon$ ) in  $[0, \infty) \times D$ ,  $-\varepsilon_f < \varepsilon < \varepsilon_f$ ;

and instead of (ii) it is required that

- (jj) for  $a \in D$  with  $f(a) = 0$  we have that  $J_f(a) \neq 0$ , where  $J_f(a)$  is the Jacobian matrix of the function  $f$  at the point  $a$ .

For a proof of Theorem 11 see [4] section 3.

#### ACKNOWLEDGEMENTS

The first author is partially supported by a MICINN/FEDER grant MTM2008-03437, by a AGAUR grant number 2009SGR-0410 and by ICREA Academia. The second author is partially supported by the grant FAPESP 2011/03896-0 The third author is partially supported by a FAPESP-BRAZIL grant 2007/06896-5. The first and third authors are also supported by the joint project CAPES-MECD grant PHB-2009-0025-PC

#### REFERENCES

- [1] N. N. BOGOLIUBOV, *On some statistical methods in mathematical physics*, Izv. vo Akad. Nauk Ukr. SSR, Kiev, 1945.
- [2] N. N. BOGOLIUBOV AND N. KRYLOV, *The application of methods of nonlinear mechanics in the theory of stationary oscillations*, Publ. 8 of the Ukrainian Acad. Sci. Kiev, 1934.
- [3] F. BROWDER, *Fixed point theory and nonlinear problems*, Bull. Amer. Math. Soc. **9** (1983), 1-39.
- [4] A. BUICA AND J. LLIBRE, *Averaging methods for finding periodic orbits via Brouwer degree*, Bulletin des Sciences Mathématiques **128** (2004), 7-22.
- [5] P.T. CARDIN, T. CARVALHO AND J. LLIBRE, *Limit Cycles of discontinuous piecewise linear differential systems*, Int. J. Bifurcation and Chaos **21** (2011), 3181-3194.
- [6] C. C. CHICONE, *Ordinary Differential Equations With Applications*, Texts in Applied Mathematics, **34**, Springer, 1999.
- [7] P. FATOU, *Sur le mouvement d'un système soumis à des forces à courte période*, Bull. Soc. Math. France **56** (1928), 98-139.
- [8] A. F. FILIPPOV, *Differential Equations with Discontinuous Righthand Side*, Mathematics and Its Applications, Kluwer Academic Publishers, Dordrecht, 1988.
- [9] YU. ILYASHENKO, *Centennial history of Hilbert's 16th problem*, Bull. Amer. Math. Soc. **39** (2002), 301-354.

- [10] M. HAN AND W. ZHANG, *On Hopf bifurcation in non-smooth planar systems*, J. of Differential Equations **248** (2010), 2399–2416.
- [11] S.M. HUAN AND X.S. YANG, *The number of limit cycles in general planar piecewise linear systems*, to appear in Discrete and Continuous Dynamical Systems-A, 2011.
- [12] J. LI, *Hilbert's 16th problem and bifurcations of planar polynomial vector fields*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. **13** (2003), 47–106.
- [13] J. LLIBRE, D.D. NOVAES AND M.A. TEIXEIRA, *On the periodic solutions of a generalized smooth and non-smooth perturbed planar double pendulum with small oscillations*, arXiv:1203.0498v1 [math.DS]
- [14] J. LLIBRE AND E. PONCE, *Three limit cycles in discontinuous piecewise linear differential systems with two zones*, to appear in Dynamics of Continuous, Discrete and Impulsive Systems, Serie B.
- [15] J. LLIBRE AND F. RONG, *On the number of limit cycles for discontinuous piecewise linear differential systems in  $\mathbb{R}^{2n}$  with two zones*, to appear in Int. J. of Bifurcation and Chaos.
- [16] J. LLIBRE, P.R. DA SILVA AND M.A. TEIXEIRA, *Regularization of discontinuous vector fields on  $\mathbb{R}^3$  via singular perturbation*, J. Dynamics and Differential Equations **19** (2007), 309–331.
- [17] J. LLIBRE AND M.A. TEIXEIRA, *Limit cycles for  $m$ -piecewise discontinuous polynomial Liénard differential equations*, arXiv: [math.DS]
- [18] J. LLIBRE AND M.A. TEIXEIRA, *Regularization of discontinuous vector fields in dimension three*, Discrete and Continuous Dynamical Systems **3** (1997), 235–241.
- [19] J. SANDERS AND F. VERHULST, *Averaging method in nonlinear dynamical systems*, Applied Mathematical Sciences **59**, Springer, 1985.
- [20] J. SOTOMAYOR AND M.A. TEIXEIRA, *Regularization of Discontinuous Vector Field*, International Conference on Differential Equation, Lisboa, 1995, World Sci. Publ., River Edge, NJ, 1998, pp 207–223.
- [21] M.A. TEIXEIRA, *Perturbation theory for non-smooth systems*, Encyclopedia of Complexity and Systems Science **22**, Springer, New York, 2009, pp 6697–6719.
- [22] A. TONNELIER, *The McKean's caricature of the FitzHugh-Nagumo model I. The space-clamped system*, SIAM J. Appl. Math. **63** (2003), pp. 459484.
- [23] F. VERHULST, *Nonlinear Differential Equations and Dynamical Systems*, Universitext, Springer, 1991.

<sup>1</sup> DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA, BARCELONA, CATALONIA, SPAIN  
*E-mail address:* jllibre@mat.uab.cat

<sup>2</sup> DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE ESTADUAL DE CAMPINAS, CAIXA POSTAL 6065, 13083–859, CAMPINAS, SP, BRAZIL  
*E-mail address:* ddnovaes@gmail.com, teixeira@ime.unicamp.br